



Trichonema subpalustre Herb. (sine
diagnosi) in Edwards Bot. Reg. 1847 (XXXIII ap.
totius vel ~~et~~ nov. ser.) t 40, 1. (Supra copia iconis.)

Descriptio reperitur in:

Revisioe Iridearum. Auct. Dr. F. W.

Wlatt. (Conclusio). — Linnaea XXXIV, (XVIII)
1865-66. p. 669.

16. *Trichonema subpalustre* W. Herbert.

T. scapo tereti multifloro erecto demum refle.
xo folioso, foliis filiformi-ensiformibus cur-
vatis acutis striatis scapum superanti-
bus basi longe vaginantibus, spathis bi-
valvibus, valvis lanceolatis acutis, purpu-
reo-punctatis; perigonii tubo brevi, lacini-
is lanceolatis acutis striatis albis

marginibus caeruleis, stigmatibus papillo-
sis. — Ic. Bot. Reg. ~~XXXIII~~. t. 40, f. 1.

Hab. Morea l. Bona St. Vincent (in Herb.
r. Berol.), ad ~~Thessalonicam~~ Thessalonicam,
et in insulis Ionicis, Herbert.

Scapus 2-3" altus; folia $\frac{1}{2}$ " lata. Spa-
thae 6" longae. Perigonii lacinae
10" longae, 2" latae.