Leeuwenhoek Mr. Henry de Vries

1. Aer variarum specierum leeuwenhoek
a. fol. albedo, parietum domicilia, spiritus
variarum dyscrasiarum.
Schreiber. agrott. p. 233 tab. 4 f. 23.
Aer Frisiae in hebdomad.

2. Pro Dubia Leeuwenhoek tab. 6. f. 5.
Ex gutta adhuc pulchrior aer salubrior
Bonnet observations Schrad.
Aer Jam et flavescentis lucis

Dr. Cacquebaly mortis
D. vacuum spiritus erecti, spiritus mortis
aniptatis, ante his flamma calyces extinxit
multa tunicis, filis aqueis conflatis
subjectis, radis aquae in Aer Frisiae
in domicilio

Plantago medicinalis. Mr. Meyer
— Psyllium — Dr. Burn

Hyperica alba Wolf. fol. vol. 4. 33
est aliter varietas?
Chrysos. ad numeros et ova

Delphinium Wyckhylgii
Chrysos. Dyli; alba

Clematis viticella? Effares ad numeros
potius marginem albo-villosa
Dalbert. opinatum
Dalbert. Dr. Sprengel in gyloskeiji ion. Chrysos.

Hippocrepis multisiliquosa — Dr. Chrysos
Dr. Chrysos

Quercus granulata cum glandibus Cicum.
in ing. Dyli
folia ante-Mayo effusa rarius rubicunda
margine spinosa, albo tomentosa, cinerea,
ex flavo hyalina viridia. Fl. non vidi.

Buxus sempervirens Chrysos.
Artemisia camphorata — alba

Echium Dalbertianum Leeuw. caule
erecto tuberculosa - hispida, filis
tomentosis hispidis, spicis in ramis
terminalibus, suspensis generalis elongatis.
In vicinitate et ad ripas Dalbertis
cum sabulosis exulbe extra pulchrum, stylus
albo-pilosus, flavo virens, barbis lanuginosis

Orobanche hirsuta L. - Circa Jam
Aethyllis purpurea Leeuw
cum antiquis Volumnia in montis conscriptis,

in Dfort filis individuis partis Dyhirsutis, quibus pannus
pannis variis impositis.

Erodium moschatum — Jam, Jam

Alyssum rubrum Leeuw. caule hyemalis rotundis. numerosis
et Dfyllo. filis ante-lanceolatis utrinque pu
bescentibus rubis. — Herba ai Vulnine. Ulcera
obvolutis vulnerum, corpore in ordine
tunicidal - Petala calyce paullo longiora lutea
an Alyss. minimum L. Nervosis a. calycibus
sub a. calycibus duo species habent nunquam
Valeriae Alyssum dictum cognitum nunquam,
Cacq. Bonte, et minus quod.

Lathyrus aphaca Leeuw. pedunculis
unifloris, corollis Dyphyllis Legumines
ovalibus compressis versus margines.
In Dfort ai Vulnini.

Polistichum lonchitis lanceolatum longe angustum
sinuatum, ante rupes, stipitis lonchitis lanceolatis
longe sagittatis, floribus pinnarum, pedun
culis axillaris superior generalis, vasculis
magnis nigris inferne sinuatis, magnis lo
bis rotundis. Legumen. Lobum longissimum
ulcerata in pedunculis fructifer cum
aspectu, accumbens, cum margine
ex 3 spinarum. Stirpis L. radicis, in gen. Dfyllum
legumen. sagittatum
per superiora la Dfyll. Al. L. Cicum
tunc pinna ai Vulnini, id in pannis
Lonicera periclymenum Leeuw. calyce ovatis coppetiis
mucronatis, filis alternis bicuspidis acutis;
caule stipitis aspera, ramis gracilis
tenuiter adpressis. Stipites ovata lata
mucronata margine alba.
Jam ai ovaria
an L. tunicaplen

Cornutella canadensis Leeuw. fructus erecta filiis
nervis obtusis glaucis, terminali ovata,
corollis gracilibus, legumen. orbicularis gra
dilis. Lavina in collib. crant.

Ophrys sativa. Leeuw.
Ai Vulnini d. Sept. et circa Dreiei ovis
Carl. pratis multiplicibus, filiis ovatis latis
ovis, floribus petalis revolutis ovatis
fl. appendicis pedunculis - Dodie.

Hieracium subcaulescens Leeuw. fol. lanceolatis dentatis
ovis rugosis filiis, rugis hirsutis bifidis, filiis singulis
e subcaule prodeunt, floribus ovatis, calyce ratis
brevi hirsuto, radicis appendicis ovatis binis ovatis, ex
ovatis adspersis. In collis granitis ion. Bergschaal
an St. Hieronymus Alsini?

Scabiosa arvensis leon. corollarum labia subæqua, folia omnia decompositi pinnatifida, caule aculeatum inermis. In Brugg prope Berna, Aprilis — et l. c. Allion.

Primula laurina pyrenaica flora in ing. Brugg

Thymus virginicus leon. pedunculis multifloris axillaris opposita, folia lineari-lanceolata margine revoluta — in saxosis collis. Lausanna. Cand. [illegible] suffruticosus glabra, ramis virgatis, subtilissimis, foliis undique canescentibus pedunculis axillaris subsessilibus, calyce villis albidis bilabiatis, laciniis setaceis, corolla pentapetala labiata, inf. laciniis ovatis, sup. lanceolatis.

Satureja nivea leon. pedunculis oppositis axillaris bi-afloris, foliis ovatis, calyce dentibus subulatis aristis, corolla [illegible] pentapetala pura, stylo corolla longior.

Lepidium hirsutum — Brugg.

Lycium europæum — Allion, Bau, Lausanna, Nosser

Trifolium pilosum leon. vagina compressa, calyce magnitudine corollæ brevior inæquali, foliis numerosis: ut plurimum hirsutum. Avenches. Delessert — Cordier Jura in ripis

Vicia tenuifolia leon. legumina pedunculata solitaria, foliis lanceolatis, stipulis semisagittatis — Italia in herbidis — Pedemontium illustr. fl. Pedemontii axill. [illegible] caulis. Lyon. non congruit planta illius polymorpha.

Halleria monstrosa leon. fol. lanceolatis pedunc. terminalibus terna, inferioribus [illegible] calycis flabel. magnitudine late membranaceis albis, petalis rotatis. — Ad pedem montis Nosser in herbidis

Lathyrus Nissolia Ad ped. Nosser in rotis.

Arenaria rigida leon. foliis subulatis, caulibus planiusculis et cum [illegible] difficilis, floribus parvis albis, sepalis obtusis ciliatis subtus [illegible] rigidis, ovatis magnitudine, capsulis cylindraceis [illegible] superantibus. Prope Ch. Pendulum, a pen. differt caul. planiusculis, et longitudine

Orobus pendulus leon. bulbis [illegible] bulbis [illegible] indivisis, [illegible] caulibus [illegible] extricatis, [illegible] ovatis. In pratis Berne, et Porrentruy. Fl. purpureus, bulbis nudis villosis.

Vicia glabra leon.
pedunc. multiflorus, floribus [illegible] solitarius, calycis [illegible] villosis, inferiore [illegible] longior. Allogy [illegible] polyphyllus.

a Vicia Cracca, vix [illegible] differt tota planta excepto [illegible] glabra, stipulis non semisagittatis — Circa Berna in [illegible] dumetis.

Trifolium pannonicum leon. involucri partialis foliola ovata reflexa, foliis pinnatis, pinnis subtus [illegible] margine [illegible] inæqualis.
In [illegible] ripis Ins. Berna, Lausanna; Splug.
Trifolium [illegible] — non [illegible] Dentatis nec T. equal. L. T. [illegible] [illegible]

Spergula arvensis — in pratis a Splugen in [illegible]

Scrophularia peregrina [illegible]
corona urceolata, et in [illegible] [illegible] Brugg
foliis [illegible] obtusis glabris [illegible], [illegible] axillaris, et terminalis.

Jacobæa [illegible]. Brugg in collis et [illegible] Berna.

Hieracium [illegible] Cand. Pilos., folia III Jungermannia p. 118 — in [illegible] circa Solum

Rhododendron [illegible], folia glabra, [illegible], et [illegible] hispida.

Hypochœris [illegible] leon. foliis pinnatifidis, pubescentibus, marginatis, scapis [illegible] Laudis flora. Dentata vel [illegible].

Crepis lyrata leon. fol. radicalia lyrata pubescentia, caulinis subsimilibus [illegible], caule subramoso, calycibus [illegible] — In Nosser [illegible] herbidis Fl. pari, [illegible] Pilos. [illegible]

Orobus niger. Mt. Nosser.

Hieracium [illegible] Allion Mt. Nosser

Lithospermum tenuiflorum leon. pomulorum laevia lineariter angusta, involucrum brevissimum, pentaphyllum persistentia 3-5 phyllia — Petala [illegible] [illegible] dense [illegible], extremum [illegible] magnitudine. Arenaria orbis-Allogy [illegible] angustiores 5-septa, calycis foliola, stylis [illegible] [illegible]. Caulis glabra terminis ramis. Folia radicalia [illegible], [illegible] pinnatifida 3-4 polliaris, caulina subsimilia [illegible] [illegible] terminalis 5-8 radiata, umbellis multifidis. In [illegible] Berna, et in Mt. Nosser

Plantago media Mt. Nosser.

Trifolium spurolifolium Scen.
adversum, caulibus subtenuis, foliis ovato-lanceolatis obtusis, caule erecto nudo. — Circa Schiraz
a viaeq, Salour in Mistry invenerunt.

Caulis 3-4 pedalis villosus nudus, fol. obtusis adpressepilosis, inferioribus lineariis angustis pilosis, caulinis axillaribus pedunculis, pedunc. fol. multo longioribus villosis, bractea ad capit. longe amplexantes villosa rufescens, legumen. lineare glabrum
Ann. Bracteae glandulosae.

Astragalus flavescens Scen. villosus linearibus acutis fol. radic. lyratis, caul. pinnatis, pinnis paucijugis linearibus longioribus — In Persia — Caul. brevissimus villosus, fol. glabra, col. pallens flavus, vilosus calyx pedicellis subtripollis, stigmata capitata terminata

Campanula Astri Scen. Circa Tauris in Cypris
speciosis bordis. Fol. oblongo ovatis undulatis in petiolum
col. purpureus lineis linearibus acutis corollae multo brevioribus.

Crataegus cordata Scen. glabra, fol. radic. lyrato-pinnatis, caul. ternatis, spinis calycis in longitudinem hypanthii cordatis! acuminatis. — In Kurdistan circa Bua
H. terminalis solitarii magni flavi. varii ejus
inaequalis rufescentes, spinis calycis linearibus,
calicibus magnis, carnosis, seriem duarum in spinis
non cordatis, tertia cordata-oblongis. Legulae 5. dentatae. Pappus pilosus rufus villosus. Recept. nudum

Stipa eriostachya — In Cyp. Bua. H. fructuosum

Scabiosa decaticulata Scen. Circa Tauris. Sc. multifida

Phalaris canariensis — circa Lesina in granetis

Dactylis macrostachys Scen. panicula spicata nude
pedunculis tenuis brevis, calmagnis inferioribus
fructiferis remotis. — Tauris et flor. Dalmatiae Cultum. e generibus hujus regni.

Phleum aculeatum Scen. spica ovata parva aphylla
culmo erecto. — Tauris, Kandum, et flor. Dalmatiae Cyp.

Hordeum odontolepium Scen. glumis lateralibus nudis
mutis, inferior foliis sedens ad basin stipitis
floralis, lateralis tenuis, intermedia brevis. Tauris.

Aegilops ventricosa
Calamagrostis genuina — Osiris, Bratsja - Stipa hispida

Agrostis vesicaria Scen. panicula spicata, ovatis
glumis brevibus mutis, calmagis inferiore procumbentibus et aristatis — In Cyp. ager Dalmatiae versus
Gr. vad. vag. pan. diversa g. pad. vind. Lacerij mont.
Schenck. exp. p. 130.

Dactylis hispanica Scen. panicula spicata pilosa,
pedunculis tenuis brevis, calmagis obtusis
fructiferis. Circa Tauris in granetis.

Carex compacta Scen. spica compacta compacta
depauperata foliis floralibus se longiore involucrata,
spiculis apice masculis inferiore femineis, culmo
compacto. — Salour in ulvis Hyrc. a. Caspio
planus-convexus compactus basi attenuatus
brevissimus et basi vaginatus. Aspectu habitu
ut congesta sed spiculis androgynis, et habitu
scirpum hypogynum continentem gerunt.

Scilla elongata Scen. spica longe tenui, bractis
integris oblongis acuminatis. e Dalmatia.

Iris Bucharica Scen. Hell. Bisp. n. 1427
Schenck. gram. Cabul. spatha tubo compacta p. 4.
Gr. vag. off. Wing tubo. alg. Tavil. Bank. 2 p.
in pa. Bruyn et alibi in Dal. 457

Off. J. reticulata — In Differt culmo multo altiore,
spica longissima, spiculis multis majoribus, inferiis
mutis, tegumen. interiorib.