



Plantarum in Dalmatia

observatarum

et in Systemate Vegetabili

A. P. Secundo

recensitarum

Enumeratio.

his accedent

aliquae ex Syst. Veget. et Pömer

Depromptae.

ex Volumine II. Syst. Veget. Secundo

Cruciferae.

Nasturtium Lippicense.

Rad. radicalibus petiolatis oblongis dentatis sub-
lymbois, superioribus pinnatis-petiolatis, lobis
linearibus integris, siliquis linearibus declivibus

Hab. in locis rupestribus Croatiae et Dalmatiae

Potruschley in litteris.

Dalman. 653. p. 3.

Cardamine montana.

Rad. petiolatis glaucis glabris pinnatis-petiolatis, seg-
mentis 5-7 obtusis uncinatis repandis, caulis dif-
ferentissimus, siliquis erectis glabris.

Potruschley in litteris.

Hab. in Dalmatiae peninsulae Istrianae (olim Hydruntis)
ad littora maris adriatici. Potruschley. (1)

Festuca triquetra.

Caulibus basi suffruticosis, ramis triquetris
ad quadratis, fol. canalicatis, radicalibus
obovatis petiolatis, caulibus oblongo-lanceolatis,
stilo longo deciduo, stigmate simplici. —
Calyci erecto glabro trifido, pubescente
stiletto pubescenti. Petala longa unguiculata,
limbo ovato obtuso. Brachium tomentosum ellipti-
peltatum; Stylus ovato longior; stigma intrinsecum.
Liliula elliptica, stilo deciduo mutica, peltis mi-
nime sparsis stiletto fere, 7-8 lin. longa,
3-lin. lata.

Alyssum triquetrum. Postelschley in littorij.
Hab. in Dalmatia ad rupes arid. Clissa. etc.
5. fl. albit.

ex Syst. Vegetab. Boemer et Schultze

Vol. II

p. 561.

Poa festuciformis Host; panicula
aequali patula; spiculis lanceolatis novem-
floris, flosculis basi villo connatis obta-
sis 5-nerviis, fol. calicis, ligula oblonga,
radice repente. Willd. Enum. et. Berol.

I p. 106. Host Gram. austr. III p. 12
t. 18. Huc quoque, teste Schrad.

Festuca palustris, panicula erecta,
spiculis 5-6 floris nudis distichis muticis,
foliis planis, culmo unguiculato v. Leenuy
Heise Tot. Dalmat. p. 72.
Culmi glabri teres, & geniculis inferioribus

indicant; 1/2 ped. et ultra; folia conj.
sed, firma, facie canaliculata scabra,
ad oras et carinam aculeata. Vagina
glabra. Ligula pubescent. Panícula
erecta aut incurva, rachis scabra
vires, ramis scabris, superioribus 2-3nis,
mediis pluribus. Spicula subtortis, sa-
vina et flavisente vire; aliquid glauca
in quibus aculeis sunt, majoribus hi-
nervi.

In Dalmatia insula Ugliano
prope Zarem. 2.

Ibid. Vol. V. p. 93.

sp. 14. a. fol. quinibus

Campanula Dostenschlagiana

caule ascendente, foliis omnibus
cordatis veniformibus dentatis, sinuibus
basos rotundatis, pedunculis axillaribus
divisis, submultifloris.
Caulis glaber, ut tota planta, prope
sulcatus. Pedunculi foliorum et ultra, basi
dilatati. Folia 5. circiter lineis longa, 9
lin. lata, dentibus acuminatis apice glandu-
loris; superiorum parum angustiora. Pedunculi
inferiorum simpliciter, petiolo brevioribus, superiorum
divisis; foliis longioribus, unde habitus plantae sub-
compositus. Corolla cum geminis $\frac{3}{4}$ circiter phlo-
is; hoc semisphaerum, 2. fere lin. in diametro.

calycis laciniis linearibus lanceolatis, mucronatis
patentibus; cor. infundibuliformi campanulato,
dilute fuscis. Stigma profunde
de triquetro laciniis revolutis. Siccitate
resiccit, et consistens quasi Impatiens,
attamen nullo modo ea Erinoidibus. Ab
Clatone in the diglo major differt.

Deit. cl. Portenschlag sine nomine
cum plantis aliis in itinere suo Dilma-
tinae lectis. Ab omnibus nobis nobis dicitur
sa toto celo.

Ibidem l. c. p. 136.

^{6. foliis subij}
Campanula pumilio de Portenschlag.

suffruticosa, caespitosa, subcaulis, foliis
densissimis imbricatis linearibus recur-
vis margine revolutis, supra nitida-gla-
berrimis, subhinc nivo-pilosis, floribus
terminalibus sessilibus, calycibus anno-
tinis persistentibus peruliformibus.

De Portenschlag in MS.

Facies exactissime caespitij Sileneae
fol. angustioribus. Radix apice divisa est
in lignis Androsacibus, Saxifragis &c. Folio
anne priorum emarid, argenteo, archi-
me imbricat, presentij vernationis resissa,
rigida, 9-10 lin. vix $\frac{1}{2}$ lin. lata (siccata).

et reducta adeo, ut pagina inferior
plane abscondatur: fasciculus pilorum
ad apicem juniorum, supra sub foliis
lanceolata et acuta, aculeo imbricata.

Cal. vix 3. lin cum germine parvo,
hispidulus, laciniis linearibus rigidulis,
erectis duplo longioribus, cor appressa
infundibuliformi-campanulata duplo bre-
vioribus. Stigma in vestris floribus blotum

Calyx anni prioris persistens, laciniis
revolutis, diametra $1\frac{1}{2}$ lin. et ultra.

Ad specimen D. de Kottenschley.

Omnium elegantissima densissimisque caespiti-
bus summae jugis montis Diuoro senora
vestris. D. de Kottenschley.

Sementi di Fiori
Gifia - Putera - Tulipani -
Sali di.

Phalangium liligo -
Papaver corniferum -
Adephus nigra -
Nigella damascena -
Digitalis purpurea -
Aphodius fuscicornis -
Anthrenum varicosum -
Mora pavonis -
Heliotropium indicum -
Thlasia arvensis -
Anthrenum arabicum -
Anthemis perfida -
Aphodius fuscicornis -
Stenobothrus -
Muscus, musculus orient. -
Lunaria annua -
Digitalis purpurea -
Pinnia elegans -
Zinnia -
Gomphrena globosa fl. albo et rubra -
Anemone pulchra bicolor -
Aquilegia bellata -
Vinca multiflora -
Oryzium -
Kochia -

Paris

in Padova.

Bisi odoriferi color laevis
— Di Vaga

Lithospermum in ins. Arabia ad vici
viciq. Jentes. J. thamm. et thamm.
Judea ann. singula flos. Caules ter-
ribiles et subpungentes. Stipulae
tota pili albi adpressi, ut the plant
caulis hispida stipulae apice fruticosa
Hinc linearis-epetulae oblique
marginatae. Stipulae ovatae
apice foveolatae. Corolla
nuda. Antherae fere distinctae.
Stylus corollae secundum
fere inter insertus. In longis
caulis. Calyx 5-partitus. Stamina
caulis tubo parvis brevioribus. Stylus
hypogynus. Fructus capsulae




Arabia, flos


Thyris *Hesperis* *Arabia*
fructus parvis brevioribus
Stylus superioribus minoribus. Antherae 5-ae
in im-parte tubo. In calyce. Semi-4 antherae
apice acutiusculis brevioribus apice.

orchy

hymenocallis



11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100



[Faint, illegible handwritten notes and sketches on the left page, including a small drawing of a flower bud.]

- 18 Signe de sel.^{co} a Capouso per mare
 19 a Capouso id.
 20 a Bovoglia a S. Maria
 21 a Capel Nuovo a Oragnino
 22 a Salona Clivo a Spalato
 26 a S. Pietro per mare
 28 a Nevezi. S. Pietro a Almisia
 29 Almisia Uvullia a Macarista
 2 Luglia Sul Diokoro de S. ant. alle Jan.
 5 a Brissac nel Primorio per mare
 6 a Hastioy id.
 7 a Metkeich, Vido e Vegorze
 8 de Vegorze a Sotiza
 9 de Kozija a Duaro
 11 de Duaro ad Ughjiam a Siga
 15 de Siga a Muck ad Ozon
 16 Sulla Sirlia
 17 a Sestica

Stachys *Spicivanda* Capelli grasso
 Vran in un botto di 2 sp. junc. la rian.
 Vestiboli 6 flori. Cor. ^{alba} calice duplo largiore
 labio superiore fimbriato intes. calice
 pubesce, internod. trilob. lobis later.
 lobis reflexis obliq. ovatis, internodi
 junctura ostendit ochroleum lineis ^{seriatis} rubris
 lobis utro subulatis; filamentis albis
 calicis purpurascens, antheris flavis
 calicibus, setis rubris; calicibus
 obliq. tetragonis, angulis mucronatis
 adve flante in partibus.

Stachys *Spicivanda* Capelli grasso
 Vran in un botto di 2 sp. junc. la rian.
 Vestiboli 6 flori. Cor. ^{alba} calice duplo largiore
 labio superiore fimbriato intes. calice
 pubesce, internod. trilob. lobis later.
 lobis reflexis obliq. ovatis, internodi
 junctura ostendit ochroleum lineis ^{seriatis} rubris
 lobis utro subulatis; filamentis albis
 calicis purpurascens, antheris flavis
 calicibus, setis rubris; calicibus
 obliq. tetragonis, angulis mucronatis
 adve flante in partibus.

Hyssopus ~~capitatus~~ *gibbi*? sive *Torretta* &
Spatol
Veritatem longior aerea, aerea alij. Capite
dentate ovata, apice uncinata, ciliata.

Aliam speciem: Sp. et *Quercus* de Capaci

Petala emarginata staminibus basi dilatatis
spatulis tertia longior. Umbellae alij quid bulbosae

Hyssopus flammulae *crabata*, petalis
apice lobatis sive obtusis, sublimi in ligno
stipite persistente orbiculata, foliis integris
radicibus ^{spatulas uncinatas} subrepentibus, caulibus *linum*-
spatulatis, imbricatis. *Trigoni* *unij*
by *ellij* *Male* *Mulina* *dicti*. *apiti*.

Hyssopus *triqueter*. Caulis basi suffruticosus
ramis teretibus, exsiccatione angulosis,
stigmata biloba

Onobrychis *fundata*? *Dr.*

caulis supino, foliis lanceatis obtusis mucronatis
scaberrimis, racemis 2-4 floris, calycibus
ciliatis corolla brevioribus, petalis inequalibus
leguminibus pubescentibus dorso cristato, crista in
laciniis 5 inequalibus integris, apice uncinatis
disco disco aucto ~~sub~~ pubescente *fundata*
que *propendit* ^{in fund} a *propis* pubescentibus

Hab. in agro Spalatensi in colle la
Torretta ☉ Pl. Apiti.

Radix annua simplex fibrosa. Caulis supi-
ni sericeo-pubescentibus striatis. Folia impari-
pinata nunc cuneata sur angustior, nunc
obovata s. latior, obtusa mucronulata subse-
ricis. Racemi pauciflori (2-4 fl.) pedunculatis
triplo brevioribus. Stipulae oppositifoliae angustissimae
by *ovatis* acutis late lanceolatis.

Flor. vrbri glomerati. Calyx 5-fidus villosus
laciniis linearibus inaequalibus, inferioribus minoribus.
Corolla calyce longiore ^{partibus} inaequalibus, veris
scilicet carina longiore, carina alia. Legumen
(immaturum) eadem parte adpressum ^{induratum}
albiflorum, dorso unguato, unguis in laciniis
late lanceolatis (4-5) integris ^{reticulatis} apice adunatis
inaequalibus partibus, lateri interno sive pediculi
cubum respiciente plano ^{latiore} ~~subtriangulari~~
disco aculeato foveisque ad eundem ^{etiam in parte}
usque profundius adpressis ^{sericeis}; foveolis
inaequalibus, duabus lateri interno proximis
caeteris multo maioribus. Semen unicum
subrotundum.

Videtur differre ab *O. fauolosa* DC. Prodr. 2. p. 346
disco leguminis enervio, vix pubescente foveo.
Usque pariter pubescentibus.

Cerastium Samnitanum? Ser. DC. 1. p. 421
s. longistylum Tenore.

tomentoso-virescens
caulis diffusis geniculis incurvatis, foliis
^{angustissimis} linearibus margine revolutis internodiis
longioribus, floribus dichotome glomeratis
bracteis scaris partibus, internodiis longioribus
pedunculatis, ~~bracteis~~ sepaliis arcuatis margine
et apice scaris basi carinis, petalis calyce
duple longioribus usque ad testam partem
bifidis unguicatis, staminibus decem
corolla duple brevioribus, stylis 5 apice
revolutis pubescentibus. Capsula subcylindrica
calyce longiore

Orchis flaccuens
bipunctata

Stachys muricata

Lithosperma crassipes

Onobrychis foedata?

Trifolium pauciflorum

Alyssum dalmaticum

Cerintho purpurea

Cerastium stachyfolium

Cerintho atrocubus

corollae dentibus acutiusculis pectatis, bracteis
atro-purpureis, fol. caulinis ^{cordatis} amplexicaulis
by 1 radicalibus spatulatis

radix perpendicularis caerulea subterris nigra
caulis teres ramosus cylindricus;
folia radicalia spatulata pectatis, caulina
cordata ^{ovata} amplexicaulis sessilia,
omnia integerrima laevia et obliqua. Bracteis
cordatis amplexicaulis, superioribus oblongis, atro-
purpureis. Calyce profunde 5-fido lobato.
Corollae tubulosae emissae apice angustatae
5-dentatae, dentibus triangularibus acutiusculis
pectatis, fere albis, tubo ~~obovato~~
fere ~~triangulari~~ flatissimo, partem inferiorem
albida, medioque majora ^{dentata} atro-purpurea, superior ex
albida in purpuream atro-purpuream dentibus pectatis

Stamina media tubi infera, antheris sagittatis -
linearibus plumbeis. Stylus filiformis
subrectus glaber. Nuc. una (forma alba
dura) bilobata pedunculata.

Cos. corio-lanceo versus apicem deorsum inflato
limbo connatis dentibus strigosis reflexis
bracteis. Tubi basi albidus, dein tenuis
involucris, pinnatis, deorsum gibbatus ibique
ruga subterminali, albidus, limbo involucri
dentibus, denticulis

Profunditate ^{conspicua} calyce 5-fido, dentibus

3. superioribus saepe majori ex parte coalitis.

apice lobis ovatis angustis; cordatis

lobis superioribus pinnatis erectis integris,

inferioribus trilobis, lobis ^{intra} ovatis strigosis
intermediis longioribus, nec cordatis lobis.

Pl. cordata-ovata villosula.

In gressibus in ripis & agris.

Phanera fl. Nivis ovata pedunculata, pinnis
vix alijando glandis Stamina pinnatis

[Faint, mostly illegible handwritten text on the reverse page]

Scilla bulbis ovato conico concentricis tunicis
ext. externis albidis; foliis omnibus radicalibus
lanceo-lanceolatis marginibus
tenuissimis ciliatis, basi vaginatis
glabris, scape breviter; scape tereti
vires (3-4 palmi) in parte floriferam
angulata corollae-virescente clausa. filii
longioribus violaceo-aequalibus
filibudibus multifloris, filamentis brevibus
distinctis, bracteis albis minimis angulis
raris antipetalis, epica petalibus, pediculis
clausis adpendentibus teretibus, filamentis
semipetalicis, fructibus longis; corolla
hexagona petalis, laminae ante-oblongis
obtusis virescentibus aequalibus limbo media
^{ignominia} filii longioribus
obtusis, filamentis adpendentibus basi

obtusis, subulatis, antipetalis immixtis
petalis (paucis) aequalibus virescentibus
angulis. foliis brevibus epica subulatis
obtusis virescentibus.

In saxibus umbrosis collibus in insula
supra Nulla prope Lebanonem Agriensis.

Allium luteum. foresis in collibus paucis
bulbis ovatis simplex viso duobus major.
Stapus foliatus, foliis semiteretibus parvis
superis longis, basi longe vaginatis
spatis brevibus, bracteis basi membranaceis
albidis, neris 5-7 in collibus prominentibus
epica subulatis, altum duobus longioribus.
Umbellae laxae multiflorae, pedunculis aequalibus
vires aequalibus, petalis oblongis obtusis

luteis staminibus fere duplo brevioribus,
filamentis simplicibus.

Die 19 Junii

Centaurea alba? Hispanica Castellae. 21 Junii
calycibus ovatis, squamis inferioribus apice spinosis
sive spinis flexilibus partim, medio ciliis inaequalibus
apice vix spinosis, ~~interioribus~~ ^{superioribus} ciliis apice
scarioso-membraceis, floribus omnibus squamis
ciliis nec radiatis.

Stachys. 29 Junii Hispanica
Cordubensis, calycibus pubescentibus, lobis
reflexis, interioribus obtusis, compositis
punctatis, ~~exterioribus~~ ^{exterioribus} ~~obtusis~~ ^{obtusis} ~~compositis~~ ^{compositis}
Calycibus calycibus lanugineis, dentibus

Concedatis aristatis duobus inferioribus
brevioribus ^{superioribus} ~~compositis~~ ^{compositis} bracteae ovatae compositae
acuminatae, ciliis integris, bracteae
setaceae, verticilli sub 7-flori. In floribus
dentibus fetet. Hab. inter Duzen
et Mincum, et etiam in Biskonia ^{et Juvie}

Galicum? ad viam Romae, Biskonia Biskonia.
29 Junii.

Corolla unguiculata-campylata 4-peta. lobis
vix oblique, staminibus duplo brevioribus, antheris
luteis. Pistillum ^{apice bipartito} ~~triloculare~~ ^{lobis} ~~staminibus~~ ^{staminibus}
brevius. Stigma ^{apice} ~~triloculare~~ ^{lobis} ~~marginibus~~ ^{marginibus}
subtus rotundum acutiusculum. Pediculus in
vix vixem lignosum.

Andropogon - Dobruza prope Vulturem.
ad mare in gromis
Herba grassa, characteribus Andropogoni Persici
exacte instructa.

Scrophularia. Crescit ad vias die 26. Nomin
in Bicklow die 2. Julii.
Corolla polyloba resupinata lobis ovatis
in magna laciniis lateralibus albis, raris
media atropurpurea lobis lobis ovatis, super
iore minime ovato viviscente. Stamina
dynamis filamentis pilosa glandulosis, filis
injetibus, antheris luteis horizontaliter
affixis longioribus exteri. ^{filis} ^{longioribus}
~~condonatis~~ ^{condonatis} lobis lobis ovatis, stylo calycis
lobis adnatis. Calycis lobis et radice suffi
gras. In viis ~~vulgis~~ ^{vulgis} ^{vulgis}

Carduus maritimus Lat. Petala ^{vel etiam} ^{recte} ^{carduus}
gianta, quod non cernit in iuncis ^{in iuncis} ^{in iuncis}
Maonia laciniis ^{vel tunc} ^{tunc} ^{sub} ^{sub}
conspicuis. Circa ^{in iuncis} ^{in iuncis} ^{in iuncis}
Abat Duare ad vias super ^{in iuncis} ^{in iuncis} ^{in iuncis}
Pentameris.

Centaurium aspidota 2 Julii Bicklow ^{vulgi}
^{maonia} ^{speciosa}
Radix perennis, subterrena prima super
perennis. Caulis subterrena foliis angulatis
foliis ovatis, ovatis lobis, et sub flori
incompleti. Stibi coriacea ^{in petalis}
~~radicibus~~ ^{radicibus} ^{radicibus}
radicibus ovatis lobulatis glaberrima, ^{vulgi}
angulata subterrena, lobis lobis ^{vulgi}
angustiore integris, superna subterrena
super ^{in iuncis} ^{in iuncis} ^{in iuncis}
ovatis subterrena (ovatis lobis in
positione folii axilla). Calycis ovatis
oblongis, ovatis lobulatis lobis
lobis ovatis, in dimidi ovatis ^{in iuncis}
ovatis, lobis lobis albis, ovatis lobis
lobis ovatis, florum ^{in iuncis} ^{in iuncis} ^{in iuncis}
lobis, lobis lobis lobis, lobis ^{in iuncis}
lobis; ^{in iuncis} ^{in iuncis} ^{in iuncis}
antheris in lobis lobis lobis

Physiciana. Soliga interlegum et
Lappa Zygua d. di g. hylis in saxoia gij.
Covokta pirocarina dein semiaperta
5-partita. Stratum basi dilatata ibiye alia
Flory sparsim glanata peltantef.
Pila glabrata cinerea unicolor nigra
artibginat - argute dentata.

[Faint, mostly illegible handwritten text, likely bleed-through from the reverse side of the page.]

Physiciana. Soliga interlegum et
Lappa Zygua d. di g. hylis in saxoia gij.
Covokta pirocarina dein semiaperta
5-partita. Stratum basi dilatata ibiye alia
Flory sparsim glanata peltantef.
Pila glabrata cinerea unicolor nigra
artibginat - argute dentata.

[Faint handwritten text, possibly bleed-through or a second entry, including the word 'Physiciana' and other botanical descriptions.]

Satureja ^{fring. utall. scin} ~~corolla ultra~~ ^{lobis} lobis ^{impunctis} ^{3.} ^{imbricatis}
 coloribus, galea ^{obscure} ^{dentata}
 sursum involuta, ^{frons villosa}, calyx angustis
^{dentibus} ^{imbricatis} ^{minimus}, hinc bilobatus
 lobis ^{complanatis} ^{ambulantibus}, ^{lobis} ^{caucibus}
 apice ^{imbricatis} ^{caucibus} ^{caucibus} ^{caucibus} ^{caucibus} ^{caucibus}
 nigrae ^{complexa} ^{super} ^{glabra}, ^{lobis} ^{lobis}
 Odor ^{diversus} ^{1.} ^{maxime} ^{Muret} ⁱⁿ ^{lobis}

Satureja ^{specie} ^{corolla} ^{albida}, galea ^{convexa}
 apice ^{retusa}, ^{bore} ^{biloba} ^{lobis} ^{subglobatis},
 nec ^{multo} ^{paucis} ^{majoris}, ^{frons} ^{punctis} ^{roscis}
^{albida} ^{partibus} ^{lobis}

Genista dalmanica. Differt videtur a Genista
 thebotanica in spicis brevibus crispantibus
 vixit purpureis ^{rectangulo}-^{vel} ^{sub} ^{trianis} ^{divanis}
^{partibus} ^{apice} ^{partibus} ^{minimis} ^{ultra} ^{partibus} ^{gemma}
 in majus in ^{dentibus}, ^{vixit} ^{partibus} ^{etiam} ⁱⁿ ^{galea}
^{vixit}. ^{Habitus} ⁱⁿ ^{vixit} ^{partibus} ^{etiam} ⁱⁿ ^{galea}

Spina ^{fringis} ^{triflorae}, ^{spicibus} ^{impunctis}

Navis ^{fringis} ^{triflorae} ^{spicibus} ^{impunctis}
^{pl. caucibus} ^{bilobis} ^{lobis} ^{impunctis} ^{1.} ^{caucibus}
^{lobis} ^{impunctis} ^{lobis} ^{impunctis} ^{lobis} ^{impunctis}
^{lobis} ^{impunctis} ^{lobis} ^{impunctis} ^{lobis} ^{impunctis}
^{lobis} ^{impunctis} ^{lobis} ^{impunctis} ^{lobis} ^{impunctis}
^{lobis} ^{impunctis} ^{lobis} ^{impunctis} ^{lobis} ^{impunctis}
^{lobis} ^{impunctis} ^{lobis} ^{impunctis} ^{lobis} ^{impunctis}
^{lobis} ^{impunctis} ^{lobis} ^{impunctis} ^{lobis} ^{impunctis}
^{lobis} ^{impunctis} ^{lobis} ^{impunctis} ^{lobis} ^{impunctis}
^{lobis} ^{impunctis} ^{lobis} ^{impunctis} ^{lobis} ^{impunctis}

Ad vixit S. thymum in Spat. Febrem

Crocus

Juxta mentem Th. Fay *Crocus* ~~relinquimus~~
noster non est nec *C. edneyi* Div. nec
C. serotinus Salisb., species aliunde inchoata
bene distincta, sed *C. Pallasii* Goldb., ita
evidenter ab inspectu unius specimenis
a *Reichenowii* eidem magis ex nostris.

Potest igitur *Crocus* versicolorum, a *C. versum*
bene distinctum (sic ille) adesse in *Detentione*

in libris ad cl. Willden., 1831.

Calendarium

1832

Dennig

6 Febr.

28 Scen:

Veronica agrostica

Helleborus niger

Laminium purpureum

2 Febr.

Senecio vulgaris

Viola odorata

Mercurialis annua

Thlaspi bursa-pastoris

Euphorbia helioscopia

Laminium maculatum

Camphorosma officinale

Ulmeny
18 J.

Crocus versum

~~Amantur~~

Medic. Practica

legit. abbas medicis officio in
reale Collegio de Madrid sub anno

1822

AB

Asphodelus luteus et ramus in spulibus
a Hispania tunc abbas S. Georgii
Allium odorum Vivianae Castellae
Astragalus monspulensis ibi
argenteus in Moscov.

Arum Drummeily Solis per traditionem

Alison Plantago Castellae

Allium rubicundum Almod. de Lamei

Aegilops trivene. sp. Iran

Andryala lutea Boeckh

Althya rosea in m. Male Mutine

pro Imperium

— Cannabina Castellae

Aconitum napellus. pro Poyosa. e Boeckh

Alnus glutinosa } Narva

Althya sp. } Narva

Anthemium monspulense pro Mutine e Boeckh
denn e Verke

B
C
D
E
F
G
H
I
L
M
N
O
P
Q
R
S
T
U
V

Erven lantid

[Faint handwritten text, likely bleed-through from the reverse side of the page]

Bieberzeit - Cantoreo

Baldianwurzel - valeriana-radix
Brockwein, stein - Tartaro emetico
Brandwein - aquivite

Polysantif

Andrych lous

Lody hufjung

Solly Drupfen

Wij maxim Ing.

Bromus pinnatus ad usum

Butenmy

Dyghtholm schiffen in ylorling ad
jedem monat Surujarra
contu Dygl

B
C
D
E
F
G
H
I
K
L
M
N
O
P
Q
R
S
T
U
V

Campanula cordata hibernica Inguini et
Spatia in cultis.

Cynoglosson pictum ib. et Alpe, ^{Thyris} _{Wij.}

Coronilla sanguinea Inguini et Spatia.

Convolvulus tenuiflorus in Mt. Sclon. in alpe

Cyclamen europaeum Nivum de Capella

Crocus immixtus ib. Sp. Wij.

Cyrtis argentea in Alpes. Br. Wij.

Clypeola Frag. Sp. ^{Wij.} _{Wij.}

Coronilla varia Br. Wij.

Callitriche ^{varia} _{Wij.} in alpe

Cephalanthus ^{varia} _{Wij.} in alpe

Chrysanthemum ^{varia} _{Wij.} in alpe

Chrysanthemum ^{varia} _{Wij.} in alpe

Chrysanthemum ^{varia} _{Wij.} in alpe

Chrysanthemum ^{varia} _{Wij.} in alpe

Chrysanthemum ^{varia} _{Wij.} in alpe

Chrysanthemum ^{varia} _{Wij.} in alpe

Chrysanthemum ^{varia} _{Wij.} in alpe

Chrysanthemum ^{varia} _{Wij.} in alpe

C
D
E
F
G
H
I
L
M
N
O
P
Q
R
S
T
U

Cyperus longus in partibus humidis Salinis, Alm.

Cirsium - Missa, Alm. - Scogli di Naxos

Chenopodium - ambracorum Naxos

Conium maculatum - Naxos

Chenopodium - Naxos

Cladonia - Naxos

Cypripedium - Naxos

Campanula trachelium - In Naxos, et in aliis
bus qui signum.

obliquifolia? circa Xuffa.

glomerata in his signis et Naxos

Dysferus subsp. speciosa
latini dicitur methoicis
ad flos

| |
|---|
| D |
| E |
| F |
| G |
| H |
| I |
| L |
| M |
| N |
| O |
| P |
| Q |
| R |
| S |
| T |
| U |

~~afgättan efers aactoro~~

Evrum lantoidy Inyuni Alotini

Ephedra prope Clifan

Echin in arce Clifan

Euphorbia spina Kapus

Euphorbia Mije spina di Capiti

Euphorbia unidentata N. A.

Euphorbia patens N. A.

Euphorbia - vert.

e

7

8

HN

9

L

m

N

o

2

2

2

5

7

u

Genung - aron, drage -

Galim - M. M. go inst. 100 p.
van fern? 100 p.
venn inst.

Phylogeny - methosist affy 8
Galego ib.
Epilipff detohit
Ladman } 100 p.

8
KH
9
L
m
N
o
a
2
a
5
7
u
A

~~Hyacinth - pratensis, pitillensis~~

Hyacinth romanus in agro Libanensi
trans ulam S. Georgii
et in vivariis Lepalli

Hyacinthus comosa Mispur Bouchi
Helleborus hyemalis Bouchi Lagonis, Klerke

Hyacinthus vulgaris In Hanoa

Helleborus vulgaris Inter hunc et

Helleborus Sphond - Bell -

KK

g

L

m

n

o

p

q

r

s

t

u

v

w

x

y

~~Saturago clathrata~~

Linum halepense ^{in cultis}

monspulana ^{ib. in sacris}

Lycium . . . sp.

Lithospermum purpureum

Ligustrum vulgare ^{ib. ad seip.}

Leucospermum

Lycium

Lobelia

Laurus

Laurus

Linum

Lycopus

Linum

Chrysanthemum

Scrophularia

Lithium

Lycopodium

L
M
N
O

2
2
2
5
7
u

Orvitha pyrenaeum Cappelli

Orvitha pyrenaeum Bickner

Orvitha pyrenaeum Jago

Orvitha pyrenaeum ^{sign} *Orvitha pyrenaeum*

Orvitha pyrenaeum

Orvitha pyrenaeum

Orvitha pyrenaeum

Orvitha pyrenaeum

Orvitha pyrenaeum

0
2
2
2
5
7
11
14

^{Poa. 2^{um}}
Panicum ~~gambellianum~~ in agro libanico.
Sobique trans altem S. Georgii, et in Borraja

ad iug

Borria borraja - Inyana

Polygala vulg. Pinea di Casale

Ab. Sp

Phytolacca decandra Ab. Inyana

Passiflora - Scopidi Nor - Inyana

Pempellia Inyana - Ab. Mar. 11.

Populus alba Nor - Vert.

Potamogeton ~~fluctans~~ Nor

fluctans ib.

pepliat ib.

Phellandrium Nor

Passiflora Vert.

Populus tremula ^{Vert.} Inyana

2

2

2

5

7

4

1

1

[Faint, mostly illegible handwriting on the left page, possibly bleed-through from the reverse side.]

Queck Silber - mercurio

[Faint handwriting on the right page, likely bleed-through from the reverse side.]

2
2
5
7
u

Commonly millefoliated in eye to bene.
trans. collin. S. Georgii, et in Borja.

Necesse lutea ubi in spiritali. Inguv. sp.

Commonly aris. In partij Iny. sp.
epithagay in inq. Saloniq
repens
et var. fl. pl. } ib.

Commonly in Bo. Capta
corrip. Sp.
ballus. Sp. Capta

Commonly in jungle Inu ad reg. et
(Kerhavin)

Rottbulla inu. v. h. sp. S. Capta

Commonly in Myra. Inu. h.
Lingua Vidua flus

Rub. p. m. in. v. l. in. v. g. et
Rogus, circa Blato, et circa Durick
in eye d. m.

2
5
4
u

Stachys muricata Pennel

novae horti botanici et Medicorum

in horto botanico et Medicorum

Solanum dulcamara L.

Sida acuta Methon

Sida acuta Methon

Sida acuta Methon

Sida acuta Methon

Sida acuta Methon

Sida acuta Methon

Sida acuta Methon

Sida acuta Methon

Sida acuta Methon

Sida acuta Methon

Diplazium tomentosum } in agris Libanici. stricte
resignatum } by trinis collibus S. George
ibidem

Diplazium repens ⁴⁵⁵ in agris Libanici et Spalati

Diplazium complanatum ib. et Libanici

Diplazium ib. _____

Diplazium complanatum 1 Libanici de Libanici ad agrum

Diplazium longum Triplici, polchi, Marone

Diplazium in Libanici. agrum ad Marone

Diplazium polytrichum L.

Diplazium unguiculatum abique in Libanici

Diplazium longum Marone

Diplazium ~~complanatum~~ Marone

Diplazium ~~complanatum~~ Marone

Diplazium ~~complanatum~~ Marone

Diplazium ~~complanatum~~ Marone

Diplazium ~~complanatum~~ Marone

7
u

[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]

Article of the [illegible] of [illegible]

[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]

[Small handwritten mark or signature at the bottom right corner.]

varmisaam - sama sauto

Valerian offic Junonia piper

Witum Jamus Brygg Almq. ibid.

Coposikal

Verbasum Lychwitz vuly - in litt.

vigvum in sybentij dula. mont.

Botanische Sammlungen

[Faint, illegible handwriting]

Capitulum

[Faint, illegible handwriting]

Launth vinnige Vög-
Lanthier No. 12

