

Compendij ordinum Florae Delmaticae

Classis I. Cryptogamae.

- Ordo I. Characeae  
II. Equisitaceae  
III. Lycopodiaceae  
IV. Filices.

Classis II. Phanerogamae  
Monocotyledones

- Ordo V. Graminae  
VI. Cyperoideae  
VII. Funceae  
VIII. Irideae  
IX. Amaryllidaceae  
X. Liliaceae  
XI. Colchicaceae  
XII. Smilacaceae  
XIII. Dioscoreae  
XIV. Orchidoideae  
XV. Callaceae  
XVI. Typhaceae  
XVII. Najadaceae  
XVIII. Alismaceae  
XIX. Butomaceae  
XX. Hydrocharitideae

Classis III. Phanerogamae  
Dicotyledones

Subclass. I. Chlamydoblastae

- Ordo XXI. Asarineae  
XXII. Cytineae  
XXIII. Nymphaeaceae

Subclass. II. Gymnoblastae

Legio I. Monochlamydeae

- Ordo XXIV. Coniferae  
XXV. Amretaceae  
XXVI. Salicinae  
XXVII. Juglandaceae  
XXVIII. Urticeae  
XXIX. Santalaceae  
XXX. Thymelaeae  
XXXI. Laurinae  
XXXII. Polygonaceae  
XXXIII. Phytolacaeae  
XXXIV. Chenopodiaceae  
XXXV. Amaranthaceae  
XXXVI. Scleranthaceae

Leg. II. Dichlamydeae

Cohors I. Monopetalae

- Ordo XXXVII. Plantagineae  
XXXVIII. Plumbagineae  
XXXIX. Globulariae  
XL. Dipsacaceae  
XLI. Valerianaceae  
XLII. Compositae  
XLIII. Campanulaceae  
XLIV. Vacciniae  
XLV. Ericinae  
XLVI. Styracaceae  
XLVII. Ebenaceae  
XLVIII. Samoliniae

XLIX. Primulaceae -

- I. Lentibulariæ
- II. Scrophulariæ -
- III. Boraginæ -
- IV. Verbenacæ
- V. Labiatæ -
- VI. Acanthaceæ
- VII. Convolvulaceæ
- VIII. Cuscutæ
- IX. Solanaceæ -
- X. Boraginæ -
- XI. Gentianæ
- XII. Asclepiadæ -
- XIII. Apocynæ
- XIV. Rubicæ -
- XV. Caprifoliacæ
- XVI. Jasmineæ -

Coh. II. Polypetalæ

Ordo LXVI. Loranthæ

- LXVII. Umbellatæ -
- LXVIII. Hederaeæ
- LXIX. Berberidæ
- LXX. Paeoniæ
- LXXI. Ranunculæ -
- LXXII. Polygalæ
- LXXIII. Resedæ
- LXXIV. Fumariacæ
- LXXV. Papaveræ
- LXXVI. Cruciferæ -
- LXXVII. Capparidæ

LXXXVIII. Cucurbitaceæ

- LXXXIX. Graminæ
- LXXXX. Nopaleæ
- LXXXI. Astinæ -
- LXXXII. Violæ -
- LXXXIII. Damariscinæ -
- LXXXIV. Hyacinthæ
- LXXXV. Paronychiæ
- LXXXVI. Potulacæ
- LXXXVII. Caryophyllæ -
- ~~LXXXVIII. Graminæ~~
- ~~LXXXIX. Saxifragæ~~

XC. LXXXIX. Halimogæ -

- XCI. LXXXX. Lythariæ
- XCII. Onagriae -
- XCIII. Granatæ
- XCIV. Myrtacæ
- XCIV. Tiliacæ
- XCVI. Malvacæ -
- XCVII. Geraniacæ -
- XCVIII. Linæ -
- XCIX. Oxalidæ

~~XCIX~~ Samentacæ

- CI. Meliacæ *Malpighiæ*
- CII. Acerinæ - *Hydrocotylæ*
- CIII. Euphobiæ -
- CIV. Rhamnæ -
- CV. Aquifoliæ
- CVI. Celastrinæ
- CVII. Dichamnæ
- CVIII. Rutæ
- CIX. Zygophyllæ
- CIX. Cassiæ
- CXI. Rosacæ -

- CXII Pomaceae  
CXIII Bryadeae  
CXIV Spireaceae  
CXV Amygdaleae  
CXVI Leguminosae. —
-



- |   |  |  |
|---|--|--|
| 14. Glumij compressij - - - 15                  | Glumijis ventricosij                                 |  |
| 15. Glumellij basi longe pilosij - 16           | <u>Milium</u> XXIX                                   |  |
| 16. Anistij glumella multa longiorij            | Glumellij basi pubescentiorij                        |  |
| <u>Stipa</u> XXX                                | <u>Agrostis</u> XXVIII                               |  |
| 17. Spiculis spicatis 18                        | Anistis glumellam vix excedentibus                   |  |
| 18. Spiculis omnibus nudis 19                   | <u>Calamagrostis</u> XXVII                           |  |
| 19. Gluma bivalvis 20                           | Spiculis paniculatis - - 22                          |  |
| 20. Valisij linearibus acuminatis-<br>aristatis | Spiculis involucentis - - - 24                       |  |
| <u>Secale</u> LII.                              | Gluma univalvis                                      |  |
| 21. Omnibus involucentis pinnatis               | <u>Lolium</u> LIIV.                                  |  |
| <u>Cynodorus</u> XLVIII.                        | Valisij lanceolatis muticis vel abrupte<br>aristatis |  |
| 22. Spiculis subtrifloris 23                    | <u>Triticum</u> LV.                                  |  |
| 23. Spiculis oblongis 24                        | Infirmis bractea ovata                               |  |
| 24. Spiculis ovatis 25                          | <u>Sesleria</u> XLIX                                 |  |
| 25. Floribus omnibus fertilibus                 | Spiculis multifloris - - 26                          |  |
| <u>Aira</u> XXXVIII                             | Spiculis orbiculatis                                 |  |
| 26. Glumellij aristatis acuminatisve 27         | <u>Beckmannia</u> L.                                 |  |
| 27. Paniculis coarctatis 28                     | Spiculis lanceolatis                                 |  |
| 28. Panicula aequali                            | <u>Avena</u> XXXIX                                   |  |
| <u>Koeleria</u> XI.                             | Flora terminali sterilibus                           |  |
| 29. Glumellij apice aristatis acuminatisve      | <u>Melica</u> XXXVII                                 |  |
| <u>Festuca</u> XLII                             | Glumellij muticis 30                                 |  |
| 30. Glumellij subovatis - - 31                  | Paniculis laxis - - 29                               |  |
| 31. Spiculis ovatis elongis 32                  | Panicula secunda                                     |  |
| <u>Poa</u> XLV.                                 | <u>Dactylis</u> XLIII                                |  |
| 32. Glumella glabra                             | Glumellij sub apice aristatis                        |  |
| <u>Glyceria</u> XLVI.                           | <u>Bromus</u> XLI.                                   |  |
| 33. Floribus nudis                              | Glumellij oblongis - - - 30                          |  |
| <u>Antheranthum</u> XVI.                        | Spiculis late subcordatis                            |  |
|   | <u>Briza</u> XLVII.                                  |  |
|   | Glumella basi lanata                                 |  |
|   | <u>Lorundo</u> XLIV.                                 |  |
|   | Floribus monandris                                   |  |
|   | <u>Ptilurus</u> LII.                                 |  |

- |    |  |    |   |
|----|--|----|---|
| 34 | <i>Horibus polygamis</i>                                   | 35 | <i>Horibus monois</i>                                   |
| 35 | <i>Spiculis involucri</i> <sup>ca. XV.</sup>               | 36 | <i>Spiculis nudis</i> <sup>ca. XXI.</sup>               |
| 36 | <i>Gumella nutica</i><br><u>Setaria</u> XXXII              |    | <i>Gumella aristata</i><br><u>Hordeum</u> LVII.         |
| 37 | <i>Gumij ibactis</i>                                       | 38 | <i>Gumij exhy echinatis</i>                             |
| 38 | <i>Spiculis uni-bifloris</i>                               | 39 | <i>Lappago</i> XXVI<br><i>Spiculis tri-glandifloris</i> |
| 39 | <i>Gumellis quibusdam aristatis</i>                        | 40 | <i>Aegilops</i> III<br><i>Gumellis omnibus nuticis</i>  |
| 40 | <i>Gumella maris aristata</i>                              | 41 | <u>Lonicum</u> XXXI<br><i>Gumella maris nutica</i>      |
| 41 | <i>Spiculis geminatis unifloris</i><br><u>Pollinia</u> XIX |    | <i>Spiculis sparsis bifloris</i><br><u>Holcus</u> XVII  |
| 42 | <i>Spiculis digitato-spicatis</i><br><u>Andropogon</u> XX  |    | <i>Spiculis paniculatis</i><br><u>Sorghum</u> XVIII     |

- |   |   |
|---|---|
| 32. <i>Plumella glabra</i><br><u>Glyceria</u> XLVII.              | <i>Plumella basi lanata</i><br><u>Arundo</u> XLIV                 |
| 33. <i>Flosculo hermaphrodito triandro</i> - - - 36               | <i>Flosculo hermaphrodito diandro</i><br><u>Arthroxanthum</u> XVI |
| 33. <i>Hortus monosicy</i> - - - 34                               | <i>Hortus polygamus</i> - - - 35                                  |
| 34. <i>Hortus masculis paniculatis</i><br><u>Lea</u> XV.          | <i>Hortus omnibus spicatis</i><br><u>Heteropogon</u> XXI          |
| 36. <i>Spiculis involucreatis</i> - - - 37                        | <i>Spiculis nudis</i> - - - 38                                    |
| 37. <i>Plumella mutica</i><br><u>Setaria</u> XXXII                | <i>Plumella aristata</i><br><u>Hordeum</u> LVIII                  |
| 38. <i>Plumis inermibus</i> - - - 39                              | <i>Plumis extus echinatis</i><br><u>Lappago</u> XXVI              |
| 39. <i>Spiculis uni-bifloris</i> - - - 40                         | <i>Spiculis tri-quadrifloris</i><br><u>Astilops</u> LIV           |
| 40. <i>Plumellis <sup>quibusdam</sup> aristatis</i> - - - 41      | <i>Plumellis omnibus muticis</i><br><u>Panicum</u> XXXI           |
| 41. <i>Plumella maris aristata</i> - - - 42                       | <i>Plumella maris mutica</i> - - - 43                             |
| 42. <i>Spiculis geminatis unifloris</i><br><u>Pollinia</u> XIX    | <i>Spiculis sparsis bifloris</i><br><u>Holcus</u> XVII            |
| 43. <i>Spiculis digitatis - spicatis</i><br><u>Andropogon</u> XX. | <i>Spiculis paniculatis</i><br><u>Sorghum</u> XVIII.              |

XV Lea I

Sen. pl. d. II p. 341.

Ch. *Flora monosicy*: masculis racemosis-paniculatis; spiculis bifloris, plumis glaberrimis bivalvatis; foemineis spicatis sessilibus confertim axi spadiciformi bracteis vaginatis impositis; spiculis unifloris, plumis plumelliguis bivalvatis, stylo longissimo filiformi pendulo glabro. Caryopsij subtendens pedicellata foveae axij adhaerens. — Cl. Monoclea Ord. Triandria Lyth. 100

34. Lea Mays L. sp. pl. p. 1133. Vis. It. p. 51. Reichb. fl. ex. p. 54  
Alsch. fl. jedr. p. 17.

L. foliis integerrimis, margine rubris.

Icon. Duchs. hist. pl. p. 825

Blackw. herb. 2. tab. 547

Lam. Enc. bot. Pl. 149

Matth. Disc. ed. Valgr. p. 117

Myr. Kükürur, Gölkud.

var.  $\beta$ . gracex : minor

Myr. Berzohod, Berzohodar.

Hab. in America meridionali, colitur vero in oeconomia usque ubique in Dalmatia praesertim montana. H. Julii, Augusto. Caryopisid occurrunt albae, lacteae, nigrae, et ex his coloribus variatae.  $\odot$

Ufus Caryopisid hujus plantae in primum molitur, et ex aqua coctae nutriendi alimentum praebent agriis. Soli libenter affumentur a quadrupedibus.

### XVI Anthoxanthum L.

Gen. pl. ed. II p. 14

Ch. Flores polygami paniculato-spicati. Stamina bivalvia triflora, filamentis latensibus nec trij. univalvibus inferius basi, superioribus sub apice aristati, medio hermaphroditi & andro, bivalvia, multia. <sup>Stigma flavo-</sup> A. Diandria. Ord. Digynia Syst. ix.

35. Anthoxanthum odoratum L. sp. pl. p. 40. Vis. Herp. Dalm. p. 25. Host fl. austr. p. 65. Reichb. fl. exc. p. 55. Alsch. fl. jard. p. 8. Retz. bot. wegw. n. 24.

A. panicula oblonga spiciformi, filamentis subsessilibus aristis longioribus, filamentis exterioribus carina ciliatis, vaginis ova barbatis foliisque glabris.

Icon. Host Ic. gram. austr. 1. tab. 8.

Lam. Enc. bot. Pl. 23

Engl. bot. t. 647

Reichb. Ic. pl. rar. XI. tab. 106 f. 1822-23

Hab. in herbis, collibus, dumis totius Dalmatiae copiose. Flor. Agati Majis.  $\odot$ .  $\text{S}$ .

Obs. Spica flavifera apice attenuata. Differt ab affini A. gracili Bivon. Herp. sic. man. 1. p. 3 panicula longiore, filamentis ciliatis, aristis floris brevioribus, foliis glabris, dum illud paniculam gerit fore ovatum, filamentis glabris, aristis praesertim florum superiorum eodem sequentibus, folia pilosa.

### XVII Holcus ef.

Ovdo VI. Cyperoidae

Russ. gen. pl. p. 26

Floris glumacis hermaphroditis aut unisexuales, spicati et tunc singuli sessiles in axilla  
(Penyjnium) ~~bracteolae~~ <sup>bracteolae</sup> ~~vel~~ <sup>glumae</sup>, raro solitarii in gluma bracteoli aut multivalvuli. Perianthium  
(flores vidi) vel nullum vel utrinque apice perovium, vel eschij hypogynis conflatum. Stamina  
tria, raro unum vel duo, hypogyna, ~~antheris~~ <sup>antheris</sup> erectis connectivis distinctis, stylus  
unius stigmatibus binis trisive interdum bifidis, ovum libero uniovulato. Larva  
rara saepe trigona, raro complanata, quibusdam styli basi rostrata. Semen liberum,  
albumina pinnacea, embryone in basi axis albuminis parte, centico.

Herbae annuae aut rhizomate perennantes aut heteropae, culms simplicissimis, basi  
foliato aut vix foliato, raro ramoso foliato, folijs altheris simplicibus viximantibus,  
ligula nulla. Spicae solitariae, aut paniculatae aut umbellatae aut  
spicatae aut capitatae, saepe bracteatae.

Clavis analytica generum

- |   |  |
|---|--|
| 1. Floribus hermaphroditis vid. No <sup>m</sup> 2   | Floribus unisexualibus   |
|   | <u>Carex</u>   |
| 2. Staminibus distinctis . . . . . 3  | Staminibus unisive imbricatis . . . 5  |
| 3. Perianthio nullo . . . . . 4   | Perianthio setoso  |
| 4. <sup>Stigmatibus</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> inferioribus distinctis | <sup>Blugnes</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> inferioribus distinctis, brevissimius |
| <sup>Stigmatibus</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> inferioribus unisive       | <sup>Stigmatibus</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> inferioribus unisive              |
| <sup>Stigmatibus</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> inferioribus unisive       | <sup>Stigmatibus</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> inferioribus unisive              |
| <sup>Stigmatibus</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> inferioribus unisive       | <sup>Stigmatibus</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> inferioribus unisive              |
| 5. Carexula sicca . . . . . 6   | Carexula dracopaea   |
|   | <u>Cladium</u>   |
| 6. Stylo basi bulbosa persispirata  | 6. Stylo basi filiformi deciduo . . .  |

- |   |   |
|---|---|
| 7. <u>Heterochaeris</u>   |   |
| 7. Perianthio nullo . . . . . 8   | <u>Carex</u> L.   |
| 8. <del>Stigmatibus</del> <sup>Stigmatibus</sup> bifidis . . . 9            | Stylis <sup>bifidis</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> bifidis       |
| 9. Floribus <del>Stigmatibus</del> <sup>Stigmatibus</sup> unisive . . . . . | Floribus <sup>Stigmatibus</sup> <del>Stigmatibus</del> <sup>Stigmatibus</sup> unisive |
|   | <u>Heterochaeris</u>  |

Ch. Floris monici spicati imbricati, singuli bracteola suffulti. Perianthium utrinque  
apice perovium persistens. Stamina tria. Stigma binum vel ternum. Carexula perianthio test.  
- Cl. Monocia Ovd. Jurae N. Syst. ex. -

\* Stigmatibus binis, carexula complanata  
Vigrea Beauv. Reichb. fl. ex. p. 55. Nees Gen. pl.  
p. 111.

+ *Carex divulsa* Good. trans. of Linn. Soc. II, p. 60 Hort fl. austr. 2. p. 582 Alph. fl. jav. p. 221. Pfl. bot. wegw. n. 204.

*C. spica suberosa laxa elongata, spiculis androgynis sessilibus ovatis alterius superne masculis, superioribus approximatis, inferioribus bracteis elongatis angustifolius, fructibus ovatis plano-convexis nervis bidentatis lacinis marginatis crebris, calicibus bracteatis ovatis acutis.*

Syn. *Vignea divulsa* Reichb. fl. ex. p. 59

Icon. Schk. hist. des car. tab. 33 et. W. W. n. 89.

Michx. nov. gen. pl. t. 53 f. 10

Engl. bot. t. 629.

Hab. ad ripas et locis umbratis circa Zava, Selenice, Dernij, Palato, Ragusa, et in insula Lerina. Spiculae flavae-virescentes, aetate virides. Fl. major 2.

+ *C. mucronata* L. sp. pl. 2. p. 1582 Hort fl. austr. 2. p. 581

*C. spica subcylindrica, spiculis androgynis sessilibus ovatis compressis superne masculis, fructibus stellatis divergentibus plano-convexis ovato-acuminatis bidentatis obsolete nervosis lobato-nervatis, bracteatis ovatis cuspidatis.*

Syn. *Vignea mucronata* Reichb. fl. ex. p. 57. Nees loc. cit.

Icon. Hort Scav. gr. austr. 1. t. 54.

Engl. bot. t. 1092

Nees loc. cit. p. 1092.

Schreb. Agrost. t. XI. f. 5.

Hab. in praealpibus et ad ripas circa Schura et Dernij. Spiculae cum bracteis ferrugineae.

Fl. Aprilis, Majo: 2.

Obs. *C. divulsa*, *mucronata* et *viridis* tam parvis vix, nec constantibus, ab invicem distinguuntur, ut vixim unam eandemque speciem constituant viderentur. <sup>et proinde C. divulsa</sup> Hort Scav. gr. austr. 1. t. 54. Reichb. Agrost. t. XI. f. 5. Nees loc. cit. p. 1092.

+ *C. vulpina* L. sp. pl. 2. p. 1582 Hort fl. austr. 2. p. 583 Alph. fl. jav. p. 221. Pfl. bot. wegw. n. 310.

*C. spica oblonga ramosa compacta, spiculis androgynis sessilibus ovatis compressis superne masculis, fructibus patentibus compresso-triangularibus ovato-acuminatis bidentatis nervosis, margine apicis scabris, bracteatis ovatis cuspidatis, ulmi acutis membranaceis.*

Syn. *Vignea vulpina* Reichb. fl. ex. p. 59

Icon. Hort Scav. gr. austr. I. t. 56.

Engl. bot. t. 309

Schk. hist. des car. tab. C. f. 10.

Less. fl. herb. t. 14. f. 15.

Hab. ad fontium marginem et ripas humidas terrae Dalmaticae. Fl. Apr. Majo. Spiculae ferrugineae. 2.

+ *C. diffusa* Good. trans. of Ann. Soc. II p. 157 Hort. fl. austr. 2. p. 576. Alsch. fl. jard. p. 221.  
*C. spica abbreviata* congesta, spiculis androgynis sessilibus oblongis alternis congestis superne  
 nupulif., basi bracteatis, fructibus erectis ovatis convexo-concavis identibus nervosis  
 subo-marginatis, bracteatis ovatis mucronatis.

Syn. *C. schenoides* Hort. fl. austr. 2. p. 576. Le. gr. 1. p. 55

Rymer Disser. Reichb. fl. exc. p. 58

Leon. Schk. hist. des lar. tab. 3 n. 18

Engl. bot. tab. 1096

Hort. le. gr. austr. 1. tab. 48

Hab. in arenis siccis in insula B. 1100, et circa Lava, Derain, Salona, Nagusa. Spicularum  
ferugineae. Del. Meje, Juni. Phytoma articulatione rugosa. 4.

\* Obs. Differentia adducta a cl. Koch in syn. fl. helv. et germ. p. 880 *C. schenoides* a  
*C. diffusa* minime diffingunt, nam fructus in utroque ovatis et nervosis esse ex ipsius  
Hookii descriptione et iconis leuclenter evincitur.

\*\* Stigmatis terminis, cavacula subtrigona

Cavex Beauv. Reichb. fl. exc. p. 61. Nees Sens. p. 1014.

+ *C. Linckii* Schk. hist. des lar. II. q. pl. 11. p. 223 (1805) Reichb. fl. exc. p. 62.

Alsch. fl. jard. p. 221. Pitt. bot. meyv. n. 200 excl. syn.

*C. spici* alternis subternis diplobis androgynis sessilibus linearibus superne nupulif.,  
 infime pedunculata, fructibus ovatis-triquetris binerviis liguibus, utrinque  
 acuminatis apice <sup>nupulif.</sup> bracteatis oblongo-lanceolatis in fructu gubulif., bractea  
 basilari foliacea.

Syn. *C. gymnomane* Benth. ex. pl. stud. Dec II, p. 43 (1806.)

*C. Klumbp* Lois. fl. gall. II. p. 629

Leon. Schk. loc. cit. t. 366 f. 117-118.

Hab. in collibus, et graminosis circa Lebenio, Nagusa, et in montibus Villedith  
 et Bischor. Doret Ag. Meje. Spicularum bracteatis margine ferrugineis. 4.

Obs. Nomen Schleckerii primitivis jure ceteris anteprecedendum.

F# *C. alpestris* Allion. fl. ped. II. p. 270 Reichb. fl. exc. p. 65 Alsch. fl. jard. p. 222

*C. spica* mascula solitaria testis pedunculata, feminis terminis b. quinquepartitis, binis  
 approximatis subsessilibus, infime radicali longipennis pedunculata, fructibus ovatis-  
 triquetris apice inopulatis identibus multinerviis liguibus, bracteatis ovatis mucro-  
 natis, bractea basilari laterea.

Syn. *C. diversiflora* Hort. ic. gr. austr. 1. p. 53 fl. austr. 2. p. 573

*C. gymnotris* Willd. hist. des pl. de Dauph. 2. p. 206 Schk. hist. des lar. p. 92

Leon. Hort. ic. gr. austr. 1. t. 70

Schk. loc. cit. tab. 9. n. 35.

Hab. in collibus et locis agris inanis. Chevo, circa Zara, Sebenico et Salona. Hort.  
Agr. Major. Spicae dilute ferrugineae. 24.

+ *C. digitata* L. sp. pl. p. 1384. Hort fl. austr. 2. p. 587. Reichb. fl. ex. p. 83

*C. spica* mascula tereti. Stiparia, feminisq; sublongis pedunculatis linearibus demum  
diffunditibus, fructibus ovatis - trigonis aequis emarginatis brevibus puberulis  
nitidij, bracteis ovatis - truncatis, bractea basilari gathypomii margine membranacea.

Icon. Hort. Ic. gr. austr. 1. t. 60

Engl. bot. t. 615

Schk. l. c. tab. H. n. 78

Michel. nov. gen. pl. t. 92. f. 9

Leesl. fl. herbom. t. XVI. f. 4.

Hab. in sylvestrij montium circa Verulica. Hort. Martii, Agypti. Spicae pallide ferrugineae.  
x.

Obs. Spica mascula scilicet proxima feminis pedunculata brevis est, hinc potius la-  
teatly quam terminalis videtur. *C. ornithopoda* W. ab hac aequa et vix quidem  
distinguitur.

+ *C. humilis* Leysse fl. hol. p. 255 Hort fl. austr. 2. p. 585. Reichb. fl. ex. p. 83

*C. spica* mascula oblonga solitaria pedunculata, feminisq; tenuis vix subtrifloris  
inclusis breviterq; pedunculatis, fructibus ovatis - trigonis longis trisulcatis brevibus  
sine restantij subuliginis, bracteis canisq; oblongis obtusis, inferiortibus majoribus,  
bractea basilari vaginante setacea - mucronata oblonga, margine membranacea.

Syn. *C. clandestina* Gooden. Trans. of Linn. soc. II. p. 167. W. sp. pl. IV. p. 254

Schk. hyp. des Car. p. 194.

Icon. Hort. Ic. gr. austr. 1. t. 67

Engl. bot. t. 2124

Schk. l. c. tab. K. n. 43

+ Michel. nov. gen. pl. xxxii f. 8.

Hab. in herbisq; montis Diskara ubi legit Dommasini. Pl. Major. Spicae ex alto et ferrugineae vix  
24.

+ *C. ovata* Vill. hist. des pl. de Dauph. 2. p. 204 (nec Schk.) Reichb. bot. weym. n. 209.

*C. spica* mascula subclavata pedunculata solitaria, feminisq; geminis approximatis oblongis  
subsessilibus, fructibus ovatis - trigonis puberulisq; integris, bracteis ovatis acutis, bractea  
basilari setacea basi dilatata vaginante variosa.

Syn. *C. procerx* Jaeg. fl. austr. t. 446. Schk. l. c. p. 67 Hort fl. austr. 2. p. 591. Reichb. l. c. p. 84. Albid.

*C. filiformis* Leesl. fl. herb. p. 204 (nec L.) fl. p. 222.

Icon. Hort. Ic. gr. austr. 1. t. 68.

Engl. bot. t. 1099.

Schk. l. c. tab. 7. n. 27

Leesl. l. c. tab. XVI. f. 5.

Hab. in pratis et collibus circa Zara, Sebenico, Spalato, Salona, Cliffa et Dagusa. Pl. Martii, Agr. Spicae  
ferrugineae. 24.

228. *C. muricata* L. sp. p. 127. 1389. Hort. Fl. austr. 2. p. 581.

*C. spica* sub cylindrica, spiculis androgynis sepilibus ovalis congestis superne masculis, fructibus stellatim divergentibus plano - convexis ovato - acuminatis bidentatis obsolete nervosis utraque surculis ~~marginatis~~, bracteolis ovalis cuspidatis.

Syn. *Signea muricata* Reichb. f. ex. 1. p. 87.

Hort. Hort. Fl. austr. 1. p. 58.

Engl. bot. 4. 1097.

Lehmann. Agricul. 4. XI. p. 5.

Hab. in pratinis et ad ripas inna Lebenico et Dernid. Spicula virides.  
Florib. Aprilis, Mayo.

229. *C. vulpina* L. sp. p. 127. 1389. Hort. Fl. austr. 1. p. 580. Alach. Fl. jard. p. 221. Lehmann. bot. wegm. 21. 210.

*C. spica* oblonga remota congesta, spiculis androgynis sepilibus ovalis congestis superne masculis, fructibus potentillae compressa triquetris ovato - acuminatis bidentatis nervosis, superne margine scabris, bracteolis ovalis cuspidatis, utrinque triquetris angulis membranaceis.

Syn. *Signea vulpina* Reichb. f. ex. 1. p. 89.

Hort. Hort. Fl. austr. 1. p. 56.

Engl. bot. 4. 207.

Lehmann. hist. des Lav. p. 6. p. 10.

Lerch. Fl. herbarn. 4. 14. p. 15.

Hab. ad fossarum margines et sepes humidos in tota Darmatia frequens. Florib. Aprilis, Mayo. Spicula ferruginea.

~~228~~ *Sigmetibus* tenuis, caruncula subtriangula.

Gynemans Det. fl. ital. dec. II d. 1. p. 43 (1806)

220. C. Linkii ~~subtus~~ <sup>Hist. de lar.</sup> ~~Reichb. Bot. Voy. Sicil.~~ Willd. sp. pl. IV p. 228. Reichb. fl. ex. 1. p. 62. Alach fl. jod. p. 221. Cell. bot. voye n. 208 (eod. synon.)

C. spicae alternis subternis distichis androgynis sepalibus lineari-  
ribus, infima pedunculata, sursum masculis, fructibus ovato-  
triangulis bimeris utrinque auriculis levibus apice subfili-  
tis, bracteolis oblongo-lanceolatis vel fructa parvis, bractea  
basilari solitaria.

Syn. C. <sup>Linkii Schk. 1872. in lar. II p. 10. p. 223. Fuchs fl. ex. p. 62. g.</sup> ~~Gynemans Det. fl. ital. dec. II d. 1. p. 43.~~  
C. huberana Loisel. fl. gall. II p. 629.

Non. Schk. ~~1872. in lar. II p. 10. p. 223.~~ 1872. in lar. II p. 10. p. 13.

Hab. in collibus circa Albenico Loisel. multif. et in montibus Velle  
bi et Albenico et Loisel aprilis Mayo. Sp. uide. Deluche ferru-  
gineo. 28

231. C. alpestris Allion fl. pedum. II p. 210. Reichb. fl. ex. 1. p. 65.  
Alach. fl. jod. p. 222.

C. spica mascula solitaria tereti pedunculata, foecundis tri-  
bus si-quinquefloris, binis approximatis, sepalibus, in-  
fima radicali longissime pedunculata, fructibus obova-  
to-triangulis apice inequaliter bidentatis multinerviis  
levibus, bracteolis ovalibus micromatis, bractea basilari  
solitaria.

Syn. C. divorsiflora Hort. Ic. p. austr. 1. p. 53. Fl. acastr. 2. p. 593.

C. gynobasis Willd. hist. Bot. pl. Du. Dauph. 2. p. 206. Schk. 1872. in lar. p. 92

Non. Hort. Reichb. anstr. I. p. 70.  
Schk. 6. ut. no. Gen. 35.

Hoab. in calibus et locis apertis in valle Cherso, circa Kara, Sebenico, et  
Salona. Flores aperti, nigri. Spica vitata ferruginea. 2.

232. C. digitata Less. sp. pl. 2. p. 1234. Hort fl. austr. 2. p. 587. Reichb.  
fl. ex. 1. p. 63. ~~alt. l. 1. p. 63.~~

C. spica mascula tereti sepius solitaria, foemineis subternis pedunculatis  
linearibus demum distantibus, fructibus obovato-triquetris  
apice emarginatis binerviatis pubescentibus nictidis, bracteis  
obovato-truncatis, bractea capulari spathe formi margine anem  
bracteata.

Icon. Hort Sc. gram. austr. v. 1. t. 60.

Engl. bot. t. 618.

alt. l. 1. p. 63.

Michx. nov. gen. pl. 4. 32. f. 9.

Less. fl. herbarn. 4. XII. f. 4.

Hoab. in sylvaticis montium prope Karliha. Flores masculi, aperti. Spica  
caerulea ferruginea. 2.

Abt. Spica mascula sepius proxima foeminea pedunculata brevior est,  
hinc spiculis lateralibus quam terminalibus videtur, ~~pedunculata~~  
~~sepius~~, C. ornithoglossa 78, ab hac acriter et vix quidem dif-  
tinguitur.

233. C. humilis Less. fl. nat. <sup>p. 235</sup> Hort fl. austr. 2. p. 588. Reichb. fl.  
ex. 1. p. 63.

C. spica mascula oblonga solitaria pedunculata, foemineis  
ternis remotis subtropis imilise breviterque pedunculatis,  
fructibus obovato-triquetris nervatis tribracteis brevissimis  
restratis subintegris, bracteis variis oblongis obtusis imperio

ribus majoribus, bracteis capsulari vaginante. lobis mucronata ob-  
tusis, margine membranaceo.

Syn. *C. clandestina* Loerd. Trans. of Linn. Soc. II. p. 167. Willd. sp. pl. IV  
p. 251. Schk. hist. des Car. p. 104.

Non. Hook. Fl. gr. austr. I. t. 67.  
Engl. bot. t. 2124.

Scheuchz. agrar. I. x. f. 1.

Nichol. nov. gen. pl. t. 32. f. 6. Schk. l. c. tab. K. n. 43.

Herb. in herbicid. montis Piskova. D. Tommasini. Floret. major. spicae  
caeruleae et ferrugineae varieg. 28.

- 234. C. ovina Vill. hist. Des. pl. du Dauph. 2. p. 204. (non Scheuchz.) Pell.  
bot. nigr. n. 209.

*C. spica* majuscula subclavata pedunculata solitaria, feminis et ge-  
minis approximatis oblongis subsessilibus, fructibus obovatis tri-  
gonis pubescentibus integris, bracteis ovatis acutis bractea caps-  
ulari lobata basi dilatata vaginante scariosa.

Syn. *C. praecox* Engl. Fl. austr. I. tab. Hook. Fl. austr. 2. p. 571. Richard.  
fl. ex. I. p. 64. Alsch. Fl. jud. p. 142.

*C. filiformis* Leers Fl. herbarn. p. 204 (non Linn.)

Non. Engl. Fl. austr. p. 114.

Hook. Fl. gr. austr. I. t. 68.

Engl. bot. t. 1099.

Schk. l. c. tab. F. p. 27

Leers Fl. herbarn. t. XXI. f. 5.

Herb. in pratib. et collibus circa Nara, Dernis, Sabinico, Spalato, Salona,  
Ulcipia, et circa ambula prope Ragusa. Floret. et Martio, a grile.

*Blysmus* Panz.

in K. et Sch. syst. veg. mant. 2. p. 41.

Char. Flores hermaphroditi bracteolati in spicam compositam terminalem disticha imbricati, spiculis multifloris bracteis suffultis. Perianthium e sechis hyggyis subseris. Stamina tria. Stylus filiformis bipartitus. Caruncula in styli apicem tunc articulatione attenuata. — Cl. Triandria Ord. Monogynia Syst. 100.

*Blysmus compressus* Panz. l.c. Reichb. fl. ex. p. 24.

B. culms angulatis, spiculis 6-8 floris, foliis planiusculis subter carinatis

Syn. *Carex uliginosa* L. sp. pl. 1381. et Mant. alt. p. 494

*Schoenus compressus* L. l.c. 65 et Mant. alt. p. 511

*Saxifraga compressa* Pers. Syn. pl. 1. p. 66

*S. canis* Metz. prod. p. 16 W. sp. pl. 1. p. 292 Host Ic. gr. austr. 2. p. 39 fl. austr.

1. p. 55.

Panz. Host Ic. gr. III. t. 57

Flor. dan. tab. 1622

Nees Sem. pl. fasc. IX Blysmus fig 1-12.

~~decc fl. herb. t. 1. p. 1.~~

Hab. in pascuis montium Vellebit. Fl. Suis, Aug. spica castanea. 24.

Obs. Differt ab affini *B. v. v.* Lh. charactibus sechis, et sechis perianthii venterum aequalis, nec erecto-pubescens, ut ventera venteris cl. Koch syn. fl. germ. et helv. 2. p. 244.

*Schoenus* ~~L.~~ Beauv.

in Less. Ess. sur les Cyper.

~~fl. p. 20 (ext. spec.)~~

Char. Flores hermaphroditi in spicam terminalem compositam bracteis albis suffultis capitato-congestis, spiculis uni-gamistis bracteis distichis, inferiorum planiusculis minoribus sterilibus. Perianthium nullum. Stamina tria. Stylus basi teres deciduus, signatim teris. Caruncula <sup>in styli apicem</sup> subseris. — Cl. Triandria Ord. Monogynia Syst. 100.

*Schoenus nigricans* L. sp. pl. p. 64 Host fl. austr. 1. p. 50 Reichb. fl. ex. p. 24 alt. fl. jav. p. 16.

*S. foliis filiformibus*, spica ovata bracteis binis erectis, alba spicam opposita, involucris.

Loc. Host Ic. gr. austr. III. tab. 54

Engl. bot. t. 1121

Lam. Enc. bot. Pl. 38 f. 1.

Nees Sem. pl. fasc. IX (Schoenus) fig. 1-14

Hab. in arenosis et saxosis montibus tum continenter tum insularum. Fl. Maj. Jun. spica nigra.

177

+ *S. mucronata* L. sp. pl. p. 63. Hort. fl. austr. I. p. 49-50. Fiehb. fl. exc. p. 74.

*S.* filij planij, spica haemisphaerica capitata, bractij 2-6 palmatifidij spiculae multae  
evidentibus involucentis.

Icon. Hort. Ic. gr. austr. IV. t. 70

Sibth. fl. gr. Cent. I. t. 40

Schemb. agrostogr. tab. 8. f. 1.

Nes. Sen. pl. p. 18. cit. f. 15-16.

Hab. in arenis montium prope Nana. Prof. Alshinger. Flor. Tunis, Julia. Flumina  
= rubra spicibus marginibus pallidioribus.

### *Cyperus* L.

Gen. pl. ed. II. p. 20.

Chin. filij hemisphaerica, in spica bractij radiantibus involucentibus digitatis, spiculae multae  
filij compressae, bractej apice imbricatij, infimij (tenuj) similibus. ~~Paniculae~~  
nullae. Stamina 2-3. Stylus bi-trifidus basi tenui deciduo. Caracula conflua  
- *C. Diandria* *Del. Monophyllum* Syd. ex.

+ *Cyperus flavescens* L. sp. pl. p. 65. Vj. sing. detm. p. 35. Hort. fl. austr. I. p. 52. Alsh.  
fl. exc. p. 17.

*C. stylo bifido*, spiculae subcompositae umbellatij involucentibus triphyllis brevioribus, spiculae  
compressae oblongae, bractej erectae obtusae, filij angustissimij & viciae fibrae

Syn. *Cyperus flavescens* Beauv. Fiehb. fl. exc. p. 72

Icon. Hort. Ic. gr. austr. III. tab. 72

Sibth. fl. gr. Cent. I. tab. 47

Koch. fl. austr. t. 21

Lam. Enc. bot. Pl. 98. f. 1.

Hab. in prope ~~littoribus~~ palustribus circa Zam, Zuvij, Salona, Budua. Florae  
Juliae, Septembris. Spiculae flavescens, demum bridae. ©

+ *C. pusillus* L. sp. pl. p. 69. Hort. fl. austr. I. p. 52. Fiehb. fl. exc. I. p. 72. Alsh. fl. exc.  
p. 17.

*C. stylo bifido*, spiculae subcompositae umbellatij involucentibus triphyllis brevioribus, spiculae  
compressae linearilanceolatae, squamis palmatis angustis, filij late linearibus,  
viciae fibrae.

Icon. Hort. Ic. gr. austr. III. tab. 73

Leaf fl. herb. t. 1. p. 2.

Sibth. fl. gr. Cent. I. t. 48

\* var. *β. viridens*: spiculae basi compressae

Syn. et Icon. *C. viridis* Koch. fl. austr. I. tab. 13 (vicij)  
*C. viridis* Hoffm. R. Germ. I. p. 21. Fiehb. fl. exc. p. 72

Hab. in locis arid., apertis circa Zam, Sebenico, Duvio, Krin; vixit prope Budua. Flor.  
Septembri. Spiculae in specie obfuscae. ☉

Obs. Ductus in hoc albi vixit teres elongatus utrinque acuminatus aculeis triquetris in  
procedente sublenisculis, oblique triquetris, punctulatis. scabris et nigris. Cr. Sm.  
1. p. 32  
fl. gr. prodr. et Jussieu. fl. sic. prodr. 1. p. 41.

+ C. olivaceo Jussieu. in Mem. della Soc. Nat. tom. 13. p. 338 (Sicil. fl. exc. p. 23)

C. stylo triplici <sup>capitata</sup> umbellato, involucris subtrichophyes brevibus, filisque apicem usque  
marginis et carina serrulatis. scabris, spiculis albis diffunditibus lineari-elongatis  
aculeis, bracteolis ovatis oblongis, infimis divaricatis, radice repente tuberosis subovatis.

Syn. C. radiceus Sibth. et Sm. fl. gr. prodr. I. pag. 30 fl. gr. 1. p. 32

Icon. Sibth. fl. gr. Cent. I. tab. 45.

Hab. ad fontem secum vixit circa Ragusa, et Budua. Flor. a Mayo ad Novembrem. Spiculae ferrugineae. subsessilis. ☿

+ C. longus L. sp. pl. p. 67 Herb. fl. austr. 1. p. 56 (Sicil. fl. exc. 1. p. 23. Arab. fl.  
prodr. p. 16. Pet. bot. voyag. n. 521.

C. stylo triplici, umbella secundae composita, involucris 4-6 phyllis longissimis carina et  
marginis serrulatis. scabris, spiculis lineari-elongatis aculeis, bracteolis ovatis imbricatis  
oblongis, radice repente.

Icon. Jussieu. Mem. 2. tab. 297

Hist. Sc. gr. austr. III. tab. 76

Sikensky. Journ. tab. 3. p. 12

~~Math. dis. sup. Dissert.~~

Hab. ad fluviorum ripas, loca palustria, et in partibus uliginosis circa Zam, Tauri, Castelli, Sa  
lona, Ragusa, Cattaro, et in insulis. Flor. Junii, Julii. Spiculae bidae. ☿.

+ C. badius Desf. fl. atl. 1. p. 48. (Sicil. fl. exc. p. 23)

C. stylo triplici, umbella secundae composita, spiculis in radice apice subsessilibus conjungitur lanceolatis  
lineari-bus, involucris tri-trichophyes longissimis carina et marginis scabris, bracteolis ovatis  
imbricatis oblongis, radice repente.

Icon. Desf. 1. c. tab. 3. p. 2

Hab. confusis involucris carina, et Carvata. Spiculae bidae. Flor. Junii Julii ☿

Obs. A procedente, cum quo a quibusdam conjungitur, differt spiculis brevibus, lobatis  
in apice radice congestis, nec late nervos ut in illo. Item Desfontainensis quoad  
spiculum formam minus bona.







*Nlyx. Brulla.*

Hab. in agris circa Zam, Carden, Selma, Aloniza, Merin, et Montem. Her. Mex.

Junio 2.

Ulyx. Culmi hujus glabris strobis contortis videntur.

+ *S. glaucus* Sm. Engl. fl. c. p. 57 (non Lam. ill. 1. p. 142)

+ *S. radice ruga, culmo tereti glabris bafi vagini fligrii tecto, spici aris terminalibus  
corymbis subulatis, bracteis brevibus emittentibus, spiculis emarginatis fimbriatis,  
mijunz brevibus glabris.*

Lyn. Heliconia glaucum Pich. fl. ex. p. 87

Non Née Gen. pl. loc. cit. p. 14-15.

Hab. in agris subulatis lauis rebralis Morinje prope Schin. Mont. Mex., Indis. Spicula  
furfuracea spiculis subulatis. 2.

Obs. Indivisi <sup>bracteis</sup> altera elongata foliacea viridi, altera abbreviata pedicula margini  
albo nervo. Spicula breviter pedunculata, vel omnino nuda ut in Engl. bot.  
t. 2921, Flor. Arn. t. 454, et Hort. Ic. gr. austr. III t. 61.

+ *S. maritima* L. sp. pl. p. 74 Hort. fl. austr. t. p. 59. Pich. fl. ex. p. 79. Dulch. fl.  
Ind. p. 16.

*S. culmi hujus folia supra retrosum cyma, spiculis ovatis - longis pedunculatis ter-  
minatibus multibracteis, <sup>bracteis</sup> spiculis recurvis teretibus, ~~spiculis~~ mjmatis  
ternis pedunculatis.*

var. ~~beta~~. a. <sup>cyma laxa</sup> ~~laevigata~~ <sup>laevigata</sup>: cyma laxa, spiculis pedunculatis sessilibus. Pich. fl. ex.

Non. Hoff. Ic. gr. austr. III. tab. 67 f. dext.

Form. fl. camp. fasc. XII ic.

Engl. bot. t. 542.

var. ~~beta~~. <sup>beta</sup>: cyma densa spiculis omnibus sessilibus

Non. Hoff. Ic. gr. austr. l. c. f. sinistr.

Kr. l. fl. id. tab. ~~15~~ xv.

Hab. in partibus vicijs, plerumque <sup>montanis</sup>, ad fluvium rivi ubique in Dalmacia Mont. Nili.

Angl. Spicula purpurea densa fuscis. 2.

var. ~~gamma~~. <sup>gamma</sup> ~~marginata~~ <sup>marginata</sup>: spiculis cylindricis elongatis

Lyn. Sc. marginatibus W. En. It. Herb. 1. p. 78

Non Scheuchz. Agrost. p. 395. tab. 9. f. 28

Hab. in partibus vicijs, <sup>montanis</sup>, marginibus ubique frequens. Mont. Nili, Angl. Spicula purpurea, fuscis.

2.



Holothroaeum Link

Hort. bot. Berol. 1. p. 293

Ch. Hony hermaphroditis nuda, <sup>superior</sup> spicis multifloris <sup>inferior</sup> subglobosis in capitula sphaericali sublobata  
incomposita congestis brachiis flexis, terminali: culmum subnatis, involucris,  
incomposita congestis, brachiis multifloris imbricatis petalis, segmentis minutis  
staminibus. Stylus filiformis deciduus longus, stylis subglobosis. Anthera  
bryonia mucronata. — Cl. Nicotiana Ord. Monogynia Syst. v. x. —

Holothroaeum vulgare Link loc. cit. Nees gen. pl. fasc. IX.

H. capitulis tuberculato-flores, stylis longis sessilibus, vaginis culmi integris, segmentis filiformibus  
filiformibus. infima aphylla.

Icon Scheuchz. Agrost. tab. 8 f. 2-5  
~~Blatt~~  
Sched. hist. bozd. t. p. 987 fig. 1000.  
Engl. bot. t. 1612  
Nees Gen. pl. fasc. IX. (Holothroaeum)

Syn. Scirpus Holothroaeum L. Sp. pl. p. 22 Hort fl. austr. t. p. 59. Sched. fl. jard. p. 16  
Pelt. bot. wegw. n. 458  
Holothroaeum Linnaei Reichb. fl. exc. p. 26.

— var. *p. romanum*: capitulis ovatis subsessilibus.

Icon Scheuchz Agrost. tab. 8 f. 6.

Syn. Scirpus romanum L. Sp. pl. p. 23.

Hab. in frutibus arboribus, locis aquosis, palustribus sicuti in Sclavonia. Capitula parva: bestia glauca.  
Flort. hinc, Augusto. 4.

Dichostylij Nees et Es.

Cyper. in Linnaea IX. p. 289

Chen. Flores hermaphroditi neis & spicis multiplicibus in capitulum terminale compositum  
bracteis filicibus involucentibus congestis, bracteis multiplicibus imbricatis subaequalibus.  
Stamina tria. Stylus basi vix emisso persistens, bipedus, stigmatibus glabris. Caruncula  
trigona styli basi mucronata. — Cl. Triandria 27. Monogynia Syst. IX. —

Dichostylij Micheliana Nees loc. cit. et in Gen. pl. fasc. IX.

D. culmi trigonis prostratis filicibus, involucre polyphylo, bracteis ovato-lanceolatis adpressis.

Non. Tilli Cat. hort. pison. tab. 20 f. 5  
Hort. Ic. gr. austr. III tab. 69  
Nees Gen. pl. fasc. IX.

Syn. Cyperus Micheliana L. sp. pl. p. 86. Hort. fl. austr. 1. p. 61

Isolopis Micheliana Roem. et Sch. Syst. veg. 2. p. 114  
Dimorphostylij Micheliana Reichb. fl. ex. p. 75 (excl. syn. Linkii)

Hab. in locis aridis, agrorum insularum Uegia et Osseri. Doct. Noi. Capitulum pellucidum  
viride. Hort. Aug. Sept. ☉

Obs. Cyperus Micheliana Delile in Roem. et Sch. syst. veg. Mant. in vol. II p. 67 (1824)  
Link in th. bot. I p. 203 (1822) et Savi in Mem. della Soc. Ital. tom. XXI, est

planta omnino alia, et non Cyperus spicis, ut jam monuit cl. Burdington in Flor. Ital.  
1. p. 304, et cl. Nees in Gen. pl. loc. cit. Planta haec aegyptiaca, quam habui a cl.

Savi ex herbario ill. Nadi, collata cum Dichostylij Micheliana, cujus authenticus speciemina  
etiam liberalitati ejusdem clarissimi professoris, differt ab hac habita refectione  
etiam liberalitati ejusdem clarissimi professoris, differt ab hac habita refectione

elatione capituli & spicis solitariis nec fasciculatis compositis, spicis planis, bracteis  
utrinque delectis  
stylis, infima duplo longior, omnibus lanceolatis subulatis apicibus fimbriatis, quae propter speciem  
pectinatae videtur. Caruncula in utroque ovuli-oblonga trigona apice styli basi persistens  
te et vix emisso mucronata, stylus filiformis bipedus, stigmatibus glabris. Haec planta

aegyptia Dichostylij est, ut bractearum dispositio eam Cyperus, et inter hos Pycnos  
conjugat. Affinis est C. pygmaeus Roem. et Sch. p. 20 tab. XIV f. 5, qui tamen  
differt parvitate omnium partium, spicis in capitulo fasciculatis minimis, bracteis utrinque  
quibus sensim apice subsistentibus, infima spicem vix superante, dum in illo duplo brevior.  
Hinc stipites aegyptia falso nominis, utpote quod falsis synonymis superstructis, hactenus  
iniquitate, ad confusionem tollendam reviserit et aliter nuncupanda, nomine auctoris  
qui primam ejusdem mentionem fecit, de quo flora aegyptia multipliciter est, saluberrima

Cyperus Delilei.  
Culmi D. Micheliana in specimenibus Micheliana longior ac in planta paterna et delila  
haec, ac fere ut in illis iconis cl. Nees supra citatis, in qua tamen caulibus erectis, quo  
nec in aegypto nec in flora Italiae representat.





C. Bauh. prod. p. 12 f. int.

Hab. in pascuis et locis agrorum humidis, hinc Slavonia. Heri in pascuis, capfula castanea  
viresca, unde nomen Schistaceum (α δάμπερα νίτα).

Obs. affinis *S. Schimper* Schk. colon. in 76 qui differt phyllis ovatis, obliq. capsula subquadrata  
et *S. arvensis* Schk. l. c. n. 66, qui diverget phyllis ovatis acuminatis, aristis indivisis  
et longioribus capsula hemisphaerica. In quibusdam siccis originis vixit inveniuntur plantae in planis  
in hinc lineatae reticulatae elongatae, quae hanc plantam videntur a C. Bauh. in prod.  
p. 12 <sup>1000</sup> *arvensis* videri per multos. *Labridum* em, id <sup>1000</sup> *arvensis* vixit inveniuntur  
hinc vult, per specimen d. Schrank in Bavarica fl. 1. p. 615, et d. Schlecht. in  
fl. bot. 1. p. 178.

Hist. Veget. III p.

*S. bulbosus* L. fl. suec. d. II, p. 113. sp. pl. ed. II, p. 466. Hist. Veget. III p. 83

cap. utriusque floris compositis, fl. linearibus canaliculatis, vixit decomposita, perianthia phyllis  
obliq. subrotundis subulatis, capsula subglobosa <sup>1000</sup> *arvensis* perianthia longioribus

Syn. *S. composita* Jacq. en. v. ind. p. 66 et 235 Host fl. austr. 1. p. 458 Reichb. fl. ex. p. 95 (non  
Humb. Bot. v. ger. 1. p. 235, Jacq. Mij. Syn. jun. p. 16)

+ *S. bulbosus* L. fl. suec. d. II p. 113. sp. pl. ed. II p. 466. Hist. Veget. III p. 83. Alth. fl.  
jard. p. 83.

Icon. Host Veget. austr. III. tab. 89  
Leop. fl. herb. t. 13. f. 7  
Hid. t. 431  
Dart. 2. tab. 247

Hab. in pascuis humidis vixit Linn. Scop. Koen. H. v. v. C. Bauh. fl. capfula castanea  
viresca.

*Luzula* DC.

Fl. fr. III p. 170

*L. Pennsylvanica* glumosa. <sup>base</sup> *glaberrima*. *Shimura* n. o. *Capsula* *Stigmata* *in* *glumosa*. *Capsula*  
*trilobis* *umbellatis* *tridisperma*. - *L. Howardii* C. Murray in Syst. ex.

+ *Luzula composita* DC. fl. fr. III p. 161 (<sup>et v. v. v.</sup> ~~excl. v. v. v.~~) Reichb. fl. ex. p. 91. Alth. fl. jard. p. 84

*L. Nuttalliana* DC. foliis <sup>linearibus</sup> *glanis* *marginis* *platis* *venis* *glaberrimis*, *spicis* *ovatis* *umbellatis* *perianthiis*  
*umbellatis* *trilobis*, *perianthiis* *phyllis* *acuminatis*, *capsula* *subglobosa*, *obliq.* *membrata* *longioribus*  
*filamentis* *ardentibus* *humidibus*

Syn. *Nuttalliana* composita a L. sp. pl. p. 466  
*L. Nuttalliana* Host Veget. austr. III. p. 64  
*L. composita* H. austr. 1. p. 455

Icon. Host Veget. austr. III. tab. 92. f. 1-5.  
Leop. fl. herb. t. 13. f. 5.  
Linn. bot. t. 82 f. 1  
C. Bauh. prod. p. 15 f. int. *L. Nuttalliana* DC. fl. fr. III p. 161 et 541

var. p. <sup>congesta: pinnata</sup>  
whiter more diffusa, spicis subumbellatis congestis

Syn. <sup>longipes</sup> Thunb. fl. par. 1. p. 119.

Hab. in grassy ~~soil~~, et in approximatis circa Zam, Vellela, Siga, bar. a monte  
Dikou. Fl. mat. maj. Fl. white. 4.

Obs. Non varietas ad <sup>habitus</sup> *Juncea congesta* Thunb. fl. par. 1. p. 119. (Syn. a *glaberrima*)

varietas <sup>habitus</sup> *Juncea multiflora* Lij. <sup>et pinnata ac</sup> *pectus* <sup>et in</sup> *vestib*, *pectus* *debat* *debat*, *non* *spicae* *in*  
*spica* *viride* *pedunculata* *int.* *et* *in* *scabris* *Deak.* *theat.* *bot.* *p.* *104*, *Moj.* *herb.*  
*skp.* *ex.* *sub.* *8* *t.* *g.* *f.* *1.* *rep.* *restituta.* *Neluy* *quod* *non* *Flora* *Senica* *1892*,  
*id* *istum* *in* *hae* *hijis* *major*, *et* *spicae* *numerosae* *ac* *in* *flora* *Senica*.

+ *L. albida* *St. f.* *cat.* *p.* *159* *Thunb.* *fl.* *ex.* *p.* *92*

*L. folijs* *linearibus* *marginibus* *pilosis*, *apice* *pyramidalibus* *effusa*, *capitulis* *subpinnatis*,  
*floribus* *phyllis* *exterioribus* *numerosis* *capitulis* *trijugis* *elipticis* *involucris* *longioribus*,  
*antheris* *subumbellatis*.

Syn. *Nanus* *albidus* *Hoffm.* *fl.* *germ.* *ed.* *1.* *p.* *126* *Hort* *Sc.* *yo.* *num.* *III.* *p.* *60* *et* *St.* *aug.*

<sup>p. 456</sup>  
*L. nitens* *Leop.* *fl.* *herb.* *p.* *91* (nec *L.*)

*Non* *Hort* *Sc.* *yo.* *num.* *III.* *p.* *95*

*Leop.* *l.* *c.* *tbl.* *13.* *p.* *9.*

*C. Barb.* *herb.* *p.* *16* *f.* *ext.*

Hab. in grassy <sup>montium</sup> *Vellela* *et* *Dikou*, <sup>et</sup> *St. Jul. Aug.* *Fl.* *indiv.* *alt.* *4.*  
*et* *in* *scabris* *apice* *subumbellatis*. *Fl.* *St. Jul. Aug.*



*Hed.* species in agris agris agris *Sobociensis* loci *Pasinae* hils. *varietas* *ijda* loci  
 circa *Zam.* <sup>fragrans</sup> *Hed.* flowers in specie <sup>substantia</sup> *complanata* *varietas* *caeruleo-virescenti*.

*Obs.* Species *polytricha* hirsuta *pappi* *myrtidina* *lytham* et *br-gendij* *hilsian*.  
*filij* *hirsuti* ~~longioribus~~ <sup>et brevibus</sup>, et *longioribus* *agrostoidibus*, *filij* *minij* *glauces*, *super* <sup>scapo</sup> ~~scapo~~  
*hirsuti*, *si* *scapo* *caulis* *interdum* *brachyformis* *inter* *plantis* *partem* *caespiti*  
*adhaerent*; *perianthia* *toti* *visu* *filij* *longiora*, et *filij* <sup>forma variabilis</sup> *exsertenti*, *visu* *in*  
*utroque* *brachy* et *longioribus*, *limbo* *chilo* *major* *aut* *minor*. *Hinc* *visus*  
*a* *specie* *minima* *differt*, *cum* *inter* *utroque* *forma* *omnis* *intermedias* *occurrit*.

*Trij* *Chamaecrista* *el.* *Bastlois* <sup>in</sup> *agris* *indica*, <sup>est</sup> *ut* *ex* *artypis* *habeo* *indij*  
 et *ex* *amissione* *d.* *saifil.* <sup>et</sup> *et* *d.* *liberenti* *Luna* *longi* *differt*, *visu* *in* *india*  
*iblican.* <sup>et</sup> *d.* *liberenti* *La* *Reverend.* *Bot.* *p.* *285* *ab* *d.* *specie* *habeo*

*visu* *in* *india* *prospicere*, *et* *habeo*, *caespitibus* *longioribus*, *visu* *in* *habeo*  
*differt* *in* *india* *visu* *in* *india* *idque* *minimam* *speciem* *probat*.

*in* *complanata* *visu* *in* *d.* *saifil.* *Reverend.* *Bot.* *p.* *285*.

1. *Gladia communis*

2. *G. segetum*

3. *G. trifloris* Sibth. mss. ex fl. gr. jard. i. p. 25 *Benth. fl. Ital. i. p. 225*

*Gladia communis* L. sp. p. 2. p. 52 Herb. fl. austr. i. p. 44  
ex loc. (excl. sign. Sibth.)

*G. spica multiflora*, floribus <sup>residui</sup>, antheris filamentis subaequalibus, filisaeae <sup>membranis</sup> ~~sub~~ <sup>subparallelis</sup> ~~sub~~ <sup>subparallelis</sup> nervosis

*Icon Cur. bot. mag.* tab. 86

*Lam. Enc. bot. pl.* 32. f. 1

*Bull. herb. de la fr.* tab. 8

*Reich. europ. myc.* tab. 110 (inc. 183)

*Park. jard.* p. 159. t. 191 f. 1

*Hab.* inter lapides in silva et circa Lagosa. Hort. Major. Pinarum rubro-violaceo.

*Obs.* Specimen Livigiana synonymum C. Bauh. (*G. floribus* cum infra dicitur *Pin. 41*) jam in  
pina dicitur *Spec. Plant.* a Livigno ipse mentionem, et in ditione altera ejusdem operis item  
pina supra citata dicitur designatum, quod synonymum et icon et palmi ~~in~~  
floribus supra operis eadem, hinc omnino spectum arboris, nec *G. segetum* fact.  
Vix floribus dicitur jam fronte ad illa recedit, et a Livigno ipse in *est. duodecim*  
*syn. not. p. 176*  
*syn. p. 176* pro illis var. *G. trifloris* quae cum synonymum C. Bauh. (*G. utriusque floribus*  
*pin. 41*).

*G. segetum* Savt. in bot. mag. p. 419 *Reich. fl. eur.* p. 83

*G. spica multiflora*, floribus <sup>residui</sup> distinctis, antheris filamentis longioribus, filisaeae <sup>membranis</sup> ~~sub~~ <sup>subparallelis</sup> ~~sub~~ <sup>subparallelis</sup> nervosis

*syn. G. communis* L. *ex loc*  
*syn. G. communis* L. *ex loc*

*G. Livigiana* Van Hook. *herb. fl. it. mag.* p. 1. *Reich. bot. mag.* v. 182

*G. byzantina* Wy. *it. mag.* p. 36 *ex Mell.*

*Icon Sibth. et Sm. loc.* tab. 37

*Icon. Peign.* *Savt. fl. v. cent.* vi. f. 819

*Savt. bot. mag.* t. 519.

*Sibth. et Sm. fl. gr. i. tab.* 37

*Robt. in. mag.* p. 99. f. *ixixiv.*

*Ulyss. Mac. Macinas. Sablan.*  
*Hab.* inter lapides in silva dicitur prope. Hort. Ag. Major. Pinarum rubro-violaceo.

Obs. *Sifon* a fl. *typharia* Mill. Diet. n. 3. Seibb. ic. fl. nr. cent. VII. p. 22. ic. 866  
floris melle viscosis, laciniis univ. pteris, ped. lanceolatis univ. vix.  
Sedley Montigny d. Pav. in 1414. p. 208. ~~Seibb. ic. fl. nr. cent. VII. p. 22. ic. 866~~  
Nt. hinc affinis, sed univ. *Sifon* var. ~~... ..~~

*S. hispidus* Sibth. in fl. gr. pad. I. p. 25. ~~Seibb. ic. fl. nr. cent. VII. p. 22. ic. 866~~

*S. spica subtota*, floris secundis, anthod. flavens biniat. filij linearis univ. formis.  
but. tenuis. ~~... ..~~

syn. fl. communis Seibb. ic. fl. nr. cent. VI. p. 26. et fl. nr. p. 83  
fl. communis ~~... ..~~ Sibth. et Sm. fl. gr. pad. I. p. 25. et fl. gr. I. p. 28.

non. Seibb. ic. fl. nr. VI. p. 811  
Sibth. et Sm. fl. gr. I. p. 38. 38

Lanich. ist. de p. imp. v. p. 53, ad caput inter Pappoch et Montigny,  
in p. 53. ~~... ..~~ *Sifon* ~~... ..~~  
Hab. inter septa ~~... ..~~ et in floris ~~... ..~~ *Sifon* ~~... ..~~  
~~... ..~~ *Sifon* ~~... ..~~ Hort. Moja, ~~... ..~~ *Sifon* ~~... ..~~

Obs. *Sifon* minor, sub *hispidus*, argenteis *Sifon* ~~... ..~~  
... *Sifon* ~~... ..~~ C. Bauh. et *Sifon* ~~... ..~~  
~~... ..~~ *Sifon* ~~... ..~~ Seibb. in *Sifon* ~~... ..~~  
non. syn. fl. gr. p. 699, *Sifon* ~~... ..~~ *Sifon* ~~... ..~~  
... *Sifon* ~~... ..~~ *Sifon* ~~... ..~~

*S. illyricus* Koch syn. fl. gr. p. 699

*S. spica subtota*, floris secundis, anthod. flavens biniat. filij linearis univ. formis,  
tenuis. ~~... ..~~

Hab. inter septa, et in montanis circa ~~... ..~~ et in floris ~~... ..~~ *Sifon* ~~... ..~~  
... *Sifon* ~~... ..~~ Hort. Moja, ~~... ..~~ *Sifon* ~~... ..~~

Obs. ~~... ..~~ *Sifon* ~~... ..~~ *Sifon* ~~... ..~~  
... *Sifon* ~~... ..~~ *Sifon* ~~... ..~~  
... *Sifon* ~~... ..~~ *Sifon* ~~... ..~~  
... *Sifon* ~~... ..~~ *Sifon* ~~... ..~~

E

#

- *J. communis* d. p. pl. 3. p. 52 Hist. fl. austr. 1. p. 64 (exl. syn. et ic. libla)

Nicht. fl. ex. p. 83

*J. communis*  
*J. floribus secundis*

antheris filamentis subaequalibus, filis aequalibus.

Dion. lurs. bot. maj. 16. 86.

Nicht. flo. pl. nr. cens. 11 p. 818

dam. enc. bot. plant. 32. p. 1.

Flor. mong. imp. 16. 110 (nec 103)

Dob. h. imp. p. 98 p. Text.

Hb. inter sax. insulae Leina, et una Wagosa. Flor. maj. Perianthii subaequali. 2.

66. *Specieo lionensem* synonymo C. Daul. (*J. floribus utroque oppositis major et gravior.*

Pan. 41) jam a prima odore spec. Plant. a Liugo esse nuntiat, et icon

quasi supra citata (ob eadem eandem <sup>quidam</sup> <sup>interdum</sup> ad huc altera spec. pl. p. 82.

(<sup>hic</sup> ~~hic~~ <sup>est</sup> ~~est~~ <sup>typographia</sup> icon hujus nomen post 103) <sup>est</sup> ~~est~~ <sup>quod</sup> ~~quod~~ <sup>synonymum</sup> ~~est~~ <sup>et</sup> ~~et~~ <sup>icon</sup>

in nostra opinione <sup>est</sup> ~~est~~ <sup>hanc</sup> ~~hanc <sup>omnem</sup> ~~omnem <sup>sp. 103</sup> ~~sp. 103 <sup>est</sup> ~~est~~ <sup>quod</sup> ~~quod~~ <sup>synonymum</sup> ~~est~~ <sup>et</sup> ~~et~~ <sup>icon</sup>~~~~~~

in <sup>est</sup> ~~est~~ <sup>hanc</sup> ~~hanc <sup>omnem</sup> ~~omnem <sup>sp. 103</sup> ~~sp. 103 <sup>est</sup> ~~est~~ <sup>quod</sup> ~~quod~~ <sup>synonymum</sup> ~~est~~ <sup>et</sup> ~~et~~ <sup>icon</sup>~~~~~~

ab <sup>est</sup> ~~est~~ <sup>hanc</sup> ~~hanc <sup>omnem</sup> ~~omnem <sup>sp. 103</sup> ~~sp. 103 <sup>est</sup> ~~est~~ <sup>quod</sup> ~~quod~~ <sup>synonymum</sup> ~~est~~ <sup>et</sup> ~~et~~ <sup>icon</sup>~~~~~~

p. 15 <sup>est</sup> ~~est~~ <sup>hanc</sup> ~~hanc~~ <sup>omnem</sup> ~~omnem <sup>sp. 103</sup> ~~sp. 103 <sup>est</sup> ~~est~~ <sup>quod</sup> ~~quod~~ <sup>synonymum</sup> ~~est~~ <sup>et</sup> ~~et~~ <sup>icon</sup>~~~~

- *J. triflorus* <sup>est</sup> ~~est~~ <sup>hanc</sup> ~~hanc~~ <sup>omnem</sup> ~~omnem~~ <sup>sp. 103</sup> ~~sp. 103 <sup>est</sup> ~~est~~ <sup>quod</sup> ~~quod~~ <sup>synonymum</sup> ~~est~~ <sup>et</sup> ~~et~~ <sup>icon</sup>~~

*J. triflorus*  
*J. floribus secundis*

antheris filamentis brevibus, filis aequalibus.

Agri et Dion. *J. communis* var. Blas. et Sm. l. c. p. 28 tab. 38

Danisch. ist. plant. ser. tab. 39.

Hb. inter Leina et Wagosa non Wagosa, et in placidia Dion. et

Leina in Wagosa Wagosa. Flor. maj. Perianthii

subaequali. 2.

66. *Circa minor*, submissifolia, <sup>est</sup> ~~est~~ <sup>hanc</sup> ~~hanc~~ <sup>omnem</sup> ~~omnem~~ <sup>sp. 103</sup> ~~sp. 103 <sup>est</sup> ~~est~~ <sup>quod</sup> ~~quod~~ <sup>synonymum</sup> ~~est~~ <sup>et</sup> ~~et~~ <sup>icon</sup>~~

C. Daul. reparatorem a Liugo pro J. communis citatum. Dion. et

Tunicy submissifolia, antheris filamentis brevibus, filis aequalibus. Dion. et

pl. 3. p. 225-227 inducit in placidia Dion. et Wagosa. Perianthii

subaequali. 2.



f

Clavis analytica generum

Herbarij caulibus vel. N. 1. Herbarij radicebus vel. N. 2.

- |   |   |
|---|---|
| 1. Perianthii tubo vix, limbo exteriori reflexo | Perianthii tubo immixto, limbo exteriori erecto |
| <u>Iris</u>                                     | <u>Stadibus</u>                                 |
| 2. Perianthii tubo brevissimo                   | Perianthii tubo longissimo                      |
| <u>Ranunculus</u>                               | <u>Crocus</u>                                   |

10  
Icon. Reichb. pt. crit. I p. 193.

Column. expt. E. p. 209 fig. sup.

Hab. inter segetes Dalm. Forette, Siberico, Spalato, Salona, Ujfa, Regusa.  
Flores apertis, saepe. corolla carnea. &

Obs. caulis scaber, foliis radicalibus obovato-spathulatis integris, caulinis lanceolatis obtusis basi incisae pinnatifidisque, bracteis adpressis ovato-lanceolatis ciliatis margine membranaceis. Fructus hirsutus pubescente angulatus, antice exaratus planiusculus, corona intus callosaque glabra nervosa marginata, et reticulata, dentibus ovatis lanceolatisque apice subulatis, quandoque alatis patentibus, semper uncinatis. Corona intus glabra differt ab affinis *F. coronata* D. C. quem confer in Prodr. syst. nat. IV p. 628. Non *Reichenbachii* bona quoad fructum, sed folia in nostra basi incisae. Non *Columna* huic pertinet, consentiente etiam A. Gussone in fl. siut. prodr. I. p. 35, cuius *F. coronata* huic referenda.

63. *F. discoides* Loisel. notie. Hort. fl. austr. I. p. 43. et Reichenb. fl. ex. 2. p. 199. quoad locum natalem (excl. descript. et icon.)

*F. fructu* oblongo subtriangulo antice exarato corona rotata, dentibus subsens lanceolatis obsolete nervosis inciso-dentatis apice breviter subulatis uncinatis, caule dichotomo divaricato diffuso.

Syn. *Fedia discoides* Vis. Strep. Dalm. p. 35. (an *Vahl.*?)

Deser. caulis ad angulos scaber dichotomus diffusus divaricatus, foliis obtusis glabris, radicalibus oblongo-obovatis incisae, caulinis oblongo-linearibus basi linearibus pinnatifidis. Florum capitulo terminali oppositi hemisphaerici, vel sub-globosi. Bractea lanceolata obtuse margine membranaceae ciliatae fructui adpressae. Fructus oblongus

subtetragonus villosus, corona rotata centro villosa, dentibus quinque  
ad septem lanceolatis ciliatis, plerisque inciso-dentatis, apice  
subulatis uncinatis, glabris, tenuissime nervosis.

Obs. Differre videtur a *V. diffoidea* Loeb. caulibus divaricatis, et  
coronae dentibus nunquam duodenis integris ut in Reichb. Pl. vit. II.  
p. 226. vel subsenis integris ut in Morison. hist. 3. sed. 7. t. 16. fig.  
sed subsenis inciso-serratis. An reapse distincta, et *V. divaricata*  
appellanda?

Hab. in apricis herbidis circa Sebenico, Travi, Spalato. Floret aprili  
Majo. corollae roseae. ☉

XXI. Crocus Linn.

Ch. Stylis radicalibus. Perianthium <sup>sen. pl. p. 17. vich</sup> tubo longissimo, limbo <sup>inequaliter</sup> scapartito  
laciniis <sup>erectis</sup> alternis majoribus, spatula hypogyna. Stamina tubo inserta.  
Stigmata tria dilatata involuta <sup>inuis</sup> dentata. Capsula trilocularis trival-  
vis polysperma, seminibus subrotundis. Ord. nat. Liliac. Herb.  
Et. Siquan. Did. Monog. v. 4. p. 44.

Crocus vernus Allim. fl. pedum. 1. p. 84. Hook. fl. austr. 1. p. 42. Reichb.  
fl. cauc. p. 28. Deff. bot. veyr. n. 209.

f. tunicis radicalibus fibroso-intertextis, spatula florali monophyl-  
la, tubi fauce pilosa, Stigmatibus abbreviatis incisus tubulosis plu-  
catis Stamina superantibus.

Syn. Crocus sativus  $\beta$ . vernus Linn. sp. pl. p. 50.

Icon. Tenor. Mem. sur Crocus napol. 1826. Tab. I  
Engl. bot. t. 344.

Mull. herb. de. la. p. 1. 351.

Nag. fl. austr. v. 8. 36.

Reichb. Fl. bot. tene. de. pl. var. cent. t. 7. 1250

Hab. in montibus Mojzech, Cromina, Svitloja, Piskavo, Starke, Mospor, et  
Sironiakh

<sup>bulbos major</sup>  
Δ In montibus subrotundis floris lacrima longior pera explanata, flus minor: in montibus laetis  
aspingit e contra. 2.

lois asperis collinis in Dalmatia montana. Floret a Febuario, ad Majum.  
Flores albi, vel basi violaceo-erulei, vel margine laciniarum albi  
dorso purpurei, vel utriusque utriusque violacei, vix odorati. Varietates  
cont. in G. Reich. v. p. m. Cent. X. p. 1250-54. Δ  
C. reticulatus Reich. a. v. p. m. Cent. X. p. 1250-54. Hausskn. fl. v. p. m. Cent. X. p. 1250-54.  
Kov. ex Dom. in v. p. m. Cent. X. p. 1250-54. Hausskn. fl. v. p. m. Cent. X. p. 1250-54.  
austr. 1. p. 123. Reich. fl. v. p. m. Cent. X. p. 1250-54. Hausskn. fl. v. p. m. Cent. X. p. 1250-54.  
C. tenuis radicalibus vix fibris reticulatis, spatula florali  
diphylla, basi fauce glabra, stigmatibus <sup>to p. m. Cent. X. p. 1250-54.</sup> elongatis missis stami-  
na subsequentibus.

Syn. C. variegatus Hoopp. et Hornsch. Preis ad. Mer.

Mon. Nieberst. Cent. plant. v. p. m. Cent. X. p. 1250-54.

Hoopp. et Hornsch. loc. cit. in  
Reich. v. p. m. Cent. X. p. 1262-65  
Hab. in praevis et locis apricis asperis Pontanniae, et Batinibus pro  
pe Kara, et in insula Rago. Perianthium alido-violaceum, lacini-  
nis externis lineis tribus ceruleis pinnato-striatis, interioribus  
unicoloribus, omnibus basi lutescentibus ibique ceruleo-striatis,  
striis in tubum decurrentibus. Floret Martio, aprili. 2.

66. C. pusillus Tenor. fl. v. p. m. Cent. X. p. 1250-54. Hausskn. fl. v. p. m. Cent. X. p. 1250-54.  
C. tenuis radicalibus membranaceis nitidissimis supra basin circum-  
suis spatula florali diphylla, basi fauce glabra, stigmatibus <sup>proprie trifida</sup> elongatis indivisis stamina superantibus.

Syn. C. triflorus Ait. v. p. m. Cent. X. p. 1250-54. Hausskn. fl. v. p. m. Cent. X. p. 1250-54.

Syn. C. pusillus Tenor. fl. v. p. m. Cent. X. p. 1250-54. Hausskn. fl. v. p. m. Cent. X. p. 1250-54.  
C. biflorus var. c. Reich. fl. v. p. m. Cent. X. p. 1250-54.

Mon. Tenor. Mem. sui (roth. lab. II. p. 1250-54.)  
Reich. v. p. m. Cent. X. p. 1250-54.

Hab. in praevis montanis apricis in Lemese inter Duris et  
Vertika, in Mariano prope Spalato, in Padigone et Belyak, et  
in montibus Ragusinis. Floret mens. Martio. Flores omnino

albi, vel basi cum tubo livida coerulesc. 24.

Obs. Aequa nos semper uniflorus; hinc et Ferrucii nomen licet  
posteriori adhiberi. Folia nunc flore breviora et tardiora  
sive hysteranthia, nunc flore longiora et coactanea, sive  
synanthia. Haec sola species a nostratibus Stigmatibus gaudet  
constanter integris, dum in reliquis Stigmatibus ludunt  
integra, crenulata, missa vel etiam partita, hinc <sup>11. 15.</sup> nulli <sup>pro</sup>  
~~haec~~ <sup>specimen</sup> <sup>minus tate</sup> discrimina pro speciebus distinguendis suaperidant.  
Non citata nostris optime congruent, huc potius quam Cili-  
neate. Haec pertinet, juxta specimen a cl. Quetere commu-  
nicata.

5. 67. C. Pallasii Marsch. a Reichb. fl. haur. canu. suppl. p. 35.  
Goldb. dispers. de proo n. 13. Alsch. fl. jadr. p. 14.

C. hemicis radicalibus fibroso-interbaccatis, spathe florali  
diphylla, tubi fauce fasciculatim villosa, Stigmatibus eton-  
gatis clavatis erosis incisive Stamina subaequantibus.

Syn. C. odoratus Vis. Stirp. dalm. p. 32. Pett. bot. vngar. n. 307. (non  
Pisoni-Bernardi)

C. longiflorus Reichb. fl. ear. & in add. p. 140. 13. n. 589. quae  
patriam dalmat. et cael. descript. (non Paspinisque)

Desov. Bulbus tubi orbiculatus depressus hemicis fibroso-inter-  
baccatis, nec lamen caute reticulatis oblectus. Spathe binae  
a bulbi apice ortae equistantes obtuse, scariose, folia flores-  
que includentes. Folia coactanea, sed post anthesin longiora,  
longa, flore breviora vel subaequantia, glabra vel margine hemicis  
sine ciliata, supra alba linea notata et via canaliculata

Non. Dem. loc. cit. hb. iv.  
Reichb. de. fl. n. cent. X. p. 1281.

margine

margine utroque subtus unijulcata, medio plana. Spatha  
florales binae lanceolato-linearis equitantes acuta a basi ger-  
minis orta, folijs equals vel isdem breviores. Flos unicus, vel  
geminus (raro tres), pediculo scapiformi suffultus, tubo lon-  
gissimo tracti gracili folia superante, albido vel livescens,  
saucio dilatata violacea, intus tuba staminifera, ibique  
fasciculatio villosula, limbo campanulato, lacinijs ovalibus  
obtusis violaceis venosis. Stamina tria filamentis angula-  
tis ochroleucis antheris linearis-sagittatis luteis aequantibus.

Pistillum gemine pedicellata, stylo longiforme, stigmatibus croceis ele-  
vatis, quandoque infundibuliformibus, truncatis, obliquis, erosis missi-  
ve, stamina superans vel isdem brevis. Capsula ovali-oblonga sexstrata.

Hab. in collibus saxosis apricis, et in dumetis, circa Kara, Dennis, Kagoric,  
Skilivri, Sabinio, Poraja, Trau, Salone, Spalato, et in insulis Buca,  
Solha, Brarka, Seluda. Floret octobris Novembri. Flores violacei

Illuv. coerulesci raro albi. <sup>21</sup>  
Cab. Ejovan. Bendiska. Supran. <sup>3</sup> quem esse in d. flor. leg. var. x. p. 122-23.  
Differt a f. odoro. Paivan, cum quo <sup>olim</sup> ~~conspicitur~~ committitur, stig-  
matibus nunquam multifidis, saucio villosa, et praecipue spatha diphyl-

Udud. Stigmata hujusce in Dalmatia colliguntur et venduntur pro illis  
C. sativi isdemque usibus tum medicis tum oeconomicis inserviant. pro-  
cum suspiciant subtiliorem sed magis fragrantem, qui saucio cul-  
tura crispior fit, et orientali simillimus.

n. 6  
Romulo Masat.  
Pl. rarit. et sativa. p. 13.  
**XXXIII. Trichanema Boerl.**

Ch. Flores radiales. Perianthium tubo recto brevissimo limbo carn-  
praecente et d. Boerl. <sup>in Exp. Scientif. de Morca l. c. p. 21</sup>  
pan. qui habitus et utriusque generis cum uno confuit. Propolis filamentum  
muss - antheris in quatuor dividit. Peria de pichb. instrum. videlicet exhibet.  
Stigmata alij <sup>provisus</sup> ~~trichanema~~ generum longioribus.

panolato. patente equaliter seapartito. Spatha hypogyna diphylla. Stig-  
mata tria angustissima bipartita. Capsula subglobosa trilocularis tri-  
valvis polysperma, seminibus subrotundis angulatis. <sup>ad nat. Sides. Vulp.</sup>  
<sup>U. Mundi Ord. Ichneumon</sup>  
~~1688~~ Bulboodium ~~Bar. Reichb. fl. cae. t. p. 83. & best. et Maur. fl. rom. p. 17.~~

Sp. foliis linearibus sulcatis basi canaliculatis flexuoso-recurvis scapo  
subunifloro longioribus, spatheis floribus brevioribus, Stigmatibus stami-  
na superantibus.

Syn. Naia Bulboodium. Linn. sp. pl. qu. tit. Nid. stirp. danm. p. 38. Hook. fl.  
austr. t. 1. p. 44. Alsch. fl. jard. p. 14. Willd. bot. siccogr. n. 521.

Icon. <sup>Trichonema Bulboodium Kex in Guk. of bot. t. 1. p. 22. Reichb. fl. cae. p. 83</sup>  
Sibth. fl. graec. cont. I. t. 26.  
Hay. icon. rar. v. 1. p. 241.

Engl. bot. t. 2549.  
Hort. gen. pl. p. 5. (Trichonema)  
Hab. in pratis herbidis apricis tum insularum tum continentis totius  
Dalmatiae. Flores jam a Januario in insulis (Zurula, Lisina, Medea  
etc.; Martio ubique. Perianthium fundo flavescente, intus albo viola-  
ceum, catus virens. &

Obs. Scapus quandoque biflorus vel triflorus et hunc profunde  
divisus non ramos seu pedunculos elongatos, quamquam prima  
frons flos inferior sessilibus videatur, eo quod pedunculi sunt infimi  
cum spatheis conjuncti. Spatha sub gemine inserta diphylla, valvis  
lanceolatis. Folia basi tantum canaliculata marginibus conniven-  
tibus, apicem versus ob margines simul coalitos compressa.

**XXIV. Gladiolus Linn.**

Ch. Flores spicati. Perianthium <sup>Sp. pl. p. 18.</sup> infundibuliforme, tubo incurvo, limbo  
inequaliter seapartito. Spatha floralis <sup>diphylla</sup> bivalvis. Stamina adpendentia.

*Stigmata* tria oblonga indivisa. capsula ovalis trilocularis trivalvis,  
#2 seminibus alatis. ~~ad. n. Ludov. herb.~~ - (L. Pinandri or. Vnnoyini  
~~radiculis communis L. sp. pl. p. 52. Herb. fl. austr. t. p. 114. ex. Deser.~~  
et in Sibth. Reichb. fl. ex. t. p. 83. Uloch. fl. jard. p. 16. ?  
fl. floribus secundis, antheris filamentis brevioribus.

Icon. Jurt. magat. tab. 86.

Reichenb. Pl. vici. Cent. VI. f. 81f.

Sp. in. mon. 163.

var.  $\beta$ . *triphyllus*: caule abbreviato triphylo, spica triquadriflora.

Syn. et L. *triphyllus* Sibth. fl. gr. Cent. I. tab. 38.

Hab. inter segetes prope Pragusa, et in prope Dvorsno Dicta in monte  
Krebojice prope Cattara cum varietate. Floret Majis. Perianthia  
rubra - violacea, in varietate  $\beta$ . (adnata inferior) albida.  $\gamma$ .

Obs. Distinguitur a *L. intricato* L. bulbo tubero simpliciter non duplici,  
perianthiis laciniis obtusis, seminibus ipsi longius adnatis et ideo  
perigynis, floribus majoribus rariis paucioribus. Cfr. et  
Reichenbach loc. cit.

10. *L. segetum* Jurt. bot. mag. t. 719.

*L. floribus* Distichis, antheris filamentis longioribus.

Syn. *L. byzantinum* K. Stirp. Dalm. p. 36. non Miller.

*L. Ludovici* Van herb. fl. det. sup. Pitt. bot. mag. n. 452.

*L. communis* Sibth. fl. gr. non Linn.

*L. dalmaticum* Paucsch ap. Reichb. fl. ex. t. in add. p. 140. 12. n. 176.

Icon Reichb. Pl. vici. Cent. VI. f. 81g.

Jurt. bot. mag. t. 719.

Sibth. fl. gr. Cent. I. t. 3f.

Illyr. Mac, Mozinax, Sabljan.

Hab. inter segetes totius Dalmatiae a terra ad Ragusa, nec non in insulis.

Floret apertis. Mojos. Perianthia rubro-violacea. 2.

Obst. Dignoscitur facillime a precedentibus characteribus Italij, a Th. byz.  
Kantno Mill. Dist. n. 3. Reichb. Pl. vici. VII p. 22. ix. 866. floribus  
multo minoribus, laciniis non ovatis patentibus, sed connatis  
tibus lanicatis. N. dalmaticus Tausch a profluo solummodo de  
versus floribus subspicatis erectiusculis, huc pertinere videtur,  
cum propter hanc nulla alia Radialis species in Dalmatia occur-  
rat antheris filamentis longioribus.

### XXV. Iris Lun

Ch. Perianthium infundibuliforme tubo recto, limbo inequaliter sex-  
partito, laciniis externis reflexis. Spatha bivalvis. Peridium tri-  
fidum, laciniis petaliformibus sub petala longitudinali stigma-  
tibus, Stamina incurva sequentibus. Capsula oblonga, trivalvis,  
trilocularis, polysperma seminibusque unguatis. Ord. nat. Lincei  
Suss.  
\* perianthii laciniis reflexis supra barbatis.

N. Iris punctata L. 3p. pat. p. 56. Vis. stirp. Dalm. p. 38. Hort. fl. austr.  
1. p. 26. Reichb. fl. cauc. 1. p. 80. Alsch. fl. Jade. p. 15. Pitt. bot.  
svegr. n. 511.

I. scapo brevissimo uniflora, foliis ensiformi-baccatis brevioribus.

Syn. I. (lusiana) Tausch in Reichb. fl. cauc. 1. p. 80. et in ad. 2. p. 140. 12  
n. 556.

Icon. Jurb. bot. magar. 1. 9.

~~Iris violacea parva~~ Labat. <sup>fig.</sup> icon. p. 63. f. 1. p. 64. f. 2.

Harz. fl. austr. 1. 1.

var. *p. luteus* flore luteo.

Hab. specij in campis apricis agris Liburniensis loco Pradine dicto, varias in locis similibus prope Kara. color florum in specie saturate coerulescens. Flores apicis. 2.

Obs. florissimus ~~Kausch~~ ~~Reichenbach~~ aliam ex hac plantat <sup>constituit</sup> speciem *T. Clusiana*, nomine, *fluvii et dabitur* firmatam synonymis. et *T. Dalmatica minor*. (Lus. hist. pl. p. 221 ~~et~~) floribus ~~pure~~ <sup>pure</sup> dilutioris colore predictis tantummodo nota, sine ulla descriptione vel icona, est planta <sup>opinio obsequi</sup> ~~hinc dicitur~~, et nunquam recte exhibitanda, et nulli profecto et speciebus natis veluti synonyma ~~hinc dicitur~~ exhiberi poterit, ~~certe necesse est~~ ~~et~~ ~~novam~~ ~~speciem~~ ~~aliam~~ ~~hinc~~. *T. Dalmatica minor* *Label.* icon. p. 62. est plantat quadriflora et ideo *T. Clusiana* biflora omnino dissimilis, et icon hinc est ipsissima, quam *Fludius* ipse proposuit sub diverso nomine *T. latifolia majoris* XVIII. in hist. pl. p. 222; quo probatur iconem hanc diverse etiam plantat pertinere ab illa, quam idem auctor antea dixerat *T. Dalmaticam* minorem. Harum utraque ab *Tride* nostra longe alienissima. *Tris. pumila* *Curtis* bot. mag. t. 9 est plantat uniflora, nostram optime representans, et ideo nequit ad *T. Clusianam* bifloram adduci. Nihil tamen minus et *Reichenbach* in ff. rar. ad. t. p. 140. 12. n. 556. affirmavit *Tridem* nostram a Kara suam *Clusianam* esse, quamquam icones supra dictae specij diversissimas exhibeant, et quamquam *Thiers* nostra constanter sit uniflora, et *T. pumila* *L.* et *Targuini* ad unquam similes. Nec ullum horum deserviment inter hanc et novam illam speciem intercedere constat, collatis utriusque diagnosticis, nam ex observationibus in vivo factis et repetitis colligitur, spolia

etiam in *T. pumila* locis asperis et calidioribus ortis glaucosens;  
tubum in individuis pusillis maxime spatij semper margine  
scariosis, et nec in nostra omnino herbae, inclusum et crispiusculum  
esse; in alijs loco fraxiore ortis elongati, attenuati, et spatiosior  
des, uti in herbario nostro adsunt exempla. Folia nunc flore bre-  
viora, nunc eundem aequantia. Colorem denique florum in hac  
planta plurimum variare videtur, et sibiimum, levisimamque hujus  
differentias in *T. Alessiandra* indicatas, de nullius ea Philosophia  
botanica, multo minus in planta verficolori speciem constituere  
certissimum est. Nunc *T. flusianam*, Saltim Dalmaticam, ab *T. pu-  
mila* nullimode diversam esse contendimus.

*T. germanica* L. sp. pl. p. 55. Sib. stirp. Dalm. p. 38. Koel. fl. austr.  
p. 55. Reichb. fl. ex. 1. p. 81. Uelch fl. jard. p. 15. Cett. botan.  
mag. n. 507.

*T. caele* <sup>ramo</sup> multiflora folijs coriiformibus falcatis longiore, floribus  
inferioribus pedunculatis, <sup>perianthii</sup> calathii tubo germinem superante.

Syn. *T. illyrica* Theophr. Hist. pl. l. ix. cap. 11. Diosc. Nicand. Galen. Pin.  
Anquill. parer. 1. p. 17-18.

Non. Sibth. fl. gr. I. 40.

Blackem. herb. 1. 1. 69.

Reichb. De. fl. m. cont. X. f. 1245.

Matth. disc. supra Diosc. 1. p. 19. Nees gen. pl. fr. v. Trj. p. 17.

Ulyr. *Maingar*, Xilj *modre*.

Herb. in muris vetustis, locis sylvaticis, asperis, et ruderalibus locis Dalmo-  
tia vulgaris. Floret aprili. Majo. Flores magni lacte cerulei,  
laciniis reflexis obscure violaceis basi albo-lineatis, barba al-  
bida. 4.

Ufus. Radix hujus plantae jam a primis botanicis et medicis inu-  
natis celeberrima fuit, eaque caeteris longe praestantior conseba-  
tur, quae St. illyricae nomine in Dalmaticae regione illa, quae flumina  
Drilonem et Naronem interjacet, colligebatur, ubi ex Nicandro et  
Plinio patet, unde magna foenore in usum medicum habebatur.  
Dum recens est odorem ingratum habet, acrisque sapore palatum ten-  
det; apte vero siccata Nilam odoratam suaviter olet. Hinc in re  
unguentaria apud veteres utilissima. Habetur etiam purgativa,  
expectorans, diuretica, erubina; praecipue vero siccata acris et  
corrodens est, alium violenter solvit, et in hydropi periculoso experi-  
mento quandoque utiliter propinatur. Flores recentis contusis  
in aqua ad putrefactionem digesti, dein calid commixti pulcherrimum  
colorem praebent, nomine St. illyricae Viride pictoribus notum.

Obs. Tridem, illyricam antiquorum spei St. germanicam L. jam monuit in  
Stimp. Dalm. sp. in p. 17: nihil tamen de eius Linnaeanum nomen reti-  
nendum spei censio, etiam si illo vetustissimo longe posterior et mini-  
me specificam, cum michi semper persuasissimum fuerit, ~~omnes~~  
~~illius~~ ~~spei~~ nomina vetera ante Sienkii regenerationem et philoso-  
phi ~~St. illyricae~~ ~~germanicam~~ plantis ~~spei~~ ~~indita~~, et multis ab hinc  
annis omnino oblita, <sup>per hunc</sup> <sup>in</sup> <sup>St. illyricae</sup> <sup>germanicam</sup> <sup>spei</sup> <sup>indita</sup> <sup>et</sup> <sup>multis</sup> <sup>ab</sup> <sup>hinc</sup>  
~~manus~~ ~~Sienkii~~ ~~progressibus~~ ~~confusa~~ (que synonymarum copia  
jamdudum confusa necnè pere obruta est), quam miseriam  
antitheticam, qua laborant, ostendere, rem jamdiu notam ab ipsis  
noviter restituta nomine salubandi. O. curas hominem!

\*\* perianthii laciniis reflexis in barbibus.

St. N. graminea L. sp. pl. p. 58. Herb. St. austr. p. 47. Nicandro. St. ex. 2

p. 82. Alach. fl. jard. p. 15.

*T. caule* subsiflora, ovipellis folijs linearibus ensiformibus rectis, multo brevioribus, gemminibus hexagonis.

Icon. Halg. fl. austr. 1. t. 2.

*T. angustifolia* W. Clud. hist. nat. p. 230.

Hab. Nees gen. pl. prec. v. pag. 4. 8 (p. 11.)  
in pratis subalpinis montis Nellibich. Fl. chajo. Flores dilute violacei, laciniis reflexis, rubentibus, prunas matura redo- lentibus. 7.

~~H. J.~~ pseudo-acoruf. L. sp. pl. p. 56. Hort. fl. austr. 1. p. 47. Reichb.

fl. exc. 4. p. 82. Alach. fl. jard. p. 15. Pelt. bot. voyag. n. 510.

*T. caule* multiflora folijs ensiformibus rectis longiore, perianthii laciniis crassis stigmata minoribus, gemminibus trigonis.

Jc. Blackw. herb. 2. t. 261.

Engl. bot. t. 578. Fl. num. t. 494.

Ulyr. Maxim. vadeni.

Hab. ad fossas, rivos, fluvios, paludes circa Knin, Sernis, Obdros atque Kara Spalato, Alonysal Marcitla. Fl. chajo. Flores lutei magni, laciniis basi mucosa crocea, striis fuscis. 7.

~~H. J.~~ tuberosa L. sp. pl. p. 58. Vit. Miss. Dalm. p. 38. Hort. fl. austr. 1. p. 49. Reichb. fl. exc. 4. p. 82. Alach. fl. jard. p. 15. Pelt. bot. voyag. n. 512.

*T. caule* uniflora, folijs linearibus trigonis, radice tuberosa.

Icon. Sibth. fl. gr. 1. t. 41.

Arch. bot. magak. 1. 531. Salt. bot. rom. 2. tab. 67.

Hab. in fruticibus dumetis herbidis asperis circa Kara, inter Trebacani et Ante pl. ab. Tomit. in Alach. me non prope pag. Fl. Monte Agiti. Pindus knini basi et inter fluvios, reflexi epia negri cornes, ovis pedes avete. 7.

Clusij autem plantae apud interij nomine Indij Dalmaticae  
iuncturae, scilicet 1°. Indij Dalmatica major Lobel. icon. italy. p. 61  
(quae Indij asiatica caerulea polyantha C. B. pin. 31. et Indij asiatica  
caerulea omnium Indium amplissima Dalmaticae affinis J. B. h. p.  
pl. 2. p. 216). 2°. Indij Dalmatica major C. B. pin. 31 (quae  
Indij Dalmatica major flore vulgati minus violaceo J. B. hist. pl. 2. p.  
215 et Indij chaledonia II Tabern. icon. 646). 3°. Indij Dalmatica minor  
Lob. icon. italy. 62 (quae Indij Dalmatica minor flore purpurea  
detractione J. Bauh. hist. pl. 2. p. 216. <sup>unum</sup> ipsissima Lobelii icona, et  
Indij latifolia major XVIII Clus. hist. rar. pl. p. 222 eadem icona <sup>quae</sup>  
sita partium tantummodo in verso. Quid sint duo priorae an  
species, an potius Indij germanicae variationes <sup>hottensis</sup> affirmare cupio  
quoniam in tota Dalmatia nulla Indij quae <sup>ipsa</sup> eisdem  
similis, praeter Dalmaticam hactenus inventa fuerit. Tertio  
vero in speciem propriam <sup>80</sup> recepta fuit a cl. Dausch (cfr. cl.  
Reichb. fl. ex. p. 89 et p. 140<sup>12</sup> n. 556.) Indij Clusiana  
nomine. Clusij quidem plantam per a Byzanti eidem  
misit (var. pl. hist. p. 221) ea tamen de causa Dalmaticam dixit  
se credo, quia hoc nomine inveniunt jam a Lobelio adumbatum  
quod si ex ipsius verbis non constat qui Lobelium nequidem nominat  
tamen luculenter evincitur ex icona <sup>ipso</sup> <sup>Clusii</sup> <sup>una</sup> quae ipsissima est cum icona  
Lobelii, <sup>partium</sup> <sup>tantummodo</sup> <sup>artificiosus</sup> <sup>inverso</sup>, ut sepe veritas facere  
conveniant. Nam vero plantae cl. Dausch caule plus accipiant et  
biflorae eadem <sup>fac</sup> <sup>est</sup> <sup>itaque</sup> Lobeliana et Clusiana, dubitatur licet sit



6.

9

[C. sativum L. in fl. ed. II p. 80 (excl. var. B.)  
Hort. fl. austr. 1. p. 42 Hort. fl. exl.

p. 85

C. ~~truncata~~ indicatibus tenuiter pubes. involucri,  
spatha floris monophylla, tubi faucis  
villus, stigmatibus trifidibus laciniis longioribus  
peltiformi-~~ibus~~ <sup>dentatis reflexis</sup> ~~involucris~~, stamine  
~~superantibus~~. Hort. hist. 1744 p. 441

Icon. Blakew. herb. 1. tab. 144 f. 1  
Mill. icon. pl. tab. III.  
Engl. bot. t. 343 (tab. 6. autumnalis)

25 Hab. in collibus circa Salona et in monte  
Mariano prope Spalato ~~inde misit~~  
Prof. Letter. An <sup>olim</sup> cultus? Flores violacei 4.

1/2 Illeg. Gajran. Denduska. Kupran.

Ufus Stigmata hejures plantae nonnullam  
aerariam, anodynam, dyminem emmenam  
virtutem possident, et aromate <sup>quo praestant</sup>  
et pulchritate hinc in condimentis  
~~adhibentur~~ <sup>adhibentur</sup>. Colorem prope luteum  
suffundunt.

Obs. Icon Lam. Enc. bot. ~~ed. 1. p. 30~~ tab. 30  
f. 1. haec a floribus dicti stigmatium  
breviter trifidis nec trifidibus, laciniis ~~superantibus~~ <sup>reflexis</sup>  
evidenter in nostris divergat.

Romulea

Ordo IX Amygdaleae

L. Brown prodr. fl. nov. Holl. p. 287

Floras speciosi: hemisphaerici <sup>hemisphaerici in capsula</sup> vel uterque ~~vel~~ solitarii aut umbellati, bracteis parvis inclusis.  
 Perianthium corollinum superum ~~caudum~~ sexpartitum aut hexaphyllum <sup>Staminibus x</sup> ~~Staminibus x~~ <sup>partibus parvis coronatum.</sup>  
 Stamina sex perianthiis ~~superis~~ opposita dno egypcio aut perianthio inserta, filamentis  
 liberis, aut basi in coram connatis, antheris inclusis, basi vel medio appis, locis parallelis,  
 longitudinalibus aut apice divergentibus. Pistillum stigmati tubulo aut ~~capite~~ stylo unico,  
 ovario infero trilobato <sup>multilocato</sup> ~~polygono~~, ovulis in loculorum angulo embolati biconvexis. Capsula  
 raro bacca, trilobatis ~~hinc~~ <sup>hinc</sup> lobulis medio apice ~~polygono~~ <sup>polygono</sup>, placenta ~~versis~~ <sup>versis</sup> ~~ovariis~~  
 centralibus. Semine anatrope testa membranacea aut correa, albumine caroso, embry-  
 one incluso axilli radicle hilum spectante.

Herbae bulbosae ~~perenni~~ <sup>perenni</sup> raro rhizoma perennans, suffruticeae, foliis radicalibus simplicibus  
 integerrimis linearibus ~~ovariis~~ <sup>ovariis</sup> ~~integerrimis~~ <sup>integerrimis</sup>, raro dentatis spinosis.

Clavis analytica generum

- |                                      |   |
|--------------------------------------|---|
| Perianthis intus nudo vid. N. 1.     | Perianthis intus coronato. vid. N. 4.   |
| 1. Radice bulbosa - - - - - 2        | Radice fibrosa  |
| 2. Florae pendulae - - - - - 3       | Agave   |
| 3. Perianthii phyllis aequalibus - - | Florae erectae  |
| 4. Staminiibus inclusis              | Sternbergia   |
| <u>Navisium</u>                      | Perianthii phyllis <del>inaequalibus</del> <sup>inaequalibus</sup> <del>diversis</del> <sup>diversis</sup> <del>brevioribus</del> |
|                                      | Salanthus   |
|                                      | Staminiibus <del>inclusis</del> <sup>exsertis</sup>   |
|                                      | Alnus   |

Agave L.



Ch. Perianthium infundibuliforme limbo subaequaliter sexpartito. Stamina ~~sex~~ perianthio  
 tubo inserta, filamentis liberis antheris <sup>ovario appis</sup> ~~centralibus~~. Stylus curv. apice perispermio, stigmati  
 apitato-trigono. Capsula coriacea trigona lobulata biconvexi, seminiibus ~~magis~~ <sup>magis</sup> ~~planis~~  
 compis, ~~testa~~ <sup>membrana</sup> ~~marginatis~~ <sup>marginalis</sup>, ~~testa~~ <sup>membrana</sup> ~~marginatis~~ <sup>marginalis</sup> - A. Hexandria Ord. Monogynia Syst. sex.

Agave americana L. ~~Flora~~ <sup>Flora</sup> ~~plansen.~~ <sup>plansen.</sup> ~~acut.~~ <sup>acut.</sup> ~~3. p. 22~~ <sup>3. p. 22</sup> ~~Sp. hist. delm.~~ <sup>Sp. hist. delm.</sup> ~~if. p. 23~~ <sup>if. p. 23</sup> ~~Naitz. fl. ex. p. 90~~  
 Aitch. fl. prodr. p. 81. ~~Bot.~~ <sup>Bot.</sup> ~~Wegw. n. 114.~~  
 A. phyllis flammis marginis et apice <sup>spinosi</sup>, corpa albiflora <sup>pyramidalis</sup> ~~perianthio~~ <sup>perianthio</sup> ~~ramulis~~ <sup>ramulis</sup> ~~apice~~  
 multi-convexibus, flantibus ~~hemisphaerici~~.



Hb. in part. leg. My alba <sup>in L. v. a</sup> ~~in L. v. a~~ <sup>in L. v. a</sup> ~~in L. v. a~~, et circa Polkovic <sup>in L. v. a</sup> ~~in L. v. a~~.

Hort sept. Oct. Hort <sup>lutei</sup> ~~lutei~~ 20.

Obs. In planta dlanica fls minus <sup>ae in hungaria</sup> ~~ae in hungaria~~ <sup>positivius</sup> ~~positivius~~ <sup>angustius</sup> ~~angustius~~ <sup>tubis</sup> ~~tubis~~ <sup>subaequalibus</sup> ~~subaequalibus~~, ut vete notavit  
cl. Scheuchzer; <sup>stylis</sup> ~~stylis~~ <sup>in utroque</sup> ~~in utroque <sup>longitudine</sup> ~~longitudine <sup>aeq.</sup> ~~aeq.~~~~~~

St. lutea Her. Neus. Amoy. p. 4 Hort fl. aust. 1. p. 429 <sup>Reich. fl. ex. p. 87.</sup>

Alch. fl. jav. p. 79 <sup>Pott. bot. weyn. u. 916</sup>

H. folij syriacis <sup>luteis</sup> ~~luteis~~ <sup>linearibus</sup> ~~linearibus~~ <sup>obovatis</sup> ~~obovatis~~ <sup>cauliculis</sup> ~~cauliculis~~ <sup>perianthio</sup> ~~perianthio~~ <sup>tubo</sup> ~~tubo <sup>duplo</sup> ~~duplo  
broad, lacinijs ellipticis <sup>obovatis</sup> ~~obovatis~~.~~~~

Syn. Amoy. lutea L. p. p. 420 <sup>Vif. Reg. del. p. 24</sup>

Non Sibth. fl. gr. IV. tab. 310

Cult. botan. mag. tab. 290

Clus. hist. pl. 1. p. 164

Hb. in saxij circa Zan, Schmis, Sardon, <sup>Prossin</sup> ~~Prossin~~ <sup>Deruy</sup> ~~Deruy, <sup>in monte</sup> ~~in monte <sup>Marian</sup> ~~Marian, <sup>Vergoz</sup> ~~Vergoz,  
u. Nagyp. Hort sept. Oct. H. lutei. 20.~~~~~~~~

*Leucosium* L.

Gen. p. 2. II. p. 113

Ch. *Perianthium* <sup>capitulum</sup> ~~capitulum~~ <sup>conspicuum</sup> ~~conspicuum~~ <sup>limbo</sup> ~~limbo~~ <sup>aequaliter</sup> ~~aequaliter~~ <sup>negatis</sup> ~~negatis, lacinijs <sup>ae-</sup> ~~ae- <sup>qualibus</sup> ~~qualibus,  
ut in part. <sup>lacinijs</sup> ~~lacinijs~~ <sup>seorsim</sup> ~~seorsim~~ <sup>perianthio</sup> ~~perianthio~~ <sup>parvi</sup> ~~parvi <sup>intra</sup> ~~intra <sup>perianthium</sup> ~~perianthium~~ <sup>lacinijs</sup> ~~lacinijs~~ <sup>liberis</sup> ~~liberis~~,  
antherijs <sup>basi</sup> ~~basi~~ <sup>affixis</sup> ~~affixis. Stylis <sup>longis</sup> ~~longis, <sup>stylis</sup> ~~stylis~~ <sup>simplicibus</sup> ~~simplicibus~~. Capsula <sup>broad</sup> ~~broad <sup>dura</sup> ~~dura <sup>etiam</sup> ~~etiam  
ovata <sup>lobulata</sup> ~~lobulata~~ <sup>stylis</sup> ~~stylis~~ <sup>remixtis</sup> ~~remixtis~~ <sup>globosis</sup> ~~globosis~~. — (C. Thunberg Or. Monog. 1. p. 148.)~~~~~~~~~~~~~~~~~~~~

*Leucosium caeruleum* L. p. p. 414. <sup>Vif. Reg. del. p. 29.</sup> Hort fl. aust. 1. p. 416. <sup>Reich.</sup>  
fl. ex. p. 87. Alch. fl. jav. p. 78. <sup>Pott. bot. weyn. u. 565</sup>

*L. spatha multiflora*, stylis <sup>clavatis</sup> ~~clavatis~~.

Non. Lam. enc. bot. pl. 230 f. 2.

o. Nagyp. fl. aust. tab. 203

a. Enst. bot. t. 621

(C. Thunberg gen. p. (Leucosium) No. VI

5. ~~Reich.~~ Bot. mag. t. 1210

~~Alchim. etc. etc. etc. etc. etc. etc.~~

Myo. *Spiza tubicola*. <sup>Linnaeus</sup> ~~Linnaeus~~.

Hb. prope in part. agris circa <sup>Seardon</sup> ~~Seardon~~ <sup>Non</sup> ~~Non, <sup>Deruy</sup> ~~Deruy, <sup>Knin</sup> ~~Knin, <sup>Trani</sup> ~~Trani, <sup>Castle</sup> ~~Castle, <sup>Salona</sup> ~~Salona,  
Almofa, <sup>Navarra</sup> ~~Navarra~~. Hort Agri. Majo. Hort alb. <sup>apertis</sup> ~~apertis~~ <sup>montibus</sup> ~~montibus~~ <sup>in</sup> ~~in <sup>montibus</sup> ~~montibus~~.~~~~~~~~~~~~~~

2.

## Galanthus L.

Gen. pl. ed. II. p. 113

Ch. Perianthium campanulatum, <sup>irregulariter</sup> sexpartitum, laciniis lobis interioribus brevioribus  
emarginatis, medio vecheriformis. Stamina perianthii parvis inserta aequalia, filamentis  
liberis, <sup>sub</sup> <sup>saepius</sup> <sup>in</sup> apice cuspidatis. Stylus filiformis stigmate simplex. Capsula  
ovata limbo breviter, seminibus subglobosis. — Cl. Hexandria Ord. Monogynia  
Syst. 148.

Galanthus nivalis L. sp. pl. p. 413. Herb. fl. austr. 1. p. 415. Fisch. fl. rar. p. 87

Sp. fl. <sup>carinata</sup>, <sup>ovata</sup> <sup>concolor</sup> <sup>caeruleis</sup> <sup>lobis</sup>, <sup>sepalis</sup> <sup>linearibus</sup>, <sup>perianthii</sup> <sup>laciniis</sup> <sup>exterioribus</sup> <sup>longioribus</sup> <sup>interioribus</sup> <sup>duplo</sup> <sup>superioribus</sup>.

Non. 2 Lam. Enc. bot. pl. 250

Engl. botan. tab. 19

poem. fl. europ. pro. v. ic.

Pey. fl. austr. t. 310

fl. dan. tab. 1285

1 C. F. Nees gen. pl. (Galanthus) p. VI

Herb. in demetris et praedictis circa Zagora et prope Cation. Pl. a Decemb. ad Februar.

fl. albi, laciniis interioribus <sup>maioribus</sup> <sup>caeruleis</sup> <sup>lobis</sup> <sup>longioribus</sup> <sup>exterioribus</sup> <sup>duplo</sup> <sup>superioribus</sup>.

## Narcissus L.

Gen. pl. ed. II. p. 113

Ch. Perianthium hypentriamorphum limbo subaequiliter sexpartito, <sup>ad</sup> <sup>saepius</sup> <sup>comum</sup> <sup>et</sup> <sup>paucis</sup> <sup>lobis</sup>  
paucioribus. Stamina sunt tubo <sup>inclusa</sup> <sup>obscure</sup> <sup>distincta</sup>, lobis exterioribus brevioribus, filamentis brevibus  
nig., antheris dorso affixis. Stylus filiformis, stigmate simplex. Capsula oblongo-triangulari  
breviter lobulata, debiscentibus, seminibus subglobosis. — Cl. Hexandria Ord. Monogynia  
Syst. 148.

Narcissus poeticus L. sp. pl. p. 414 Herb. fl. austr. 1. p. 416 Fisch. fl. rar. p. 88 Alch.  
fl. jav. p. 28 Bot. bot. weyer. u. bot.

fl. solij linearibus floris, saepe unicoloribus antheris, perianthii laciniis exterioribus longioribus  
<sup>superioribus</sup> <sup>exterioribus</sup> <sup>duplo</sup> <sup>superioribus</sup> <sup>interioribus</sup> <sup>inferioribus</sup> <sup>duplo</sup> <sup>superioribus</sup>

Non. 2 Bull. herb. 7. la. de. tab. 506.

Engl. bot. tab. 225 <sup>p. VI.</sup>  
C. F. Nees gen. pl. (Narcissus) p. 1-5

Illyr. 2 Parkers. pand. terr. p. 25 p. 1.  
Herb. in parvis montibus Velleth, Pomina, Solaja, Znan, Biskov, Vastiza etc.  
fl. April. fl. albi, corolla <sup>flava</sup> <sup>marginata</sup> <sup>reversura</sup>. X.





p. 32

Henry Milleri. *tbl. leg. et long. dist. 2. g. 5. 4. 10. 10 p. 28* *tbl. a. Lincea*  
 ible, *corolla unguis. cum illa ad foris videntur* *tbl. a. Lincea*  
 expiant perianthis formam ac longitudinem. *tbl. a. Lincea*

Ordo X. Liliaceae

*Del. th. elem. d. 1. p. 249*

Flaves hermaproditae *trianuly* *tbl. a. Lincea* aut racemif. aut cernitae, brachet. Liliaceae  
 collisera *tbl. a. Lincea* *tbl. a. Lincea* per minus setiform aut hexaphyllum fere videntur. Stamina  
 ut perianthis segmentis opposita sive intra ut hypogyn, plerumque oblonga, antheris interioribus  
 basi ut vixis affixis, loculis parallelis longitudinaliter delinquentibus. Pappiolum *tbl. a. Lincea*  
 triblo aut indur. *tbl. a. Lincea* stylo minus *tbl. a. Lincea* *tbl. a. Lincea* *tbl. a. Lincea* *tbl. a. Lincea*  
 ovulis in ludorum angulo biniatis. Capsula *tbl. a. Lincea* *tbl. a. Lincea* *tbl. a. Lincea* *tbl. a. Lincea*  
 vixis septifera, facientis *tbl. a. Lincea* *tbl. a. Lincea* *tbl. a. Lincea* *tbl. a. Lincea*  
 vixis. Semen *tbl. a. Lincea* testa membranacea aut mucosa, albumin exsuff. cuticula insula axili  
 indurata hilo proxima.

Herba herbacea aut terrosa scapigera aut caespitosa, plerumque in floribus *tbl. a. Lincea*  
*tbl. a. Lincea* *tbl. a. Lincea* *tbl. a. Lincea* *tbl. a. Lincea*

Classif. analytica generum

- |  |   |
|--|---|
| Radic bulbosa <i>tbl. a. Lincea</i> 3.       | Radic tuberosa aut fibrosa <i>tbl. a. Lincea</i> 10   |
| 1. Perianthis hexaphylo aut sexpartita 2.    | Perianthis sexfida aut sexdentata 9   |
| 2. Perianthis phyllis rectangulis 3          | Perianthis phyllis rotundis 4   |
| 3. Iovata nectarifera supra basin phyllosum  | Siles nectarifera longitudinali   |
| <u>Tritillaria</u>                           | <u>Lilium</u>   |
| 4. Stigmatis stylis instructis 5             | Stigmatis sessilibus  |
| 5. Perianthis phyllis ovatis 6               | <u>Dalila</u>   |
| 5. Phyllis oblongis cordatis 6               | Perianthis phyllis ovatis basi brachetis  |
| 5. Floribus corymbis aut racemis brachetis 6 | <u>Emphrasium</u>   |
| 6. Filamentis <i>tbl. a. Lincea</i> 7        | Phyllis <i>tbl. a. Lincea</i> 7   |
| 7. Antheris dorsis affixis                   | Phyllis <i>tbl. a. Lincea</i> <i>tbl. a. Lincea</i> <i>tbl. a. Lincea</i> <i>tbl. a. Lincea</i> |
| <u>Ornithogalum</u>                          | <u>Allium</u>   |
| 8. Perianthis ova aequalia aut ampliora 9    | Florum basi <i>tbl. a. Lincea</i>   |
| 9. Perianthis limbo patulo                   | Antheris basi affixis   |
| <u>Hyacinthus</u>                            | <u>Gagea</u>  |
|  | Perianthis ova contracta  |
|  | Muscari   |
|  | Perianthis limbo ovato  |
|  | <u>Bellevalia</u>   |

11. *Styctes indicus* — 12

12 *Scitibidius declivatus - asperdentatus*  
Asphodelus

Stylo  
~~Styctes indicus~~

Asphodelus  
*Scitibidius varius*  
Anthericum

10 *Styctes indicus* — 11

11 *Scitibidius varius*  
Anthericum

Trillium

*Styctes rubro-aurantivus* — 12

*Scitibidius declivatus - asperdentatus*  
Asphodelus

12 *Pentantherium phyllidivum - omnium - calicis*  
Asphodelus

*Phyllidivum omnium - calicis*  
Erythronium

Trillium L.

Gen. pl. ed. II p. 117

U. *Pentantherium corymbosum* <sup>indicum</sup> *asperdentatus* *calicis* *perle* *nervis* *oblongis*.  
lineari <sup>radice</sup> *sagittata* *prostrata*, *Stamina* *inter* *perianthia* *inserta* *et* *involuta*, *filamentis*  
*tri* *subulatis* *variis*, *antheris* *obtusis*. *Stylus* *variis* *Styctibus* *terminatus*. *Capitula*  
*Stylis* *trifloris*, *sepalis* *trifloris*, *obovatis* *marginatis* *obovatis* —

Cloje. Alexandri Ord. Maygini Sylb. n. 108.

Trillium montanum Hoff. pl. sic.

*P. foliis* *linearibus*, *inferioribus* *floribusque* *suboppositis*, *superioribus* *floribus*  
*subulatis*, *perianthiis* *lineari* *oblongis* *obovatis* *obtusis* *emarginatis*, *variis* *pallidis*,  
*perle* *nervi* *per* *strobilum*.

- Syn. *P. tenella* *Feichl. fl. ex. p. 102*
- P. pyrenaica* *Hort. fl. avst. i. p. 126. non L.*
- P. thalictroides* *Albt. fl. ind. p. 19. Bot. bot. wago. n. 119. non L.*

B. foliis *longis* *linearibus*, *caulis* *bifloris*.

Icon. P. racemosum a. Saub. bot. mag. tab. 1216.

*Hab.* *in* *provincia* *mediana* *Valleth. Provincia*, *Saraja*, *Belged. Viperovis. Mopar,*  
*Disthor.* *near* *ad* *domum* *Cajoviana: raritas* *provinciae* *cum* *non* *in* *provincia*  
*collibus* *montanis* *Stavimoron* *unde* *misit* *Neumannus* *et* *Calvisius.*

Obj. *Plantae* *non* *variae* *lehedinae*, *longitudine* *et* *dispartitione* *filamentis* *perle* *nervi* *ovarii*  
*specie*, *nisi* *inferius* *nisi* *superius*, *modo* *suboppositis* *suboppositis*. *In* *variis* *specie*  
*nullo* *characteribus* *a* *specie* *protra* *Styctibus*, *floris* *hinc*, *unde* *et* *specie* *caulis*





(Syl. Allium) Silla L.

gen. pl. 2. II. p. 119

Ch. Pennathium sexpartitum vobis-gattus perisperm. Stamina in periclythio infra  
filamentis patenti, subulic, antheris depresso medi affixis. Stylus filiformis, hinc  
obtus. Capsula oblonga trigona hinc apice delinque, loculis 1-4 spermiis, immixtis  
subulicis elongatis. — Cl. Hexandria Ord. monogynia Syst. nat. —

Silla athenensis L. sp. pl. 2. II. p. 443 Vindob. Delat. p. 42. Herb. fl. austr. 1. p. 434  
Pict. fl. exc. p. 106. Alch. fl. jard. p. 81. Pelt. bot. wege. n. 835.

S. filij ~~filiformis~~ <sup>ovata</sup> linearibus, floribus racemosis subulicis, pedunculis adscendens  
longitudinis filij, capsula oblonga aquilata.

Don. Avon. ic. III. tab. 274 f. 2.  
Uex. hist. stir. II. p. 185 f. 1, 2.

Hab. in collibus, et parvis saxosis circa Zano, Sebudo, Desio, Clima, Pappo, etc. in  
in insulis Doria, <sup>in montibus humilibus, Prussia, Masses, et in monte Silla.</sup> Florit  
Septembri, Octobri. H. ~~Septembri~~ 2.

Obi. Roman. Engl. bot. 78, et Bot. wege. 919 consulto omni, cum plantas  
nuper filij lat. linearibus, floribus racemosis a vasa hinc affixis.  
Alia ludant longis et brevibus, hystrochis, et squamulis (quae) tenui  
vix; pedunculis in antheris; floribus longis, racemosis filij subulicis; fructibus  
autem ovum filij <sup>in</sup> multo longior.

S. bifida L. sp. pl. 2. II. p. 443 Herb. fl. austr. 1. p. 434 Pict. fl. exc. p. 106 Alch.  
fl. jard. p. 81. Pelt. bot. wege. n. 836

S. filij subulicis linearibus longis racemosis apice <sup>subulicis</sup> conulicis, floribus  
racemosis <sup>subulicis</sup> subulicis, pedunculis erectis, capsula subulicis <sup>obtus.</sup>

Don. Jacq. fl. austr. t. 117  
Engl. bot. t. 24  
Gaem. fl. eur. f. 5  
Bot. wege. t. 746. f. text.

Hab. in umbrosis collibus, montibus Valleth, Prussia, Masses, Minerva,  
Shaba, Silla, Pastovich. Florit Martio, Aprili. Non comit. 2.

S. protensis Waldst. et Kit. fl. hung. II. p. 207 Pict. fl. exc. p. 106 et 110<sup>18</sup>  
S. filij trigonorum linearibus longis racemosis apice <sup>subulicis</sup> conulicis, floribus racemosis, hinc  
obtusis, oblongis, pedunculis <sup>subulicis</sup> subulicis, capsula subulicis <sup>obtus.</sup>



Icon. Allectio. herb. t. 1. 171  
Syn. Ornithogalum spicatum Jacq. in Bot. mag. 918  
Icon. Bot. mag. ii. 918

Class. herb. pl. p. 151 loco Torretta pro Spala  
Herb. Diolfa Kapelle.  
Herb. in saxoii agris circa Castellum pro Spala, capisse autem in igne  
Lilia, in Saxoii agris, circa Castellum pro Spala, capisse autem in igne  
Lilia, in Saxoii agris, circa Castellum pro Spala, capisse autem in igne  
Lilia, in Saxoii agris, circa Castellum pro Spala, capisse autem in igne

Herb. Dulcis lillae montanae in Saxoii agris et deperditis in igne  
aschant et catarrho pulmonum, tunc ferax et hygro vltis adhibetur sed pro  
pulvis, catarrho, hinc inde, vltis, catarrho, et igne. Herb. in Saxoii agris  
symptomata vltis, acerbis, propria esse vult.

(non in lilla) Allium L.

Gen. pl. ed. v. p. 143.

Ch. Penanthium hexapetalum vel septemlobum laciniis parvulis aut campanulatis con  
vinctis, papillis. Stamina in perianthio in tubo filamentis filiformibus, aut longis  
filamentis sublongis, alioquin interdu brevibus, antheris dorso nudo appressis. Stylus filiformis  
nigrescens, longior. Capsula tripartita aut triloculari, stylo communi, localiter mono  
dipertita, seminebus ovato-obovatis. — Class. Tourneforti Ord. Monogynia. Syst. 128.  
x Moly Doob. ind. alt. Leyd. bot. 2. p. 146  
herminibus demibus <sup>in Saxoii agris</sup> herbis

Allium moschatum L. sp. pl. 2. 2. 427 <sup>in Saxoii agris</sup> Herb. fr. austr. 1. p. 424. Scind. fl. <sup>in Saxoii agris</sup>  
excip. III. Arch. pl. jard. 78. Tab. bot. univ. n. 30.

A. filio bulbis oblongis tenuibus prope involucris obductis, sepalis inferne filis, filis papillis  
tenuibus, umbellatis, spatha bulbis lanceolatis acuminatis, umbellae  
pauca laxe, perianthio laciniis oblongis lanceolatis Stamina septemlobis  
filiformibus, longioribus, stylo communi, localiter mono  
dipertita, seminebus ovato-obovatis.

Syn. A. capiteum Cav. ic. pl. 3. p. 2. Tab. Bot. univ. n. 26  
A. salsuum W. et Kit. fl. rar. hung. 1. p. 50

Icon. Cav. loc. cit. 3. tab. 206 f. 1.  
W. et Kit. loc. cit. tab. 68  
C. Bauh. Pin. p. 28  
Scind. fl. ic. bot. v. p. 115

Herb. in altibus, laciniis saxoii agris circa Castellum, Spala, Cattaro, et in igne  
Drappa et Lurium. Flory alb. aut roseo, nervo laciniarum rubro, laciniis  
in Saxoii agris. Flory Aug. Sept. 24

A. saxatile Bieb. fl. tauricae in igne Besch. am Cay. near. p. 167 App. n. 39 et Flor. taur. can. 1. p. 264

A. bulbis oblongis tenuibus membranaceis obductis, sepalis inferne filis, filis subulatis  
tenuibus, umbellatis, spatha bulbis lanceolatis acuminatis, umbellae  
pauca laxe, perianthio laciniis oblongis lanceolatis Stamina septemlobis  
filiformibus, longioribus, stylo communi, localiter mono  
dipertita, seminebus ovato-obovatis.

Icon. Scind. fl. ic. bot. v. p. 5 fig. 596

Herb. in saxoij agarij, et vngaria in monte Sateja inter Dornij, et Biskero,  
et in montibz circa ~~Nasvato~~. Florit a Maio ad Julium. Heri gelidij, et in unum  
oboluceni, quodam tamen pulchre carum. 24.

Obs. Huc p[ro]p[ri]e cum cl. Koch in syn. fl. germ. p. 212, quoniam ad A. succulentam Paug.  
(ic. var. 2. p. 34 hb. 364) cum cl. Benthoni in fl. ibid. W. p. 49-50, pariter  
oboluceni flant ~~in~~ temperam a Monte Sateja et Opem, unius specimen

7. Item lecta cum vngariis omnino identice esse videtur. A. succulenta Paug. differt ab h[ab]itu  
folij floris <sup>ante in unum</sup> nec tenuibus, <sup>pubescentibus</sup> et h[ab]itu loboribz, umbellae globosae nec longioris.  
et praecipue spatula lobis lanceolatis acutis umbellae <sup>at Willdenowii Paugii</sup> ovatae brevior, nec apice in  
fistula solutae umbellae <sup>non sp[er]m[ati]t[is] p[ro]p[ri]e</sup> representat producta. Differt quoque ab A. obtusiloba  
Waldst. et Kit. fl. var. hung. 2. p. 204 a vngariis quod vngariis florum, et spatula  
et umbellae, nec non floribus duplo majoribus habet.

8. A. flexa ~~in saxoij~~ Schult. syn. vng. VII p. 102  
A. bulb. hujusmodi repenti in saxo <sup>h[ab]itu vngariis</sup> <sup>et pubescentibus</sup> repente uniusglobata, folijs tenuibus  
floribus oblongis oblongis, spatula bi-hilata pedunculata umbellae <sup>anguste</sup>  
longioris, ~~et~~ antheris exsertis

Syn. A. ~~anguste~~ var. ~~anguste~~ Hoff fl. austr. 1. p. 420. (Schult. fl. exc. p. 112 alt.)  
fl. jedr. p. 23. Bot. bot. vngar. n. 35 (var L.)

Var. A. ~~anguste~~ Hoff fl. 1. p. 196  
Schult. fl. bot. VIII hb. 235.

Herb. in saxoij, <sup>in saxoij</sup> ~~in saxoij~~ prope Zar et in saxoij montibus inter  
Dornij et Biskero. Florit Aug. Sept. Heri vngari, floris vix glomeratis. 24

Obs. Differt <sup>ab h[ab]itu</sup> ab A. succulenta L. sp. pl. ed. II. p. 430, quod tamen ~~h[ab]itu vngariis~~  
~~h[ab]itu vngariis~~ <sup>(et Schult. fl. bot. VIII hb. 235)</sup> ~~h[ab]itu vngariis~~ ~~h[ab]itu vngariis~~ ~~h[ab]itu vngariis~~  
h[ab]itu vngariis quod ex Linnaeo ipso, et et cithone Schult. fl. bot. I. p. 59. p. XI  
p. 2, est floris lobis, saepe angustis, folijs saepe h[ab]itu quadrato loboribz (et  
a Gmelino var. p[ro]p[ri]e, et vixi p[ro]p[ri]e Lich); denique in vngariis saepe lobulatum  
aequant, <sup>et</sup> intemp. floris, et floribus duplo majoribus.

9. A. nigra L. sp. pl. ed. II. p. 430. ~~h[ab]itu vngariis~~ Schult. fl. exc. p. 113. Potentilla.

<sup>vngariis</sup>  
A. bulb. ~~h[ab]itu vngariis~~ saepe h[ab]itu vngariis, <sup>h[ab]itu vngariis</sup> folijs lob. lanceolatis acuminatis floris  
floribus, spatula unilobata bi-hilata <sup>subrepente</sup> pedunculata, umbellae <sup>subrepente</sup>  
longioris, <sup>h[ab]itu vngariis</sup> ~~h[ab]itu vngariis~~ ~~h[ab]itu vngariis~~ ~~h[ab]itu vngariis~~

Syn. A. ~~anguste~~  
p. atropurpurea = pubescentibus atropurpureis Potentilla herb. edm.

Syn. A. atropurpurea Waldst. et Kit. fl. var. hung. 1. p. 16-17 Schult. fl. exc. p. 113

*A. veyron* Pet. bot. veyron. n. 31

Non. Waldst. et Kit. loc. cit. tab. 17

Hab. inter caetera <sup>in</sup> *Spalato*, <sup>et</sup> *Alpe*. <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
*Alpe*, <sup>in</sup> *caetera* <sup>in</sup> *Alpe*. <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

Obi. Plantae vid. *Chamaedry* <sup>non</sup> *Agria*, <sup>non</sup> *Alpe* <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
in *Chamaedry* <sup>non</sup> *Agria*, <sup>non</sup> *Alpe* <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

*A. subtriplex* L. sp. pl. ed. II p. 424. <sup>non</sup> *Alpe*. <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
Pet. bot. veyron. n. 37

*A. bulbis subrotundis*, <sup>non</sup> *Alpe* <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
*Alpe*, <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

Syn. *A. ciliolata* L. v. <sup>non</sup> *Alpe*. <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
*A. Aspicum* <sup>non</sup> *Alpe*. <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

Non. *Cypr.* loc. cit. tab. VI (col. pl. pl.)  
3. *locus* <sup>non</sup> *Alpe*. <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

1. *Denar.* loc. cit. tab. 151 f. 2. <sup>non</sup> *Alpe*.

Hab. <sup>non</sup> *Alpe* <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
*Alpe*, <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

*Flora* <sup>non</sup> *Alpe*. <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

Obi. <sup>non</sup> *Alpe* <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
*Alpe*, <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

1. p. 192 a. *L.* <sup>non</sup> *Alpe* <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

*Alpe* <sup>non</sup> *Alpe* <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
*Alpe*, <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

5. *A. micrum* L. sp. pl. ed. II p. 296. <sup>non</sup> *Alpe*. <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
*Alpe*. <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

*A. bulbis subrotundis*, <sup>non</sup> *Alpe* <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*  
*Alpe*, <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

Syn. *A. illyricum* Gay. ic. var. 2. p. 14

Non. *Gay.* loc. cit. tab. 365

*Sibth.* fl. gr. II. tab. 514

var. *β.* *bulbiferum*: <sup>non</sup> *Alpe* <sup>in</sup> *caetera* <sup>in</sup> *Alpe*, <sup>Flora</sup> *Agria* <sup>ma</sup> *Flora*

Syn. *A. caruicum* *Sibth.* fl. gr. I. p. 7  
*A. ambrynum* *Sibth.* fl. gr. II. p. 23

Non. *Sant.* *Ung.* in *Dase*? tab. VI  
*Sibth.* fl. gr. loc. cit. tab. 527

1 Bot. mag. ic. 978

1 Descr. fl. veg. 1. tab. 28

Herb. episcia <sup>virens</sup> in humidis et frigidis circa Zara, Sebaste, Spalato, Salona, Metesta, Ragusa  
see var. in insulis Page et Laticron, prostrata cum varietate. Floris vici. Floris  
Agulis, Tunis. 2.

Vis. 16 p. det. p. 24

b A. heterophyllum Cyr. pl. var. reag. I. p. 13, Reichb. fl. exc. p. 114 + in all. p. 140. n. 752

A. bulbis ovatis, saepe trigonis immo tetra. plus, plurius late linearibus aequantibus, cavatis, planis  
glabris, spatula uncinata acuminata pedunculatis duplo brevioribus, umbellae pappiformis, pendulis  
filis linearibus <sup>orb.</sup> <sup>ovatis</sup> ellipticis, <sup>nig.</sup> <sup>br.</sup> <sup>longioribus</sup>

Syn. a. album Sant. viing. ac montan. p. 352

Descr. fl. var. reag. I. p. 13

2 Sant. loc. cit. tab. VIII

1 Stth. loc. cit. IV. tab. 325.

Herb. ad ripas et humos habitum circa <sup>Trojan</sup> ~~Spalato~~ <sup>Cattaro</sup>. Flor. alb.  
Hort. Mart. Agr. Moja. 2.

7 A. Chamaemoly L. sp. pl. ed. II p. 439 Reichb. fl. exc. in all. p. 140. <sup>all. fl. juv. p. 11</sup>  
Pott. bot. vey. n. 28

A. bulbis ovatis, saepe hexagonis inclusa, plurius linearibus uniuersis ellipticis, spatula uncinata  
numeris pedunculatis <sup>linearibus</sup> <sup>parvis</sup> <sup>pendulis</sup> <sup>duplo</sup> <sup>brevioribus</sup>, umbellae pappiformis, <sup>linearibus</sup> <sup>pendulis</sup>

Descr. Column. Ephr. 1. p. 326

2 Martell. Pl. rar. et satun. p. 18 tab. 2

1 Bot. mag. tab. 1203

Herb. ad marginem agrorum et sepes circa <sup>Trojan</sup> ~~Spalato~~ <sup>Cattaro</sup> in insula Oros, circa Zara,  
et in insula Laticron. Flor. a Iuvenis ad Mart. Flor. alb. circa herbarium  
vicia. 2.

Obs. Non Caucasici in h. et Descr. fl. var. 3. tab. 207 p. 1 planta exsiccata saepe a plurius exsiccata,  
quod magnum videtur in siccis spontanea.

4 A. ussuriense L. sp. pl. ed. II p. 431 Hort. fl. austr. t. 2. p. 618 Reichb. fl. exc. p. 114. <sup>all.</sup>

A. bulbis oblongis, saepe trigonis hinc unguis, plurius <sup>longis</sup> <sup>parvis</sup> <sup>ellipticis</sup> <sup>linearibus</sup> <sup>uniuersis</sup> <sup>pendulis</sup> <sup>duplo</sup> <sup>brevioribus</sup>, spatula uncinata <sup>linearibus</sup> <sup>pendulis</sup> <sup>duplo</sup> <sup>brevioribus</sup>, umbellae pappiformis, <sup>linearibus</sup> <sup>pendulis</sup>

Descr. Engl. botan. t. 322

Flor. dan. tab. 757

Nep. Luch. Divlj.

Herb. in umbratis sylvarum maris Vellebita ad insulam Croatica, vicius lopy est Prof. Albinus.

Flor. Agulis, Tunis. Flor. alb. 2.

x. <sup>Ed. europ.</sup>

xx Codonopogon Turb. fl. exc. p. 111

9 a. uigra

Chimiechly amichy sifochy bei Kletk conary

A. flexuosum Sibth. et Sm. fl. gr. p. 1. p. 225  
Waldst. et Kit. fl. rar. hung. 3. p. 207

puta hules

A. saepe teni ad medium filio, filij linearibz serrulatis, canaliculatis, umbellam  
efform. supra rugositate, perianthii: obovato-trigoni linearis oblongis, communi 5, raris  
sem. perianthio duplo longioribz.

alt. a. umbella capillifera

Vis. iteg. hung. p. 24

syn. A. paniculatum, Herb. fl. aust. 1. p. 424. Gard. fl. helv. 2. p. 190  
Codonopogon paniculatus Turb. fl. exc. p. 115 (excl. syn. L. paniculatus)

non Reichb. icon. bot. v. tab. 418

Gard. fl. helv. 2. tab. XIII f. 12

beta. umbella bulbifera

A. curvatum Alch. fl. jard. p. 77. Bot. bot. wagn. n. 27

syn. A. flexuosum Waldst. et Kit. fl. rar. hung. 3. p. 207 Reichb. ic. bot. v. p. 10.

A. flexuosum Herb. fl. aust. 1. p. 422

Codonopogon flexuosus Turb. fl. exc. p. 116

non Waldst. et Kit. loc. cit. tab. 278

Reichb. ic. bot. v. tab. 603

in agris circa Zarn et Salva et in Romia, Silesia, Bohemia, non isolari. Hung.

Hab. utroque varietate in dumis montanis, Romania, Silesia, Bohemia, non isolari. Hung.

aeolate 2

Obs. Allium paniculatum L. sp. pl. ed. II p. 428 ab utroque varietate

see duplo longioribz. Cfr. Don monogr. all. p. 44 et Koch syn. fl. germ. p. 220.

non Haller's opus. p. 260 a L. ubi ab Allium perianthio longior, et perianthio depressione minus hab.

11 A. intermedium Lam. et DC. fl. fr. 5. p. 318! ex DC. herb.

A. caule teni ad medium filio, filij linearibz serrulatis, canaliculatis, supra hules

umbellam efform. supra rugositate, perianthii: oblongis linearibus oblongis, communi 5, raris  
sem. perianthio duplo longioribz.

syn. A. paniculatum Alch. herb! ad fl. jard. p. 72 (excl. syn). Bot. bot. wagn. n. 33.  
Codonopogon intermedium Turb. fl. exc. p. 116.

Hab. in saxo, in dumis, ad rivos circa Zarn, Desviz, Sabinu, Salva, nec non in sylvis, non

hinc a Romania, Silesia. Varietas promiscua est et cum specie. Hung. var. H. agr.

2.

Obs. Nam hoc varietate differt ab A. paniculato L. Don et Koch, <sup>valde dubium</sup> ab utroque varietate, cum utraque

habeat a ut. cordibus adhaerentibus speciebus hanc diffinitionem congruat A. intermedium et A. paniculatum

non et auctores, non A. curvatum Sm. Chimiechly utroque perianthio longioribus, non uno ut genus

in specie longioribus perianthio longioribus hinc a Romania a L. in rivo differt.









22 *A. viride*  
*A. myrsinaceum* Sibth. et Sm. p. 226  
*A. bulbis* <sup>aut agrostis</sup> *viridis*, *caulis* *strictus* *et* *mediis* *placis* *tenis*, *foliis* *terribus* *spiralibus*, *spatha*  
*unicuique* *longissima* *intra* *umbellam* <sup>interfolia</sup> *cauliculis* *superius* *gerens*, *gerens*  
*hinc* *partim* *longioribus* *lanceis* *oblongis*, *staminibus* *longioribus*.

Icon. Sibth. fl. gr. IV. tab. 315

Herb. in agris circa *Spagna* *veritas*, ubi *loco* *et* *communi* *Doc. Clarend.* *Herb.*  
*alibi*, *in* *via* *publi*, *inter* *loca*. *Herb. Vern.* 4

861. *Huc* *A. scrophularia* *Juss.* *in* *Act. Soc. Boton.* *just* *specimen* *authenticum* *in*  
*herb. ab. Del.* *Ne* *immixta* *l. Scrophularia* *in* *fl. Ind. W. p. 23*  
*habent* *A. scrophularia* *D'Urv.* *quod* *ab* *illo* *herbo* *Scrophularia* *gerens* *hinc* *caulis*  
*in* *media* *caulis* *inter* *aut* *per* *umbellam* *notari*. *Spatha* *in* *recessu* *longioribus*

*caudicum* *puberulis*; *capis* <sup>hinc</sup> *puberulis* *gerens*. *Habitu* *gerens* *A. rotunda* *quam* *A.*  
*scrophularia* *refert*: *Apud* *ab* *illo* *foliis* *non* *terribus* *spiralibus*, *spatha* *longioribus*  
*et* *forma*, *umbellam* *laxioribus*, *staminibus* *longioribus*; *ab* *hoc* *bulbis* *intra* *hinc* *compositis*  
*aut* *omnis* *inferi*, *non* *extis*, *oblongis*, *foliis* *terribus* *spiralibus*, *umbellam* *laxioribus*  
*foliis* *duplo* *minoribus*, *staminibus* *gerens* *hinc* *longioribus* *umbellam* *laxioribus*,  
*capitulis* *longioribus* *inter* *aut* *per* *umbellam* *notari*.

Fl. Ind. W. p. 23

23 *A. viride* *L.* *sp. pl.* *ed. II.* *p. 428* *Herb. fl. austr.* *I.* *p. 422*, *alibi* *fl. Ind. W.* *p. 26*  
*A. bulbis* <sup>ad</sup> *umbellam* *viridis*, *caulis* *strictus* *et* *mediis* *placis* *tenis*, *foliis* *terribus* *spiralibus*, *spatha*  
*unicuique* *umbellam* *longissimam* *intra* *umbellam* *cauliculis* *superius* *gerens*, *gerens*  
*hinc* *partim* *longioribus* *lanceis* *oblongis*, *staminibus* *longioribus*.

Syn. *Comen. arvensis* *part. fl. ex. p. 110*

Non. Engl. bot. t. 1934  
 1. *part. fl. ex. v. p. 110*  
 2. *part. fl. ex. t. 1934*  
 3. *Michx. bot. gr. fl. t. 24. f. 1*

*β. compacta*: *umbellam* *floribus* *in* *caulis* *bulbis* *omnis* *caulis*

Non. *Del. ic. shop* *p. 155. f. 2.*

Herb. *in* *cultis*, *et* *bulbis* *intra* *caulis* *caulis*, *et* *spatha*, *non* *umbellam*. *Herb. Vern.* *Non. austr. 4.*





Oenothera biennis L. in Kuhn's botan. et ill. Wiln. et d. v. 1. p. 1701 n. 53  
videtur ipse O. conium multum plii angustioribus...

3 O. exasperata Torr. fl. veg. 1. p. 175 ~~Fuchs~~ fl. ex. p. 109  
O. plii <sup>flor. glaucis</sup> <sup>multis</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
caulis <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
radix, apice <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
radix, apice <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>

Jan. Ten. fl. veg. 1. p. 175  
Syn. O. ~~reputans~~ <sup>flor. glaucis</sup> <sup>multis</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
Hab. in saxoia agris circa ~~Wismar~~ <sup>Wismar</sup> <sup>Wismar</sup> <sup>Wismar</sup> <sup>Wismar</sup> <sup>Wismar</sup> <sup>Wismar</sup> <sup>Wismar</sup>  
Habitat. Not. plii, quae ab ista ~~differt~~ <sup>differt</sup> <sup>differt</sup> <sup>differt</sup> <sup>differt</sup> <sup>differt</sup> <sup>differt</sup> <sup>differt</sup>

~~Ch. B. fl. veg.~~  
Obs. Scapus <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
longioribus <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
Nota. Nota <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>

5 O. ~~variegata~~ <sup>variegata</sup> <sup>variegata</sup> <sup>variegata</sup> <sup>variegata</sup> <sup>variegata</sup> <sup>variegata</sup> <sup>variegata</sup>  
Alch. fl. veg. 1. p. 175 ~~Fuchs~~ fl. ex. p. 109  
Alch. fl. veg. 1. p. 175 ~~Fuchs~~ fl. ex. p. 109

O. plii <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
radix, apice <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
Syn. O. ~~reputans~~ <sup>flor. glaucis</sup> <sup>multis</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
Nov. 2. <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>

Ch. ~~hist. fl.~~ <sup>hist. fl.</sup>  
3 Engl. bot. t. 1. p. 199  
2 Jacq. fl. austr. 2. p. 103

Reichb. icon. bot. X. p. 1225  
Hab. in ~~Mitteleuropa~~ <sup>Mitteleuropa</sup> <sup>Mitteleuropa</sup> <sup>Mitteleuropa</sup> <sup>Mitteleuropa</sup> <sup>Mitteleuropa</sup> <sup>Mitteleuropa</sup> <sup>Mitteleuropa</sup>  
Fraxina, ~~Krassia~~ <sup>Krassia</sup> <sup>Krassia</sup> <sup>Krassia</sup> <sup>Krassia</sup> <sup>Krassia</sup> <sup>Krassia</sup> <sup>Krassia</sup>  
Habitat. Not. 2.

### Gagea salicifolia

Labr. Ann. of bot. II p. 555

Ch. ~~Pentstemon~~ <sup>Pentstemon</sup> <sup>Pentstemon</sup> <sup>Pentstemon</sup> <sup>Pentstemon</sup> <sup>Pentstemon</sup> <sup>Pentstemon</sup> <sup>Pentstemon</sup>  
radix, apice <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
Cappula <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
ad. <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>

Gagea ~~ovata~~ <sup>ovata</sup> <sup>ovata</sup> <sup>ovata</sup> <sup>ovata</sup> <sup>ovata</sup> <sup>ovata</sup> <sup>ovata</sup>  
Schult. 1. p. 547 Reichb. fl. ex. p. 108

G. ~~helle~~ <sup>helle</sup> <sup>helle</sup> <sup>helle</sup> <sup>helle</sup> <sup>helle</sup> <sup>helle</sup> <sup>helle</sup>  
radix, apice <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>  
oppositis, <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup> <sup>longioribus</sup>

Syn. ~~Oenothera~~ <sup>Oenothera</sup> <sup>Oenothera</sup> <sup>Oenothera</sup> <sup>Oenothera</sup> <sup>Oenothera</sup> <sup>Oenothera</sup> <sup>Oenothera</sup>  
O. ~~minorum~~ <sup>minorum</sup> <sup>minorum</sup> <sup>minorum</sup> <sup>minorum</sup> <sup>minorum</sup> <sup>minorum</sup> <sup>minorum</sup>  
Nov. Pers. bot. cit. 1. p. 110  
H. ~~dem.~~ <sup>dem.</sup> <sup>dem.</sup> <sup>dem.</sup> <sup>dem.</sup> <sup>dem.</sup> <sup>dem.</sup> <sup>dem.</sup>

Silph. fl. græc. IV. tab. 332  
 Hb. in uulgi circa Zava, Sabinis, Sylab. Heros flori exly vivdy. Heros Robur, Agri.  
 Obs. Nis ludunt villos, siliis, aut omnino glabr, perianthis laciniis glabræ vel exly villosis, 2  
 oblongis aut mucronatis.

*S. lutea* pusilla Schult. syst. veg. VII. part. 1. p. 543 (Sylab. fl. ex. p. 107)

*S. lutea* siliatis ovata, plis radiatis unice laciniis <sup>angustis</sup> bise canalibus - longioribus, floribus <sup>5-6</sup> vivdy laciniis  
 laciniis, saepe <sup>5-6</sup> dubitans, pediculis <sup>5-6</sup> hybridis <sup>5-6</sup> glabratis, glabris, perianthis laciniis oblongis-lanceolatis  
 longis.

Syn. Orusthyge pusillæ Schmidt fl. botan. unt. II. p. 41 (Sylab. fl. ex. II. p. 8. Alth. fl. jud. p. 26)

*S. Clusii* Tausch in Bot. Zeit. 1828 p. 431

*Saga Clusiana* Schult. loc. cit.

Icon. Reichb. Ic. bot. II. f. 228, 229. opt.

Clus. hist. pl. II. p. 189

Hb. exly vivdy circa Zava, et in babilij montij Alth. Heros flori exly vivdy. Heros Robur, Agri.  
 Masi in Alth., Agri. Masi in Alth. 24

Obs. Variet saepe uni-bifloris, pediculis <sup>5-6</sup> vivdy, <sup>5-6</sup> longioribus, pediculis <sup>5-6</sup> vivdy, pediculis <sup>5-6</sup> vivdy, pediculis <sup>5-6</sup> vivdy,  
 plis <sup>5-6</sup> vivdy: hinc videntur <sup>5-6</sup> vivdy, quibus *S. pusilla* et *S. Clusiana* distiguntur Alth.  
 minima <sup>5-6</sup> vivdy.

*S. pygmaea* Schult. loc. cit. p. 547

*S. lutea* siliatis ovata, plis radiatis subgeminis limbo-<sup>5-6</sup> vivdy canalibus, floribus <sup>5-6</sup> vivdy  
 canaliculatis unilobis, pediculis <sup>5-6</sup> vivdy, perianthis laciniis oblongis-lanceolatis

Syn. Orusthyge pygmaea W. En. h. bot. p. 367 Alth. fl. jud. p. 25

Hb. in coll. apertis circa Zava, ubi legit et communicavit prof. Alth. Heros flori exly vivdy.  
 Heros Masi. 24

Diff. Planta uni-bifloris. Nis ut glandosum bina, interdum unicum. Caulis teris <sup>5-6</sup> vivdy  
 plis <sup>5-6</sup> vivdy, basi glabr, inter plis <sup>5-6</sup> vivdy. Nis tri alba laciniis - attenuata angustis  
 canaliculatis unilobis - Pediculis <sup>5-6</sup> vivdy, ut glandosum plis <sup>5-6</sup> vivdy. Perianthis  
 laciniis oblongis <sup>5-6</sup> vivdy, <sup>5-6</sup> vivdy, <sup>5-6</sup> vivdy, <sup>5-6</sup> vivdy. <sup>5-6</sup> vivdy  
 longis, pediculis, et perianthis <sup>5-6</sup> vivdy exly laciniis.

Mesasi

Journ. inst. v. herb. t. 180

Alth. Perianthis subglobosis-helipis <sup>5-6</sup> vivdy, ovae <sup>5-6</sup> vivdy, <sup>5-6</sup> vivdy. Stamina perianthis  
 hinc vivdy et involucris, filamentis <sup>5-6</sup> vivdy, <sup>5-6</sup> vivdy, <sup>5-6</sup> vivdy. Stylus vivdy <sup>5-6</sup> vivdy  
 oblongis. Capsula <sup>5-6</sup> vivdy, <sup>5-6</sup> vivdy, <sup>5-6</sup> vivdy. <sup>5-6</sup> vivdy. <sup>5-6</sup> vivdy. <sup>5-6</sup> vivdy. <sup>5-6</sup> vivdy.  
 Journ. Syst. 108.





Hab. ad marginem agrorum, et in salsis circa Dikla prope Zam, Dionysia prope Toni, et in Nivion dei Cappuli. Flores albi, extra lenticulae qualescentia, <sup>ant. et gub. antheris</sup> ~~antheris~~ fungini, psicis. Plant. Lycol.

D. Dubia Reichb. fl. ex. p. 105

D. breviter exagantulata, perianthio emissis, expis laciniis orbato-rotundis, filamentis longis, obovatis, marginibus obtusis, caulis glauco, apice subulato, capitulo profunde amovibili.

Syn. Hypocistis Dubius Jussieu. cat. Hort. Botanicif. 1821 p. 92 et 98.

Hab. in herbis circa Lesion, ubi communicavit profr. Italia. Flora imperialis comitatus, regnum abdo. insensit, ante et post antheris sordid viridis. Hort. Mart. Agri. 7.

Antherium L.

gen. pl. 2. II. p. 119 (excl. spec.)

Ch. Perianthium hexapetalum patens profundum. Stamen receptaculo inserto, filamentis filiformis; lobis patulis, antheris oblongis parvis non benè affixis. Stylus tertius trigynus subcapitato. Capitulum obtusum trigynum apice trilobis, lobulis 2-biserviatis, seminibus antipodibus dorso cavatis. —

Cl. Hexandria Ord. Monogynia Syst. sex. —

Antherium Lilium L. sp. pl. 2. II. p. 445 Hort. fl. austr. 1. p. 438 Reichb. fl. ex. p. 117 et 118 fl. jard. p. 82

A. filix concaudulata, sepalis simplicibus, stilo declinato

Syn. Phalargium Lilium Lam. enc. bot. 2. Patro. v. p. 231

Asphodelium Lilium Reb. bot. voyag. n. 93 (nec L.)

Don. Lam. 1. c. pl. 240 f. 2.

Flor. den. tab. 66b.

Jay. hort. ind. tab. 83.

Solan. may. tab. 914

Hab. in herbida in sub lago, circa Nagyndi, et Nagyndi, non in montibus Vellekth, Polyst, Biskov. Flora alba Hort et Majo ad Aug. 24.

A. variflorum L. sp. pl. 2. II. p. 445 Hort. fl. austr. 1. p. 438 Reichb. fl. ex. p. 117 et 118 fl. jard. p. 82

A. filix concaudulata floribus subulatis, sepalis parvis paniculato ramis, stilo recto.

Don. Jay. fl. austr. tab. 161.

Flor. den. tab. 1197

Loth. icon. 1. fig. p. 42 f. 2.

Hab. in alibus saxosis agris circa Dionysia, et alibi in Illyria montana. Flora Novi, Augusti. 24

Asphodelium

Reichb. fl. ex. p. 116

Ch. Perianthium sexpetalum patens profundum. Stamen receptaculo inserto, filamentis declinatis confundendis, lobis obtusis, antheris oblongis parvis non benè affixis. Stylus tertius trigynus subcapitato. Capitulum obtusum trigynum apice trilobis, lobulis 2-biserviatis, seminibus antipodibus dorso cavatis. —



*A. pedunculata* Lamour. L. p. 444. Hist. fl. austr. 1. p. 436 Reichb. fl. ex. p. 117  
alt. fl. ped. p. 61. Bot. bot. w. 127

*A. radice tuberosa*, filij late linearibus caudatis, scapo vixto, densifloro, bracteis im-  
punitis pedunculis subaequalibus, <sup>floribus</sup> glabris et basi subrotundis, ab angulo filiformibus.  
capsula subglobosa lobata hexagona

Gen. Sibth. fl. gr. IV. tab. 334

Sonn. mag. 984

Gay. austr. tab. 161.

Ches. left. fl. 2. p. 196 f. inf.

Myz. Caprij, Caprij, Caprij, Novodak.

Hab. in saxositate apertis hinc Delonchiae litoralis, probabile cum repente commixta.

Flor. albi, <sup>lucida</sup> linae pers. rubrae et hinc Ximoyi. Flor. April. Maji. 2.

*A. neglecta* Schult. Syst. veg. VII. p. 488-89. Reichb. fl. ex. p. 117

*A. radice tuberosa*, filij late linearibus caudatis, scapo vixto, <sup>densifloro</sup> densifloro, bracteis  
impunitis pedunculis duplo longioribus, <sup>floribus</sup> glabris et lanceolatis.

Hab. in saxositate saxositate <sup>lucida</sup> linae, unde vixto glab. <sup>et in</sup> Flor. <sup>et in</sup> April. 2.  
cum pro vixto et ceteris etiam albi, et quoniam in insulis Flor. April. 2.

Obs. Specij habita *A. venosa* K. nullum a qua tamen characteris hinc in dno  
Bary florum in tui glabris, marginibus vixto. <sup>quibus</sup>  
saxi Ximoyi. Nihil differunt a capsulae forma et angulis patet, et ab  
autem in dno, <sup>etiam</sup> mibi minimis covetis vis. int.

*A. alba* Mill. the good. Dict. n. 3 Hist. fl. austr. 1. p. 437 Reichb. fl. ex. p. 117

*A. radice tuberosa*, filij late linearibus caudatis, scapo singulis aut big. subaequalibus, <sup>densifloro</sup> vixto  
bracteis impunitis, pedunculis subaequalibus, <sup>floribus</sup> glabris et basi oblongis  
filij lanceolatis, <sup>etiam</sup> hinc

Gen. Blackw. herb. 2. tab. 238

Ches. left. fl. II. p. 192 f. int.

L. F. Nees gen. fl. IV. (Cappadocia)

Hab. in saxositate montis <sup>lucida</sup> linae alt. 2500 ad 3000 ped. Flor. Julio. Flor. albi, linae  
abopugena. 2

Obs. Variet. <sup>lucida</sup> linae vixto basi vixto cum altioribus autem, sed cum in forma <sup>quibus</sup> <sup>quibus</sup>  
abit. Differt in qua vixto ab *A. neglecta*, vixto nec characteris nec habitus vixto.

A. *fruticosa* L. sp. pl. 2. p. 444 Vif. 1. King. Del. 27 Herb. fl. austr. 6. p. 436  
Nicht-fl. austr. p. 117 Alch. fl. austr. p. 81 Pitt. bot. austr. 1. 125

A. *indica* (Poir.) Planch. *indica* subulosa, fimbriata, capsulae immixtae, breviter imbricatis, utrinque pedunculata, fructibus deorsum breviter, flosculis & lili. ovae magis profunde subulatis.

Non. Chos. icon. pl. no. III. p. 202  
Sibth. fl. gr. IV. tab. 355  
Chos. hist. pl. II. p. 197 f. ext.

Herb. in lecturis in Zam. Schina, Tami, Sydes, Linn, Niger. Flory albi, utriusque  
rubri. Hort. <sup>avili</sup> Maj. 2.

### Erythronium L.

gen. pl. 2. p. 118

Ch. *Pinnatifida* hexagone, capsulae reflexae, perianthio, glabro, inferne communi, & utriusque latere. Stamina extrinseca, tubum glabrum, perianthii inferni, filamentis brevibus, subulatis, cunctis, antheris brevibus. Stylus filiformis, hinc utriusque capsulae breviter, loculis glabris, cunctis, ovae breviter, paulatim appendicibus. — Claf. Heder. Ord. Monog. 1. p. 118.

Erythronium *Davis* var. L. sp. pl. 2. p. 437 Herb. fl. austr. 6. p. 427 Nicht-fl. austr. p. 102 Alch. fl. austr. p. 80 Pitt. bot. austr. 379

E. *plenum* (L.) Willd. capsulae breviter, utriusque perianthii communi.

Non. Curt. bot. mag. tab. 5  
Nag. fl. austr. opp. tab. 9  
Lam. Enc. bot. pl. 244 f. 1

Op. Haeg. gen. pl. fasc. IV. (Erythronium)

Herb. in sylvarum montium Vellebit. Vinnisija, Dalgat, Doshan, et supra Pithul. Flory rubri, utriusque latere. Hort. Apr. Maj. 2.







in *Colobium leucum* nomen, quod de Jussieu apparuit subdit in fl. Ind. Ind. p. 452  
 anno 1827 edito <sup>recepto Schumier</sup> ~~et ab Jussieu antea~~  
 Jussieu de campis specij quibus a d. Schumier pro C. montano ~~proposita~~, hinc  
 tunc reformi regit, etiam si venia a plants esse Jussieu, quod altero d. Schumier  
 non hinc regit, quibus ~~est~~ has plantas <sup>in det. mag. 1827 p. 41-42</sup> ~~et~~ Jussieu non minus quibus videtur  
~~cum~~ bulbis sicut, d. Schumier, et hinc hinc regit, plus hinc magis etiam, ut  
 color ut etiam hinc, etiam hinc a plants esse Jussieu, quod altero d. Schumier  
 cum ~~antiquior~~ fuit, etiam in eadem. Hinc addit C. montanum ~~det. fl. Ind. Ind.~~  
 p. 322 ex inspectione hinc Jussieu a d. Jussieu in hinc etiam in eadem  
 nomen, quamvis a d. Schumier ad C. montanum nomen adducatur; addit insuper  
 a nomen a d. Jussieu <sup>recepto</sup> ~~recepto~~ Jussieu botanica proposita, fl. Ind. Ind.

in ~~recepto~~ hinc Jussieu nomen, quod a nomen a plants esse Jussieu, quod altero d. Schumier  
 tunc hinc regit, quibus <sup>quod inter hinc invenit etiam in eadem</sup> ~~est~~ has plantas <sup>recepto</sup> ~~et~~ Jussieu non minus quibus videtur  
 color ut etiam hinc, etiam hinc a plants esse Jussieu, quod altero d. Schumier  
 cum ~~antiquior~~ fuit, etiam in eadem. Hinc addit C. montanum ~~det. fl. Ind. Ind.~~  
 p. 322 ex inspectione hinc Jussieu a d. Jussieu in hinc etiam in eadem  
 nomen, quamvis a d. Schumier ad C. montanum nomen adducatur; addit insuper  
 a nomen a d. Jussieu <sup>recepto</sup> ~~recepto~~ Jussieu botanica proposita, fl. Ind. Ind.

Plantam de qua hinc Jussieu nomen, d. Jussieu <sup>recepto</sup> ~~recepto~~ Jussieu botanica proposita, fl. Ind. Ind. p. 322  
 Vingt. Vingt. 1826. p. 375) Jussieu nomen in ~~recepto~~ Jussieu botanica proposita, fl. Ind. Ind. p. 322  
 tunc hinc regit, quibus <sup>recepto</sup> ~~est~~ has plantas <sup>recepto</sup> ~~et~~ Jussieu non minus quibus videtur  
 color ut etiam hinc, etiam hinc a plants esse Jussieu, quod altero d. Schumier  
 cum ~~antiquior~~ fuit, etiam in eadem. Hinc addit C. montanum ~~det. fl. Ind. Ind.~~  
 p. 322 ex inspectione hinc Jussieu a d. Jussieu in hinc etiam in eadem  
 nomen, quamvis a d. Schumier ad C. montanum nomen adducatur; addit insuper  
 a nomen a d. Jussieu <sup>recepto</sup> ~~recepto~~ Jussieu botanica proposita, fl. Ind. Ind.

Ordo XII Trilicaceae

fl. Br. Prodr. fl. nov. Hol. p. 292

Herb hermaphrodit. ut absteris unisexualibus axillaris vel terminalibus, pedicellis 1/2-petiole articulis; brachylophyis. Perianthium corollae inferum sexdigitum, rurs 1-8-partitum, fereum medium. Stamina perianthio brevius aequalia opposita corollae bafi inserta: filamentis ~~liberis~~ non ita brevibus, antheris brevibus vel longioribus, lacinis longioribus longitudinaliter subdivisibilibus. Pistillum breviter stipitatum, stylo 2-5-partito, ovario ovum 1-2-fidum, bivalve, lacinis non bi-quadribus, ~~stipitatis~~ stylo brevioribus aequalibus uniserialibus, subinde ~~liberis~~ stipitatis 2-partitis. Pericarpium carnosum induratum bi-vel tri-quadriculatum, lacinis digitiformibus, non articulis, monopermis. Semen subglobosum aequale locum in centro affixum, testa membranacea, albumina subcornea, embryo minimus in albumine curvato inclusus.

Herbes stipitate perennantes, vel perennantes, foliis alternis subrotundis, simplicibus integris, nervosis, interdum 1/2-pinnatis.

Clavis analytica generum

Planta et

floris annuis 6 Vol. N. 1. floris annuis 8-10

- |   |  |
|---|--|
| 1. <u>Staminibus</u> liberis — 2  | <u>Staminibus</u> <sup>Pan's</sup> <u>monadelphis</u>  |
| 2 <u>Perianthio</u> hexaploco 12-partito — 3  | <u>Perianthio</u> breviter sex-fido <del>liberis</del> |
| 3 <u>Perianthio</u> longioribus <sup>bafi</sup> <u>longioribus</u> <del>hexaploco</del> | <u>Perianthio</u> <sup>ovariis</sup> <u>obtusis</u>    |
|   | <u>Perianthio</u> <u>patente</u> <u>reperto</u>        |
|   | <u>Smilax</u>  |
| 4 <u>Perianthio</u> <u>complicato</u>   | <u>Perianthio</u> <u>lobato</u>                        |
| <u>Convolvulus</u>  | <u>Delphinium</u>                                      |

Paris L.

Sm. fl. 2. II. p. 139

Un. floris hermaphrodit. Perianthium tubulatum och-decaphyllum pentiformi perispermio. Stamina och-decem, filamentis brevibus vel longioribus et perianthio brevibus, antheris ovum ultra articulum aequum in aequum locum productis. Pistillum ovum quadrangulatum, stylo 4-5-partito, ovario ovum 1-2-fidum, bivalve, lacinis non bi-quadribus, stipitatis 2-partitis. Pericarpium carnosum induratum bi-vel tri-quadriculatum, lacinis digitiformibus, non articulis, monopermis. Semen subglobosum aequale locum in centro affixum, testa membranacea, albumina subcornea, embryo minimus in albumine curvato inclusus.

Clavis quadrifida L. fl. pl. 2. II. p. 139. Herb. fl. nov. p. 292. Querc. fl. ex. p. 99. Alib. fl. judr.

P. foliis subpinnatis subrotundis simplicibus nervosis, perianthio stylo lobato - bidentato, internodiis linearibus subulatis.

Icon. C.F. Nees gen. pl. pers. II (Paris)

Len. Enc. bot. pl. 319

Fl. dan. 4. 139

Engl. bot. 4. 7.





Asparagus L.

Gen. pl. 2. II. p. 120

Ch. ~~Asp. hemisphaerica~~ and ~~Asp. bari~~ in ~~tabula~~ ~~angustifolium~~ ~~abrupte~~ ~~constrictum~~, ~~laevius~~ ~~in~~ ~~superioribus~~ ~~lobis~~.  
var. ~~agris~~ ~~patens~~. ~~Stem~~ ~~non~~ ~~laevius~~ ~~peristichia~~ ~~infata~~, ~~floribus~~ ~~apertis~~, ~~et~~ ~~longis~~  
p. ~~Asp. hemisphaerica~~ ~~impositis~~ ~~puberulis~~, ~~1/4~~ ~~in~~ ~~unum~~ ~~conatus~~, ~~ovis~~ ~~retortum~~, ~~lobis~~ ~~laevibus~~.  
lobis - ~~Basia~~ ~~suberis~~ ~~hirsutis~~ ~~hexagonis~~, ~~ramibus~~ ~~globosis~~ ~~angulatis~~. - ~~Class. bot.~~  
• ~~radix~~ ~~bot.~~ ~~monog.~~ ~~lyth.~~ ~~lat.~~ -

2. ~~Asparagus tenuifolius~~ ~~Lin.~~ ~~enc.~~ ~~bot.~~ ~~tab.~~ ~~1.~~ ~~Asp. tenuifolius~~  
radix ~~Boiss.~~ ~~prosp.~~ ~~pl.~~ ~~forq.~~ ~~p.~~ ~~22~~

A. caule ~~laevibus~~ ~~lobis~~ ~~ovatis~~ ~~laevibus~~, ~~filigibus~~ ~~peristichis~~ ~~ramibus~~ ~~laevibus~~, ~~lobis~~ ~~peristichis~~ ~~in~~ ~~superioribus~~  
~~longis~~, ~~floribus~~ ~~antheris~~ ~~oblongis~~ ~~interaxillaribus~~

Syn. A. ~~maritimus~~ ~~Veitch.~~ ~~fl.~~ ~~exc.~~ ~~p.~~ ~~118~~

2 A. ~~amarus~~ ~~DC.~~ ~~cat.~~ ~~h.~~ ~~monog.~~ ~~p.~~ ~~81~~ ~~Herb.~~ ~~fl.~~ ~~austr.~~ ~~1.~~ ~~p.~~ ~~439~~

2 A. ~~maritimus~~ ~~Pitt.~~ ~~bot.~~ ~~ways.~~ ~~n.~~ ~~120~~

1 Hab. in ~~montibus~~ ~~prope~~ ~~Nona~~ (~~v. Sic.~~ ~~in~~ ~~herb.~~ ~~prof.~~ ~~Peter.~~) ~~Herb.~~ ~~lyon.~~ ~~major.~~ ~~Herb.~~ ~~bot.~~ ~~lat.~~ ~~2~~

A. ~~officinalis~~ ~~L.~~ ~~sp.~~ ~~pl.~~ ~~2.~~ ~~II.~~ ~~p.~~ ~~448~~ ~~Herb.~~ ~~fl.~~ ~~austr.~~ ~~1.~~ ~~p.~~ ~~438~~ ~~Veitch.~~ ~~fl.~~ ~~exc.~~ ~~p.~~ ~~118~~ ~~Alsch.~~ ~~fl.~~  
~~rad.~~ ~~p.~~ ~~82~~

A. caule ~~laevibus~~ ~~lobis~~ ~~ovatis~~, ~~filigibus~~ ~~peristichis~~ ~~ramibus~~ ~~laevibus~~, ~~lobis~~ ~~peristichis~~ ~~in~~ ~~superioribus~~  
~~longis~~, ~~floribus~~ ~~antheris~~ ~~oblongis~~ ~~interaxillaribus~~.

Don. C. 2 ~~Nay~~ ~~gen.~~ ~~pl.~~ ~~prosp.~~ ~~II~~ (~~Asparagus~~)

Engl. ~~bot.~~ ~~V.~~ ~~bot.~~ ~~359~~

Flor. ~~den.~~ ~~bot.~~ ~~305~~

1 Hab. ~~Sparg.~~ ~~pitomi.~~

~~Hab.~~ in ~~partibus~~ ~~indis~~ ~~exca~~ ~~Sagris~~, ~~et~~ ~~Narenta~~, ~~ubi~~ ~~legit~~ ~~Neumayer.~~ ~~Herb.~~ ~~bot.~~ ~~lat.~~ ~~2.~~

~~Uf.~~ ~~Radix~~ ~~in~~ ~~diambro~~ ~~quod.~~ ~~Turcom.~~ ~~et~~ ~~alios~~ ~~sunt.~~ ~~Species~~ ~~in~~ ~~Dalmatia~~ ~~vix~~ ~~culta.~~

2 A. ~~amphiplex~~ ~~L.~~ ~~sp.~~ ~~pl.~~ ~~2.~~ ~~II.~~ ~~p.~~ ~~449~~ ~~Vij.~~ ~~Herb.~~ ~~lat.~~ ~~p.~~ ~~27~~ ~~Herb.~~ ~~fl.~~ ~~austr.~~ ~~1.~~ ~~p.~~ ~~439~~ ~~Veitch.~~ ~~fl.~~  
~~exc.~~ ~~p.~~ ~~118~~ ~~Alsch.~~ ~~fl.~~ ~~rad.~~ ~~p.~~ ~~82~~ ~~Pitt.~~ ~~bot.~~ ~~ways.~~ ~~n.~~ ~~119~~

A. caule ~~puberulis~~ ~~angulatis~~ ~~ramibus~~ ~~laevibus~~, ~~lobis~~ ~~peristichis~~ ~~in~~ ~~superioribus~~ ~~longis~~, ~~floribus~~ ~~antheris~~ ~~oblongis~~ ~~interaxillaribus~~

Don. ~~Sibth.~~ ~~et~~ ~~Son.~~ ~~fl.~~ ~~prosp.~~ ~~IV.~~ ~~bot.~~ ~~353~~

Zaunich. ~~piant.~~ ~~bot.~~ ~~van.~~ ~~bot.~~ ~~179~~

Class. ~~hist.~~ ~~pl.~~ ~~V.~~ ~~p.~~ ~~118~~ ~~CLXXVII~~ ~~fig.~~ ~~imp.~~

1 Algo. ~~Sparg.~~ ~~Spargina.~~

~~Hab.~~ in ~~montibus~~, ~~et~~ ~~ad~~ ~~regem~~ ~~totius~~ ~~Dalmatiae.~~ ~~Herb.~~ ~~bot.~~ ~~lat.~~ ~~2.~~

~~Uf.~~ ~~Turcom.~~ ~~hujusmodi~~ ~~planta~~ ~~rad.~~ ~~est~~ ~~et~~ ~~per~~ ~~quod~~ ~~quod~~ ~~in~~ ~~superioribus~~ ~~montibus~~ ~~vix~~ ~~culta~~ ~~in~~ ~~partibus~~ ~~diambro~~ ~~quod.~~







3.	<i>Ovario vete</i>	4	<i>Ovario contorta</i> <i>aceras</i>
4	<i>Labell. bifid</i>	5	<i>Labell. indivis</i> via <i>frimbria</i> <i>Spiranthes</i>
5	<i>Labell. heteroclinus calloso</i> <i>Ophrys</i>		<i>Labell. bifid</i> baji <i>hirsuta</i> <i>Spiranthes</i>
6	<i>Ovario vete</i>	8	<i>Ovario contorta</i> <i>Cephalanthus</i>
7	<i>Labell. bifid</i>	8	<i>Labell. indivis</i> <i>Linadrom</i>
8	<i>Labell. baji hirsuta</i>	9	<i>Labell. calloso</i> <i>Nectis</i>
9	<i>Labell. imbricatus vete</i> <i>contorta vete</i> <i>Callorhiza</i>		<i>Labell. <del>contorta</del></i> <i>neobis geminata</i> <i>Erythraea</i>

Ch. *Pteranthus laevis* <sup>2. quatuor</sup> *subaequalibus*, <sup>extremis</sup> *provisis* - *connatis*, *intra* *orbis* *patulis*, *labell.*  
*ligulis* *sub* *postea* *callosis*. *Gynostemium* *breve*, *anthera* *admixta* *locali* *baji* *divergentibus*.  
*Pollinia* *tria* *caudiculata*, *retinente* *viscid.* *Staminodia* *dua*. *Stigma* *marginatum* *orbem*  
*reflexum*. - *Cl. Gynostemii* *Or.* *Diandra* *Syst.* *ix.* -

*Pteranthus bifidus* *Nich.* <sup>amst. p. 55</sup> *tab.* *Nich.* *fl.* *xxx.* *p.* *120 et 140*"

B. *filix* *radicalibus* *brachy* *oblongis* *caulinis* *lanceolatis* *parvis* *ovoidibus*, *laevibus* *perigonis*  
*obtusis*, *calcare* *ovario* *longiori* *descendente* *rectiuscula* *filiformi*

*Syn.* *Ophrys* *bifidus* *L.* *sp.* *pl.* *ed.* *tbl.* *p.* *1531* *Hort* *fl.* *amst.* *2. p.* *525*

*Non.* *Nich.* *fl.* *xxx.* *p.* *1142* *Hort* *fl.* *amst.* *2. p.* *525*  
*Hall.* *hist.* *herb.* *tab.* *xxxv.* *fig.* *2-6* *N.* *pl.* *herb.* *tab.* *xxxv.* *1.* *p.* *2-6*  
*Engl.* *bot.* *t.* *22*  
*Boiss.* *pl.* *herb.* *2.* *tab.* *xv.* *p.* *x.*  
*Carmon.* *epit.* *p.* *625* *fig.* *ext.*

3. *clavata*: *calcare* *horizontali* *aristato* *apice* *clavato*

*Syn.* *Pteranthus bifidus* *Alloch.* *fl.* *xxx.* *p.* *212*  
*Non.* *Vaill.* *bot.* *paris.* *tab.* *xxx* *p.* *7.*

*Hab.* *in* *montibus* *sylvarum* *sicula* *Perago* *prope* *Castro*, *et* *in* *partibus* *circa* *Xerax* *prope* *Zan.*  
*Hort* *Majo.* *Herb.* *albid.* *propr.* *2.*

*Obs.* *Varietas* *prope* *Charitum* *de* *quibus* *species* *specie* *videtur* *Dussin*: *dentata* *homon* *forma*  
*intra* *orbis* *calcare* *filiformi* *et* *ovario* *brachyorbite*, *quibus* *exhibetur* *in* *icon.* *235* *flor.* *Dussin*  
*calcare* *vete* *et* *ovario* *clavato* *ut* *in* *icon.* *Halleri* *l.c.* *tab.* *35* *p.* *4.*, *et* *denique* *calcare* *deflexo*  
*et* *ovario* *aristato* *et* *subulato* *ut* *in* *Carmon.* *epit.* *p.* *625* *fig.* *ext.* *Hinc* *calcare* *filiformi*







*O. piper* Jacq. fl. austr. IV. tab. 307 Herb. fl. austr. 2. p. 500 (Herb. fl. austr. p. 125)  
Alch. fl. ind. p. 211 Herb. bot. vigo. n. 654

*O. lutea* <sup>var. alba - geminata</sup> ~~var. alba - geminata~~ <sup>profunda</sup> lobis 2-partitis, lobis oblongo-linearibus, vix late  
obovata dimidia lobis mucronatis, calice sublobato ovato dente breviter, perianthio breviter  
antheris connatis, spica thyrsoidea, brachiis breviter, filij ovali-oblongis.

*O. juncea* Jacq. bot. vigo. tab. 307 p. 1334  
Herb. bot. vigo. tab. 307

Hall. bot. vigo. tab. 307

Vahl. bot. vigo. tab. 31. f. 27-28

Jacq. fl. austr. II. tab. XV p. 2

Herb. ad marginem agrorum, et in agris, circa Lam., Sibiria, Sibiria, Sibiria, Sibiria, Sibiria.

Fl. legit, Major. Sibiria Sibiria Sibiria Sibiria Sibiria, lobis alba vix geminata. 2

*O. virgata* ~~Herb. fl. austr. 2. p. 529~~ <sup>Herb. fl. austr. 2. p. 529</sup> (Herb. fl. austr. p. 124)  
Alch. fl. ind. p. 211

*O. lutea* <sup>var. alba - geminata</sup> ~~var. alba - geminata~~ <sup>profunda</sup> lobis 2-partitis, lobis oblongo-linearibus, vix late  
obovata dimidia lobis mucronatis, calice sublobato ovato dente breviter, perianthio breviter  
antheris connatis, spica thyrsoidea, brachiis breviter, filij ovali-oblongis.

*O. juncea* Jacq. bot. vigo. tab. 307 p. 1334  
Herb. bot. vigo. tab. 307

Hall. bot. vigo. tab. 307

Hall. in Suppl. fl. jen. tab. VI f. ext.

Vahl. bot. vigo. tab. 31. f. 22, 23, 24

Herb. in parvis ad Doccynge et Cyrenae prope Lam., in mont. Sibiria supra Sibiria, circa  
Sibiria, Sibiria prope Lam., et Sibiria. Flores albae-linearibus, lobis  
geminatae-geminatae. Herb. Major. 2.

*O. lutea* <sup>var. alba - geminata</sup> ~~var. alba - geminata~~ <sup>profunda</sup> lobis 2-partitis, lobis oblongo-linearibus, vix late  
obovata dimidia lobis mucronatis, calice sublobato ovato dente breviter, perianthio breviter  
antheris connatis, spica thyrsoidea, brachiis breviter, filij ovali-oblongis.

*O. lutea* Lam. fl. fr. ed. I (1778) II. p. 507 Alch. fl. ind. p. 211 Herb. bot. vigo. n. 654  
(Herb. synon.)

*O. lutea* <sup>var. alba - geminata</sup> ~~var. alba - geminata~~ <sup>profunda</sup> lobis 2-partitis, lobis oblongo-linearibus, vix late  
obovata dimidia lobis mucronatis, calice sublobato ovato dente breviter, perianthio breviter  
antheris connatis, spica thyrsoidea, brachiis breviter, filij ovali-oblongis.









Hb. in agraio prope Bragança, ubi legit prof. Alchizyo. Flor. inq. vrb. muniti. H. Maja 7

*Acens* R. Br.

Hort. kew. 2. III. h. v. p. 191

Ch. Liriodendron laevig. quinij communitatij, internodij, with angustioribus, lobis glande  
 linearibus trij gibberis. Gynostemium brev. anthera, anthera distincta, Cauda dymij  
 trij oppositij. Pelticium linea caudiculis, rotundis in unum connatis buspand  
 inlongij. Stamina obrot. Stylus coniformis gynostemii et lobis majoribus p. Cl.  
 Gynandri Ord. Standri Syst. ex. -

*Acens* antheropora R. Br. l. c. Neubl. fl. exc. p. 127

A. ~~lobis~~ <sup>lobis</sup> ~~trij~~ <sup>trij</sup> ~~oppositis~~ <sup>oppositis</sup> ~~gynostemio~~ <sup>gynostemio</sup>, multij longitudinalij prope h. p. re laevij filiformij

Syn. Ophrys antheropora L. p. pl. 2. II. p. 1343

*Clorofium* antheropora Rich. de Orch. eur. anst. p. 32

*Hieracium* antheropora Sfr. Syst. veg. III p. 674

C. P. Nees gen. pl. fac. v. n. 18.

Hell. hist. nat. vol. n. 1204

3 Vaill. bot. par. hb. 31. p. 19. 20

1 Engl. bot. t. 29.

5 Column. cephr. 1. p. 320 f. ext.

Hb. in pinet. insula Cragus prope *Schima*, et in saxij insulae *Lesina*. Flor. in dolo, magis purpur.

Obs. Non *Hemidion* sub n. 1111 ab auctoribus huc ablat. videtur gestum *Orchidanthelium*.

*Spiranthes* Rich.

Orch. eur. in Munda Mus. IV. p. 50

Ch. Liriodendron nigra laevij communitatij, brev. castrum glande, lobis acis d'favo subpita  
 bixto, basi unguiculato calcaro. Gynostemium brev. anthera scripta, laevij ventij. Pelticium  
 linea bifaria, demum parvitas apice appret. Stamina unca. Stylus subrotundus, rotunda  
 lamina bip. parvula linear. instruct. - Cl. Gynandri Ord. Standri Syst. ex. -

*Spiranthes* <sup>acuminatij</sup> ~~acuminatij~~ Rich. l. c. Neubl. fl. exc. p. 127 Alch. fl. ind. p. 214

S. ~~caud. v. p. 1343~~  
 filij <sup>altern</sup> adrotatij orb. Majorij in pectus contractij, castris adprij apiceformidij, p. p.  
 spirali, lobis <sup>altern</sup> ~~obovatis~~.

Syn. Ophrys spiralis L. p. pl. 2. II. p. 1340 (exl. m. v.)

*Nectria* acuminatij H. fl. eur. 2. p. 543

*Hemidion* ~~caud. v. p. 1343~~

Engl. bot. t. 541

H. d. m. t. 387

Hb. in ~~pinet. insula~~ <sup>pinet. insula</sup> ~~Cragus~~ <sup>Cragus</sup> prope *Lava*, ubi legit prof. Alchizyo. Flor. in dolo, lobis in dolo,

fragrantij. Flor. Aug. Sept. 7.







*Flammula sulphurea flavicans* \*  
O. ~~villosa~~ ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

O. lobis obtusis, lobis medio fere bari integro, apice trilobis, lobo medio minime appendice; fovea fere <sup>obovata</sup> recta, laciniis perianthii interioribus lanceolatis-linearibus, marginibus puberulis, gynostemio <sup>flavio</sup> ~~recto~~ ~~angulato~~

Herb. in saxoij montij Beruiffonij prope Douai, ubi legit A. Andrich. Nony perianthii lobis lobis fere ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

Desv. Tubus subglobosus. Planta bi-tripliciter graciliter biflora. Nisi usque ad ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

laciniis, basi angustis, planis, acutis-pet. Nony hinc ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

majoribus, basi lanceolatis-subpetalis, perianthii laciniis ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

vis apicibus, laciniis exterioribus ellipticis-oblongis, interioribus lanceolatis-linearibus, flavis, fovea margini ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

opposita et minime puberulis, apice obtusis, ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

marginibus ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

curvis, integris, apice emarginatis cum appendice curvata, lobis medio fere ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

trilobis, supra ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

puberulis, utrinque lobis, apice acutis exiguis, fovea recta. Gynostemio ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

laciniis perianthii interioribus angulatis.

Obi. affinis O. villosae Desf. coll. de Douai. p. 8. hb. h. Dory et Chamb. exp. simul. de Douai. bot. p. 263 ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

pl. XXXII p. 3. nec W., et magis O. villosae Fisch. icon. bot. cent. ix p. 27 f. 1172, quae ab icon. et descriptione utriusque auctoris quidam ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

fructu ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

gynostemio angulato, nec tripliciter ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

flor. duobus minoribus, lobis appendice minime ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

late subrotunda-obovata, unicolor nec ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

glauc. Desf. et Dory lobis obtusis nec subpedunculatis, hinc nec longis ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

minoribus; a glauc. ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

ultramarinae ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

O. fissa ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

O. invidios Desf. coll. de Douai. p. 6. Fisch. icon. bot. cent. ix p. 21 ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

O. lobis oblongis, lobis medio fere bari integro, apice trilobis, lobis medio minime appendice; fovea fere <sup>obovata</sup> recta, laciniis perianthii interioribus lanceolatis-linearibus, marginibus puberulis, gynostemio <sup>flavio</sup> ~~recto~~ ~~angulato~~

Icon Desf. loc. cit. hb. 3

Fisch. icon. bot. l. c. f. 1148

Herb. in demum insula Lesina, ubi legit prof. Stolzi. Flores perianthii flavo-roseis, lobis obtusis, hinc magis longis parallelis oblongis conjunctis, apicibus ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

Herb. Agri. - 2.

Obi. affinis O. villosae Lk., a qua praecipue differt lobis planis nec curvis, velutis nec ~~fructu~~ ~~sem~~ ~~bot~~ ~~cent~~ ~~ix~~ ~~p~~ ~~27~~ ~~an~~ ~~Desf~~?

O. lutea Cavan. ic. pl. 2. p. 46 Reichb. fl. ex. p. 128 et in adl. p. 140 n. 86a Alch. fl. ind. p. 213-14

O. labell about major villosa medio utriusque marginis (fimbriata, supra medium trilobis, lobis lateralibus <sup>interioribus</sup> brevibus, medio <sup>lobis</sup> obcordatis - lobis paulo brevioribus, laminae parietalis interioris oblongo-linearibus floris glabris synochromis mutis subaequalibus

Syn. O. insubrica L. sp. pl. ed. II p. 1343

Icon. Reichb. icon. bot. 18. p. 1149

Bois et Chamb. exp. univers. de Mor. bot. pl. XXXII f. 2.

Cavan. ic. pl. 2. pl. 160

Breyer. <sup>ext. fl. cand.</sup> bot. 45 p. ext.

Mons. hist. pl. 3. 5. 12 t. 60 p. 15

ubi invenit post Alchinger.

Hab. papata in parvis umbrosis circa Zava non longe a Johanne de la Botziana. Flor. junio hinc laminae exterioris indivisa, interioris lobis, labellae fimbriae, disco sanguineo, basi mucilaginea lobis oblongis ciliatis utra. Fl. Agr. Mayo. 4.

### Serapias L.

gen. pl. 2. 57. p. 332 (excl. spec.)

Ch. Pseudanthus laminae exterioris cucullatae - connatis, interioris separatis a basi. Dilata longe cuspidata, lobis acuminatis basi bilobata trilobata, lobis lateralibus brevibus medio majori reflexo. Synochromium oblongum arcuatum, apice in filamentum lamellatum productum: antheris adnatis lobis medio distinctis. Pteris binae, caudicibus subrotundis unice basivulae inclusae connatis. Straminei oblongi. Stylus oblongus curvus. — Cl. Synochromium Ord. Diandria Syst. 128. — (excl. 1794)

Serapias Lingua L. sp. pl. 2. 57. p. 1364 Herb. fl. univers. 2. p. 597 Reichb. fl. ex. p. 129

Alch. fl. ind. p. 212 (excl. syn.)

S. labellae basi <sup>glabrae</sup> ~~trilobatae~~ lobis medio ovatis - lamellatis <sup>genuis</sup> ~~floribus~~ <sup>floribus</sup> ~~parietalis~~ <sup>interioribus</sup> ~~lobis~~ <sup>lobis</sup> ~~oblongis~~ <sup>oblongis</sup> ~~linearibus~~ <sup>linearibus</sup> ~~floris~~ <sup>floris</sup> ~~glabris~~ <sup>glabris</sup> ~~synochromis~~ <sup>synochromis</sup> ~~mutis~~ <sup>mutis</sup> ~~subaequalibus~~ <sup>subaequalibus</sup>

Icon. Pich. oper. M. I. tab. 128 p. 2. Column. explor. i. tab. 321

Mons. hist. pl. 3. 5. 12 t. 14 p. 21

Lam. ill. bot. par. ven. t. 42 p. 11

Hab. in partibus herbis junceis inter cetera, ad Xerxes prope Zava. non longe a Cethon, Beger. ubi invenit <sup>in valle</sup> ~~post~~ <sup>post</sup> ~~Alchinger.~~ <sup>Alchinger.</sup>

H. Mayo, Junii 4.

Pl. Labellae longioribus medio longioribus unice, plus minusve acutis, indivisim lobis, laminae parietalis interioris cucullatae - connatis, interioris separatis a basi. Dilata longe cuspidata, lobis acuminatis basi bilobata trilobata, lobis lateralibus brevibus medio majori reflexo. Synochromium oblongum arcuatum, apice in filamentum lamellatum productum: antheris adnatis lobis medio distinctis. Pteris binae, caudicibus subrotundis unice basivulae inclusae connatis. Stylus oblongus curvus. — Cl. Synochromium Ord. Diandria Syst. 128. — (excl. 1794)

S. cordigena L. sp. pl. 2. 57. p. 1365 Reichb. fl. ex. p. 130 Alch. fl. ind. p. 212

S. labellae basi <sup>glabrae</sup> ~~trilobatae~~ lobis medio cordatis - ovatis acuminatis, floris <sup>interioribus</sup> ~~parietalis~~ <sup>interioribus</sup> ~~lobis~~ <sup>lobis</sup> ~~oblongis~~ <sup>oblongis</sup> ~~linearibus~~ <sup>linearibus</sup> ~~floris~~ <sup>floris</sup> ~~glabris~~ <sup>glabris</sup> ~~synochromis~~ <sup>synochromis</sup> ~~mutis~~ <sup>mutis</sup> ~~subaequalibus~~ <sup>subaequalibus</sup>

Syn. Helleborinae cordigena Schaff. et Maur. fl. rom. prod. p. 312



*Finn.* hexag. t. 5 <sup>iv</sup>f. ext  
Herb. in umbrosa sylvarum <sup>montif. Uolub. alt. in vally</sup> nigrae, ubi legit prof. Albrechtus Koenig, <sup>rubat.</sup> labell<sup>is</sup> <sup>ovis</sup>  
flora maculata. P. legit 7.

C. rubra Michx. l. c. Peitb. fl. exc. p. 132

C. labella ~~arven~~ exsult. linearijque primislibi omnibus cumintis, brevij florisq. ovis  
pubescent. subglobulij, plij lanceolatis.

Syn. longij rubra L. syst. veg. 2. XII p. 699 Engl. bot. 7. Flor. den. 6, 8  
Epigynij rubra Sw. act. botan. 1800 p. 232 Bot. fl. aust. 2. p. 545

Non. Engl. bot. t. 437

Pl. den. t. 348

Hall. hisp. strap. bala. n. 1299

*Finn.* iv. hexag. t. 6. p. intern.

Herb. in sylvarum montij Divon, ubi legit D. d. Clementi. <sup>Flora purgum</sup> ~~Fl. den.~~ <sup>labell</sup>  
alba, <sup>linis</sup> ~~rubr~~ <sup>pubescent</sup> ~~pubescent~~ <sup>longitudinalit</sup> ~~longitudinalit~~ <sup>undulatis</sup> ~~undulatis~~ <sup>floridij</sup> ~~floridij~~ <sup>primislibi</sup> ~~primislibi~~ <sup>vista</sup> ~~vista. P. Non. Bot. 7.~~

*Lindorum* Down.

inst. v. heb. 2. p. 437

Ch. *Penicathium* laevij omibz erect. pubulij, internij melle congestiorij, labella  
transverse quadrata, antial. bractei unguiculata calante, terminali indivij  
gynostemium marginibz angulata. Spermium elongatum erectum,  
anthera orbicly deors appax levij approssimatij. Pollen granulum  
procella *Desmon* appubinata. Staminidia nulla. Stigma obcordatum  
rotunda brevissim procella instruct. Cf. *Spermidia* W. *Diandria* 194. 4.

*Lindorum* abotivum Sw. vov. act. botan. 6. p. 80 Bot. fl. aust. 2. p. 545 <sup>fidit</sup>  
fl. exc. p. 131.

L. caule rignos asphyle, labella ovis undulata, aliam rubella ovis subangant.

Syn. Orhicly abotivum L. p. fl. 2. t. 1. p. 1336

Non. C. F. Nees gen. pl. fasc. v. hb. 11

Say. fl. aust. hb. 193

Hall. hisp. strap. bala. n. 1288

Herb. in dumosij collium circa ~~Caten~~ <sup>Caten</sup>, et ~~Nejderrich~~. <sup>Post plant isolau</sup>, <sup>floridij</sup>  
Nestler ~~florid~~. L. <sup>Fl. maji 7.</sup>

~~herb. den.~~  
act. vov. 1840 p. 33 (exl. spicib.)

Ch. *Penicathium* laevij quinij cumintibus subangulatis, labella pendula trii saepe  
cauliculis subrotunda, lob. medio productij, apice bifido. Gynostemium brev erectum,  
anthera orbicly trii appax levij medio *Dij* <sup>vista</sup>. Pollinaria tria subglobulata

rad., rhizomulo communi juncta. Stylus transversus marginatum, reticulatum laminae indivis.  
antheris, parvis melle. — Cl. Synandri Ord. Dindii Syst. sex. —

- Necker Nidus aq. Nich. annot. de orchid. p. 37 Quibb. fl. exc. p. 132  
N. caule vixente ~~aperta~~ <sup>valde</sup> ~~obscure~~ <sup>pubescenti</sup> perianthii laciniis oblongis longior.  
Syn. Ophrys Nidus aq. L. sp. pl. ed. II. p. 1339 Fl. den. fasc. IV. p. 3 x. 1.  
Lichen Nidus aq. Hook fl. lond. Hist fl. austr. 2. p. 539  
Neothidium Nidus aq. Schlechtfl fl. bot. p. LXV. C. F. Nees gen. pl. fasc. V.

Icon C. F. Nees l. c. tab. 15

Fl. den. tab. 181

Engl. bot. t. 28  
Hall. hist. stirp. helv. v. 1290  
Nees - ~~hisp.~~ <sup>stirp.</sup> v. 1290. h. 7 f. ext.  
~~tab. 181~~

Hab. in sylvis montis Birkhorn supra Macarosa in vultibus arborum perennibus. Planta  
omnino alba tunc <sup>in vultibus arborum perennibus</sup>  
<sup>in vultibus arborum perennibus</sup> <sup>in vultibus arborum perennibus</sup> <sup>in vultibus arborum perennibus</sup> <sup>in vultibus arborum perennibus</sup> <sup>in vultibus arborum perennibus</sup>  
Lign. florata <sup>in vultibus arborum perennibus</sup> <sup>in vultibus arborum perennibus</sup> <sup>in vultibus arborum perennibus</sup> <sup>in vultibus arborum perennibus</sup> <sup>in vultibus arborum perennibus</sup>

### Corallorhiza Hall.

hisp. stirp. helv. 2. p. 159

Ch. Perianthium laciniis quinque connatis, subaequalibus, lobis peltatis bracteis  
calicis lobis, lobis medio maximo linearibus, bisalpis. Gynostemium oblongum, antheris  
lobis tria oppositas transversis subparvis lobis longioribus. Pellicula vixente  
prostrata <sup>globosa</sup> <sup>globosa</sup> <sup>globosa</sup> <sup>globosa</sup> <sup>globosa</sup>. Stylus horizontalis, vixente lobis, parvis marginibus  
antheris. — Cl. Synandri Ord. Dindii Syst. sex. —

Corallorhiza imita N. Bv. in herb. den. p. 209. Quibb. fl. exc. p. 131

C. verna paniculata, perianthii laciniis acutis, ipsius deflexis, lobis oblongis oblongis.

Syn. Ophrys Corallorhiza L. sp. pl. ed. II. p. 1339 Flor. den. fasc. VIII p. 6

Corallorhiza dentata Hook fl. austr. 2. p. 547

Icon C. F. Nees gen. pl. fasc. X tab. 12

Flor. den. tab. 457

Hall. hist. stirp. helv. v. 1301.

Char. h. v. fl. p. CXX f. int. <sup>et in vultibus arborum perennibus</sup> <sup>et in vultibus arborum perennibus</sup>

Hab. in sylvis vallis montis Birkhorn cum procedente. Lign. imita Nees x.

### Epigastri Hall.

enum. stirp. helv. 1. p. 237

Ch. Perianthium laciniis quinque peltatis subaequalibus, lobis calicatis transversis  
geniculatis, articulis bracteis connatis vixente, terminalibus bifidis bilobatis. Gynostemium  
oblongum, antheris tria oppositas lobis lobis longioribus transversis  
conjugis. Pellicula vixente demum prostrata globosa appressata.  
Stylus horizontalis. Stylus subparvis lobis parvis terminatis. — Cl. Synandri  
Ord. Dindii Syst. sex. —

Egipcijski kaptivi Allim. fl. jad. 2. p. <sup>152</sup> ~~288~~ Hort. fl. austr. 2. p. 543. Nisib. fl. ex.

p. 133 Alch. fl. jad. p. 214 <sup>inter</sup> litt. bot. voy. v. 360 <sup>bractej infimij flon ~~longioribus~~</sup>

E. labella ovato-acuminata laciniis perianthii vlyzqj sublongioribus, flaj ovatis, infimij <sup>longioribus</sup>

Syn. Sengy Hallebrinae a. kaptivi L. sp. pl. 2. p. 1544  
d. kaptivi Sm. engl. bot. 4. p. 269

Icon. Sm. engl. bot. t. 269  
Holl. bot. voy. tab. n. 1295  
Niv. inv. hexag. tab. 2. opt!

Hab. in sylvis montium Vellebth, Dinara, et Veliki-Srad supra Clissa. Flory virdy  
labella videntur. H. Nivis, Julis. & <sup>ex habita et floribus et herbis ~~patet~~</sup>

Obs. Icon Camusii Egip. p. 889, ubi a Linnaeo ipso huc citatur, <sup>potest</sup> sequente extra  
bot. <sup>(1769)</sup>

E. palestina Crantz austr. 6. p. 262 Hort. fl. austr. 2. p. 544 Nisib. fl. ex. p. 154  
Alch. fl. jad. p. 214 <sup>bractej ~~infimij~~ <sup>ovatis</sup> per bractej</sup>

E. labella subrotunda, lobis laciniis perianthii angustis, flaj ovatis lanceolatis.

Syn. Sengy kaptivi V. ~~fl. austr. 6. p. 262~~  
d. palestina Seng. fl. austr. 2. p. 204 Engl. bot. engl. bot. 4. p. 280  
Hallebrina kaptivi H. du. pro. v. p. 6

Icon. C. 2. New gen. pl. fasc. X. tab.  
Engl. bot. t. 280  
Flor. du. t. 267  
Crantz l. c. t. 1. f. 5  
Holl. bot. voy. tab. n. 1296

Hab. in palestina <sup>prop. ~~bot.~~ ad</sup> Staber supra Spalat, et Ormba supra Regium. Flory rubra. Hort  
Nivis, Julis. &

E. microphylla <sup>Tabula ~~bot.~~ t. 63 ~~(1769)~~</sup> Hort. fl. austr. 2. p. 543 Nisib. fl. ex.  
p. 133

E. labella orb. acuta cuneata perianthii laciniis subaequantibus, bractej ovatis flon  
bractej, flaj lanceolatis.

Icon Waldst. et Kit. fl. rar. hung. tab. 270

Hab. in sylvis ad pedem montis Dinara supra Glavice, ubi legi n. Julis flo.  
virescentibus. Thun & Thun Spessin, sed sine huius indicatione, misit Neum.  
<sup>specimen</sup>  
yca. Flory flon virdy, maculenta. &











2. *Posidonia*

3. *Verby* ~~gründlich~~ - 2 plants <sup>gründlich</sup> ~~gründlich~~ <sup>gründlich</sup> ~~gründlich~~

Koen. anal. of bot. p. 95 2 plants <sup>gründlich</sup> ~~gründlich~~ <sup>gründlich</sup> ~~gründlich~~

Ch. Flory hermaphrodit. <sup>ovary</sup> ~~ovary~~ <sup>3-6 flous</sup> ~~3-6 flous~~ <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow <sup>flowers</sup> ~~flowers <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.  
Singulari spatula binli <sup>ovary</sup> ~~ovary~~ <sup>apert.</sup> ~~apert. <sup>antherae</sup> ~~antherae <sup>3-6 flous</sup> ~~3-6 flous <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow~~ <sup>flowers</sup> ~~flowers~~ <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.~~~~~~~~~~~~~~~~~~~~~~

*Posidonia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

Syn. *Posidonia* <sup>1791-1800</sup> ~~1791-1800 <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~~~

*Posidonia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

Non. *Posidonia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Posidonia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Posidonia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

Herba <sup>ovary</sup> ~~ovary~~ <sup>3-6 flous</sup> ~~3-6 flous~~ <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow~~ <sup>flowers</sup> ~~flowers <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.  
Singulari spatula binli <sup>ovary</sup> ~~ovary~~ <sup>apert.</sup> ~~apert. <sup>antherae</sup> ~~antherae <sup>3-6 flous</sup> ~~3-6 flous~~ <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow~~ <sup>flowers</sup> ~~flowers <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.~~~~~~~~~~~~~~~~~~~~

Herb. in man. <sup>ovary</sup> ~~ovary~~ <sup>3-6 flous</sup> ~~3-6 flous~~ <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow~~ <sup>flowers</sup> ~~flowers <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.~~~~~~~~

Herb. in man. <sup>ovary</sup> ~~ovary~~ <sup>3-6 flous</sup> ~~3-6 flous~~ <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow~~ <sup>flowers</sup> ~~flowers <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.~~~~~~~~

Herb. in man. <sup>ovary</sup> ~~ovary~~ <sup>3-6 flous</sup> ~~3-6 flous~~ <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow~~ <sup>flowers</sup> ~~flowers <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.~~~~~~~~

Herb. in man. <sup>ovary</sup> ~~ovary~~ <sup>3-6 flous</sup> ~~3-6 flous~~ <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow~~ <sup>flowers</sup> ~~flowers <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.~~~~~~~~

Ch. Flory <sup>ovary</sup> ~~ovary~~ <sup>3-6 flous</sup> ~~3-6 flous~~ <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow~~ <sup>flowers</sup> ~~flowers <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.  
Singulari spatula binli <sup>ovary</sup> ~~ovary~~ <sup>apert.</sup> ~~apert. <sup>antherae</sup> ~~antherae <sup>3-6 flous</sup> ~~3-6 flous~~ <sup>in</sup> ~~in <sup>yellow</sup> ~~yellow~~ <sup>flowers</sup> ~~flowers <sup>bract</sup> ~~bract <sup>conspic.</sup> ~~conspic.~~~~~~~~~~~~~~~~~~~~

*Zinnibellia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Zinnibellia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Zinnibellia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Zinnibellia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Zinnibellia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Zinnibellia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Zinnibellia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Zinnibellia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~

*Zinnibellia* <sup>1791-1800</sup> ~~1791-1800~~ <sup>Coen.</sup> ~~Coen. <sup>anal.</sup> ~~anal. <sup>of bot.</sup> ~~of bot. <sup>p. 95</sup> ~~p. 95 <sup>2 plants</sup> ~~2 plants <sup>gründlich</sup> ~~gründlich <sup>gründlich</sup> ~~gründlich~~~~~~~~~~~~~~





*P. uspa* L. sp. pl. ed. t. p. 185 Hort. fl. austr. 1. p. 223 Fiecht. fl. ex. p. 8  
 Alch. fl. jud. p. 35

*P.* filij omnibus submissis <sup>lineamentis</sup> membranarij pallidij <sup>lineari-  
 longis auris serrulatis undulatis, pedunculis angulatis, dugis conyviis dupo</sup>  
 oblongis apice ~~curvatis~~

Juss. C. 2. Nour. gen. pl. fasc. VI. n. 13 p. 1-8  
 Chemis. lin. cit. tab. V. f. 12  
 Engl. bot. t. 1012  
 Flor. dan. t. 927  
 Labat. icon. 1699. t. 286. f. 2.

Hab. in aquis fluminibus circa Zam, et in fluvio Kerkha, et Narente. Slav.  
 Mijo, Aug. 2.

*P. purpurescens* L. sp. pl. ed. t. p. 182, Hort. fl. austr. 1. p. 222 Fiecht. fl. ex. p. 8

*P.* filij omnibus submissis <sup>lineamentis</sup> <sup>membranarij pallidij</sup>  
 subrotatis <sup>oblongis</sup> <sup>serrulatis glanis, pedunculis angulatis, dugis conyviis dupo</sup>  
 oblongis apice <sup>curvatis</sup>

Juss. Chemis. l. c. tab. V. f. 13  
 Flor. dan. t. 196  
 Locrel. fl. pruss. t. 65.

~~Flor. dan.~~ Backh. bup. 3. f. 285 f. int. sup.

Hab. in aquis fluminibus Kerkha supra S. Arcangelis, et Narente. Slav.  
 Mijo, Aug. 2.

*P. pectinata* L. sp. pl. ed. t. p. 183 Hort. fl. austr. 1. p. 224 Fiecht. fl. ex. p. 7.

*P.* filij omnibus submissis <sup>lineamentis</sup> membranarij pallidij <sup>top. vaginatus</sup> apice  
 auris, <sup>unicurvis</sup> <sup>trihincis</sup> <sup>auris</sup>, <sup>marginis</sup> <sup>integris</sup> <sup>glanis</sup>, <sup>pedunculis</sup> <sup>angulatis</sup>  
 elongatis, apice <sup>interrogis</sup>, dugis <sup>dupo</sup> <sup>curvatis</sup>.

Juss. Chemis. lin. cit. tab. IV. f. 2 <sup>part.</sup>  
 Flor. dan. t. 1246  
 Engl. bot. t. 523  
 Vaill. bot. parif. t. 52 f. 5.

Hab. in aquis fluvii Kerkha supra Sardina, et lacus palustribus supra Domus. H. P. M. Aug. 2.  
 Obs. Simelliana *P. marina*, et per Sardina dugis <sup>curvatis</sup>. ~~affinis~~ <sup>differs</sup> <sup>per</sup> <sup>se</sup>  
 juvenis. <sup>juva</sup> <sup>discipat</sup> <sup>fluctu</sup> <sup>compso</sup>, <sup>lyali</sup> <sup>hic</sup> <sup>ubi</sup> <sup>nee</sup> <sup>adnatu</sup>, <sup>filij</sup> <sup>tri-</sup> <sup>quinguentis</sup>.





L. C. Rich. Mém. du Mus. 2. p. 365

Flory hermaphroditi: in rogi: apice strobili aut umbellae, baccat. Perianthium brevissimum  
lanceolatum breviter, stylis breviter internodis pediculis. Stamina plurima polygama  
exteriora indurata striatim, antheris in trochis terminalibus longitudinalibus obliquis.  
Ovaria 6-plurima interiora unilocularia ~~separata~~ unilocularia, stylis brevibus longis, ovum  
breviter. Carpella 6-plurima in polygama styli obliqua. Stamina plurima  
exteriora placentis parvis: articulatione unguis affixa erecta, embryone recto aut  
obverso, articulo umbilico parvis bina fructu parvis.  
Hortus paucis pulchris flos, rogi: stylis articulis, carpellis, pediculis ramis: germen  
brevi, lacinia singulis integris.

Rutaceae Tourne.

ist. v. herb. p. 271.

Ch. Perianthium stylis omnibus coloratis. Stamina novem hypogynum. Ovaria sex infera  
connata, styli: obliqui. Carpella unilocularia apice lobata, seminibus flexuosis: castis.  
- Ch. Eurpedici Ord. Herpogyni Syst. sex -

Rutaceae umbellatis L. 1. p. 41. 2. D. p. 532 Hort. fl. austr. 1. p. 488 Perid. fl. ex.

p. 99

D. stylis brevibus elongatis, carniculis: breviter: autis.

Don C. F. Mém. p. fl. p. 61. p. 20.

Engl. bot. t. 654.

Flor. dan. t. 604

Lam. enc. bot. fl. 324

Hab. in agrestibus pulchris circa Lodiis rivis, in insula Pago, et circa Navona. Hortus  
Linnæi, Indis. Flory vici. 7

Ordo XX Hydrocharitaceae

~~Ch. Braun p. fl. vici herb. p. 544~~

DC. fl. fr. ut III. p. 265

Flory hermaphroditi: in ~~rogi: apice strobili~~ <sup>umbellae</sup>. Perianthium brevissimum  
uni-bilobum placentis parvis, articulatione unguis affixa erecta, embryone recto aut  
obverso, articulo umbilico parvis bina fructu parvis. ~~Stamina plurima polygama~~  
exteriora indurata striatim, antheris in trochis terminalibus longitudinalibus obliquis.  
Ovaria 6-plurima interiora unilocularia ~~separata~~ unilocularia, stylis brevibus longis, ovum  
breviter. Carpella 6-plurima in polygama styli obliqua. Stamina plurima  
exteriora placentis parvis: articulatione unguis affixa erecta, embryone recto aut  
obverso, articulo umbilico parvis bina fructu parvis.  
Hortus paucis pulchris flos, rogi: stylis articulis, carpellis, pediculis ramis: germen  
brevi, lacinia singulis integris.

Hab. aegyptiaca et plerumque jamaica, bracteae vel longae et aedae carboventis, filij glomerati  
articulibus compositis, quibuslibet integerrimis, nervis.

*Hydrachne* L.

gen. pl. 2. II. p. 374

in scopis <sup>et aliis</sup>  
juba <sup>fructus</sup>  
juba <sup>fructus</sup>

Ch. Flor. <sup>maior</sup> *Hydrachne* in scopis aegyptiaca <sup>maior</sup> *Hydrachne*  
perianthio laevibus adhaerentibus articulis, interioribus articulis. Nervis, <sup>maior</sup> *Hydrachne*  
Jacobson, filamentis longis connatis, quibuslibet articulis, quibuslibet articulis: perianthio <sup>maior</sup> *Hydrachne*  
articulis univariis, striatis in articulis, articulis articulis, stylis brevibus, <sup>maior</sup> *Hydrachne*  
Sensu <sup>maior</sup> *Hydrachne*. <sup>maior</sup> *Hydrachne* articulis, nervis glomeratis perianthio. — Cl. *Hydrachne*  
an. *Hydrachne* Syst. 100.

*Hydrachne* Moivre *Flora* L. p. pl. 2. II. p. 1466. *Flor. pl. aegypti.* 2. p. 667. *Flora*

pl. ex. p. 14

*Hydrachne* <sup>maior</sup> *Hydrachne* articulis, hic articulis. <sup>maior</sup> *Hydrachne* nervis, <sup>maior</sup> *Hydrachne* floribus <sup>maior</sup> *Hydrachne*  
articulis pedunculatis.

Icon C. 7. Nees gen. pl. VI. n. 15

Flor. dan. t. 878

Engl. bot. t. 808

Sodomy <sup>maior</sup> *Hydrachne* p. 585 <sup>maior</sup> *Hydrachne*

Hab. in aegypti <sup>maior</sup> *Hydrachne*. *Neurospora*. *Flora* <sup>maior</sup> *Hydrachne*, *Flora* <sup>maior</sup> *Hydrachne*. *Flora* <sup>maior</sup> *Hydrachne* *Flora*  
articulis. 24.

















Up. ex hujus alioque catenis variis. *Purbotaria communis*, que in multis arboribus  
 in illis venteribus multo frequentius, in arboribus et subter landis. *Demetrium*  
*junior (Chusonia 187)*  
*Junonitium communis*, ex agris deinde in multis venteribus venteribus arboribus  
*Purbotaria* quibus venteribus ad usum venteribus communis.

P. ~~Purbotaria~~ <sup>(1872)</sup>  
*Murphy* leg. fl. cara. ad. st. tom. 2. p. 242 ~~her~~ *Heidl.* fl. exc. p. 159

P. *degruyi*, folij geminis videri admodum, *Purbotaria communis* enim aut *inclinata* venteribus  
 oblongis foliis subaequalibus, squarum foliis <sup>in opposita</sup> linearibus acutis inveni prominentibus  
 seminibus albis <sup>in</sup> oblongis, *Purbotaria communis* apice rotundata deinde brevior.

Syn. P. *Purbotaria* Haenk. bot. auf ver. nach d. viefen. p. 68 (1791) *Herb. fl. aust.* 2. p. 629  
*Walst. et Kit. pl. rar. hung.* 1. p. 160

Icon *Walst. et Kit.* 1. c. tab. 149  
*Ches.* hist. pl. 1. p. 72 f. int.  
*Deluc.* hist. leg. d. 1. p. 47 f. int.

Myr. *Purbotaria*  
*caucasiensis* *Delebith* et  
*Hab. in* ~~in~~ *montibus* *Dinarum* *in* *Albania*. *H. Nagi.*  $\frac{1}{2}$

Obs. *Purbotaria* alae seminibus venteribus lobis exhibitum in eorum pl. hung. lit. h. *Icon*  
 et *Deluc.* *hist. leg. d. 1. p. 47 f. int.*  
 autem *Walst. et Kit.* 1. c. tab. 149. p. 113. ab auctoribus hinc *Albania* venteribus  
 ad *Purbotaria*, habet *Dinarum* *Albania*, *in* *montibus* *Dinarum*, *in* *Albania* *Purbotaria*  
~~in~~ *Albania*.

sect. II *Albia*

*Albia communis* *Albia*, *in* *montibus* *Dinarum* *in* *Albania*. *H. Nagi.*  $\frac{1}{2}$   
*Albia communis* *Albia*, *in* *montibus* *Dinarum* *in* *Albania*. *H. Nagi.*  $\frac{1}{2}$

P. *Albia* *Delebith* *herb.* 2. p. 45 *Vis.* *herb. d. 1. p. 43* *Abel.* *fl. jadr.* p. 227 <sup>ex herb.</sup> *ex herb.*  
*Waga.* et *Syn.* *P. H.* *bot. venteribus* n. 701 (excl. *Syn.*)

P. folij *pectinatis* *linearibus*, *oblongis* *emarginatis*, *subtus* *albo* *bluente*, *stipulis* *ovatis* *sessilibus*,  
*foliis* *multis* *longioribus*, *squarum* *obovatis* *oblongis* *obtusis* *adpressis*, *bracteulis* *agris* *acutis* *venteribus*,  
*seminibus* *albis* *in* *oblongis* *deinde* *longioribus* *et* *lobatis*.

Syn. P. *Purbotaria* *L.* *pl. rar. hung.* 1. p. 1620 *Herb. fl. aust.* 2. p. 630  
*Albia* *Purbotaria* *Heidl.* *fl. exc.* p. 159 et p. 824  
*Al* *geobot.* *fl. rar. hung.* 1. p. 228 *Nich.* *de* *conif.* *et* *cyced.* p. 166

Icon. C. F. *Nees* *gen. pl. rar.* 1. n. 3.  
*Nich.* *loc. cit.* *tab.* XVI *f. 2.*  
*Comer.* *opit. pl.* p. 48 et 49  
*Muhlb.* *D. venteribus* *ad.* *Nagy.* 1868 p. 119

Myr. *Albia*, *Capin*  
*Hab. in* *venteribus* *montibus* *Albania* *et* *Dinarum*. *H. Nagi.*  $\frac{1}{2}$











Syr. J. breviflor, Herb. fl. austr. 2: p. 670

Non. C. F. Nees gen. pl. p. I. n. 9

Nich. de consif. et organ. bot. 2.

Engl. bot. t. 246

Dub. hist. ny. abstr. II. fol. 86 (ex Math. hist.)

Math. hist. ny. 2. Vol. 1588 p. 1156.

Pos. Karyl.

Herb. in syst. bot. Pakkenia, et Monte Sants in montibus Valle d'Alta. Hinc must. ego.

Præterea late ubi. H.

Uf. Plant. omnig., id fl. præcipue in hinc ubi dicitur: gaudens, hinc in montibus  
floriferis præcipue pectore form. pectore ubi hinc exhibetur.

Ord. XXV. Amentaceae\*

(Bivalvæ et cupuliformes Nich.)

Flora omnium amentaceae, squam. suffultis, perianthis nullis aut 3-6-fidis: max. stramineis  
4-2a perianthis aut squam. insertis, atheris uni-bilocularibus longioribus debilibus:  
perianthi ex ovario bivalvatis, loculis uni-bilocularibus, signatis ~~dentibus~~ hinc filiformibus unitis:  
by ut styl. columnari suffultis. Fructus amentacei indurati, vel cum squamis in  
stabilis speciem digni, vel in involucri cupuliformi aut capsuliformi uniti, abster. uniti:  
calycis, et plurimum monospermi, perianthibus exalbuminosis gaudens, radice vestrum fructus  
speculate

Atheris ut fructus monospermi, filij impleribus plurimorum atheris dentibus lobatis, hinc lobis  
deciduis.

Clavis analytica generum

- |   |   |
|---|---|
| Fructibus spicis — ord. N. 1.   | Fructibus in strobilum racemum aggregatis — ord. N. 4         |
| 1. Involucrum fructus distincte prægæ — 2   | Involucrum indivisum aut plerumque — 3                        |
| 2. Hærem amentis rugulosis  | Hærem amentis <del>filiformibus</del> <sup>filiformibus</sup> |
| 3. Glande in <sup>pagis</sup> <del>cupula</del> <sup>bracteis</sup> <del>cupula</del> by: immixta | <u>Castanea</u>   |
| 4. Strobili compacti imbricati — — 5  | Necesse involucri subulsi inclusi                             |
| 5. Strobili squamis membranaceis  | <u>Corylus</u>  |
| <u>Ostrya</u>   | <u>Quercus</u>  |
|   | <u>Alnus</u>  |
|   | <u>Strobili squamis lignosis</u>                              |
|   | <u>Alnus</u>  |

Ord. I. Cupuliformes

Styl. columnari, ovario perianthis dices, uniti locularem angulo centrali apertis, fructu in fructu  
involucro cupuliformi aut capsuliformi sessile.

Dagay Doum.

inst. v. herb. 1. p. 584 pl. 351

Ch. Hong in amey subflogij pumulety: mung pumulethi angumlethi s. b. fid. 1/2 quantity 8-12  
ijij loy: inserty, anthemum luntij unnetos usq. 20ijij, bap. p. ijij: pernici luntum subflogij  
dilto bifoo, pumulethi cario in luntum usqum luntum gaudet seriam tubulans p. 20  
aliter. thypope by 14ijij luntum: pumulethi aducent. Tuntij ex insulans subflogij  
Doma pumulethi unuly bing unbulans mangemay aut bap. p. ijij: insulans.  
— U. Morsini ord. Polydri sp. 100 —

Dagay syphoca L. 17. p. 2. II. p. 146. Wip. 14ijij. del. ij. 35 Herb. fl. ex. 2. p. 623  
Fiedl. fl. ex. p. 176 Alch. fl. j. d. p. 225 Pitt. bot. wyew. i. 403

F. plijit vity vity g. b. ijij, mangem subflogij subulans ijij ijij.

Don. C. F. Nay gen. pl. 1. 2. 16

Lam. enc. bot. pl. 782 p. 2.

Engl. bot. t. 1846

Jaestn. de fr. et um. 1. 1. 37. p. 2

Method. sp. 2. Uq. 1568 p.

Uly. Duk, Duk.

Herb. in silyij unnetum Velleth, Sulyij, Dikho. Pl. Capiti, Luni.  $\frac{1}{2}$

Uf. signum ad rem. un. p. 100

Castanea Doum.

inst. v. herb. 1. p. 584 pl. 352

Ch. Hong mosule uny amey filiponi loka avet formulethi, pumulethi 10ijij, pumulethi  
staminitij 8-15 ijij loy: inserty, anthemum tubulans subflogij, unnetum dilto, pumulethi  
subflogij mangemay insulans angumlethi subflogij, pumulethi insulans subflogij  
luntij 5-8 p. fid. usqum unnetum bi-10ijij luntum 6-12 arboris capiat, styl. unnetum, thypope  
luntij bi-10ijij subflogij. Tuntij ex insulans subflogij Doma pumulethi unuly 2-3  
(mangem) unbulans mangemay flous unnetum pumulethi. — U. Morsini ord. Polydri  
sp. 100. —

Don. C. F. Nay gen. pl. 1. 2. 16 (1788) Herb. fl. ex. 2. p. 623 Fiedl. fl. ex. p. 176

Alch. fl. j. d. p. 225 Pitt. bot. wyew. i. 403 (1785) Fung. Capiti L. 17. p. 2. II. p. 146.

Castanea sava Mill. Gard. dit. 2. VIII (1768) sp. 1.

Don. C. F. Nay gen. pl. 1. 2. 16  
Lam. Enc. bot. pl. 782 p. 1.  
Duk. bot. de alch. 1. 1. 50  
Method. sp. 2. Uq. 1568 p.

Allye. Koshinj. f.

Herb. forte ulmij, in insulis circa Zam, et in Niphae 16th: in sylvarum cum cortice lico et ipse

Koshinjia die ~~in insulis~~, et vasa Sklivio in Canale & Canton.

Uf. Fructus edulis, lignum ad opem vasa strickherum a phtij lignum hunc a Siaming. H. Paris f.

Quercus L.

gen. pl. 2. II. p. 354

Ch. Nony insipidi, rufi amara filiformi laxe pendula florentibus, perianthis 6-8-partitis  
alaba, stramineis 6-10 longi basi inserti, antheris haudulatis brevibus: pericarpio rugulato  
gemmiformi, brevibus et involucri unisporo in cupulae comae perianthis laxe rugo  
scapulis aut rotundis, ovum tuberculatum vasculatum, stylum brevissimum et obliquum tubu  
loso exigentem. Fructus glaucus cupulae extrae oppositae basi immixtus. - U. mucronata  
Ord. Mygdaliae L. n. 10.

1. Lign. Quercus pedunculata Willd. arb. p. 278 Fruct. fl. ex. p. 177 Arb. fl. Ind. p. 226

Quercus Robur L. sp. pl. 2. II. p. 1414 Herb. fl. ex. p. 2. p. 622 Bot. bot. v. 1. p. 119

L. folij deciduis, ramis pubescentibus, stipulis ovatis, lobis ovatis, pediculis  
apice stramineis multij, fructibus longis pedunculatis, cupulae operculi brevissimi adpressi  
Vif. Mygdaliae L. n. 10. p. 45

2. Lign. Q. pedunculata Willd. arb. p. 278 Fruct. fl. ex. p. 177 Arb. fl. Ind. p. 226

Q. perennis H. DC.  
Non. L. 2. Nony gen. pl. 1. 1. 15

H. DC. t. 1180  
Euph. bot. t. 1342  
Dich. fruct. Ind. ab. 2. t. 47

Allye. Dub. Post. et lignum Opacium

Herb. in insularibus Siaming Siamis. Hunc Meiji f.

Uf. Cortex coriis perspicuis insensit: lignum ad vasa strickherum a phtij lignum hunc a Siaming:  
cortex ipse, et folia, et galle videtur spongiosa et siccitate gaudere, hinc in  
phlegma insensibile, postquam claudicet, galeolae vasa et insipida, ubi  
hinc vasa insensibilia sub forma galeolae aut ductu, exhorum sub forma galeolae, balaui, insi  
chorij vel pyropisatij praesentem in vasa vasa. Quod quodam coffea  
modo temptat, in pulvorem hinc, obaque formae insi, in <sup>strickherum</sup> <sup>ablatum</sup> ablatum  
proprietatem. Duo vasa strickherum vasa strickherum.

Q. pubescens Willd. sp. pl. 4 p. 450 Herb. fl. ex. p. 2. p. 622 Fruct. fl. ex. p. 177

L. folij deciduis, ramis pubescentibus, stipulis ovatis, lobis ovatis, pediculis  
lobis ovatis multij, fructibus subsessilibus, cupulae operculi brevissimi adpressi.

Rem. Clus. sup. hist. hisp. p. 21 et in. pl. p. 18 f. (ab ipse fruct. et cum galle)

Herb. in insularibus Siamis cum pino. Hunc Meiji f.





<sup>Lijena.</sup>  
 Myr. Ljescnik, Bogu, Sarguzu. et pr. Ljescnik.  
 Hb. ad regu et in yskovny krtuy delovky vorkov, nec non in imply Lijem, Bogu  
 H. Herb. Martii.  $\frac{1}{2}$   
 Upr. Nunciae doly sunt.

Caupiny Douva.  
 inst. v. herb. 1. p. 582

Ch. Nary vovuti seuy amosa efidna pahle ruzp imbitu. abstrahi, pericathu  
 a ruzpam fuphu shruva 12. flum topi yente, plumenty kimpity, anthy vutbu  
 itoy substokry apic pitoy: peminu seuy amosa laxa evet dipoti budy dudy,  
 invluvi oftri phtly gomyj cozpultar bilyj petistij, pericathu cozpultar  
 limbo regu dentulata ovium bloulan bantolou vektul, style boudim, 14 p.  
 kty bilyj phtformy ovity. Nudy ex invluvi phtary vuculy khom verty  
 novozomy ad lity izvityby in vucum pndulum laxa dyffu - U. thomei v. d. h.  
 andi lept. v. -

Caupiny Dchuy L. 1 p. 2. 11 p. 146 V. kty. dl. p. 27 Hst fl. austr. 2. p.  
 626 Spich. fl. austr. p. 175 Hst. fl. austr. p. 225 ~~Spich. fl. austr.~~

C. involucri phtly kigartyj, leiny lanceoly, <sup>longior</sup> ~~intermedia~~ longior

Non C. D. N. v. fl. austr. p. 1. v. 12  
 Dub. v. fl. austr. 1. t. 49  
 Lam. Enc. bot. pl. 780  
 Gaertn. de fructum. 2. t. 89. p. 2.  
~~Meth. d. fl. austr.~~ Douva. inst. v. herb. 1. p. 348

Myr. Bide Gub.  
 Hb. viny ac regu in ruzp vorkum; legi in vort kalyz. H. Agula.  $\frac{1}{2}$

C. <sup>non 2.</sup> duryy fl. austr. 2. t. 243 Spich. fl. austr. p. 175  
 C. involucri phtly <sup>semi-cordaty</sup> izvityj cozpultar vuculy subangulyj delyj.  
 Syn. C. vityj Lam. Enc. bot. 2. t. 700 Hst fl. austr. 2. p. 626

Non. Sup. loc. ut. hb. 60 (excl. fig. int. ext)

Myr. Czernyob. Gab. et Lijem. ~~Spich. fl. austr.~~  
 Hb. vlyjstomy i velyj luyje ruzpny krtuy delovky H. Mart. lept.  $\frac{1}{2}$   
 Bti. Nigun infinis ruzpny extus viny koptimoz invluviu brypntem exhibit  
 C. Dchuy pa cozpultaryj canp vucum illu C. duryy i ruzp paxim dlym.

Ostrya Michx.

nov. gen. pl. p. 223

Ch. Flory masculi semper amara cylindrica pendula saepe imbricata abacta, perianthii & siliis simplicibus staminibus 12. glabra basi genitae filamentis brevibus, antheris unilocularibus utroque breviter, apice filiformi. femineae semper amarae oblongae erectae basi diffusae, bracteis deciduis, involucri uniloculari filiformi genitae margine callositate integro perianthium amplexuans limbo saepe indiviso ovarium bracteola bracteata obducunt, stylo bursae, et stylo hinc filiformi includuntur. Fructus ex involucri ~~receptaculo~~ in striatulis callosis uniuersis hinc longis ~~apice filiformi~~ nonnullis abactis in striatulis ~~receptaculo~~ ~~obducuntur~~ diffusis. — (C. Monroii Ord. Syll. l. p. 100.)

Ostrya carpinifolia Scop. fl. car. 2. t. 1. tom. 2. p. 243 ~~Michx. fl. car. p. 175~~ C. F. New York. p. 1. n. 13

O. striatilis Michx., filij utriusque duplices - comites, genitae oblongae.  
Syn. O. vulgaris W. sp. pl. 4. p. 469 Hort. fl. austr. 2. p. 622 Alch. fl. ind. p. 225 Bot. bot. n. 674  
Nov. C. F. New York. p. 1. n. 13  
Michx. nov. gen. pl. p. 104

Myr. Frax et lignum Fraxinam

Hab. in sylvestri <sup>colluvium</sup> circa Jann, lign. Virginis, Capens. Flor. May. Fr.  
Obs. Striatilis nunc pendula, nunc patens, nunc erecta in eadem arbor.

Sub II. Ochelinae

Stylo saepe, ovum libero, callosis dimidiatis affixis

Alnus Torr.

inst. v. herb. l. p. 587 tab. 359

Ch. Flory densa imbricata, masculi semper amarae cylindrica pendula, bracteis filiformi marginibus serratis utroque, perianthii 4-partito, staminibus 4. ejusdem longi integro apice filiformi, antheris bilocularibus breviter, femineae semper amarae oblongae, bracteis involucri genitae hinc trifido ovum bina bracteola bracteata tegunt, stylo hinc in utroque breviter filiformi, fructus striatilis & bracteis lignosis uniuersis longis lignosis nonnullis parvis. — (C. Monroii Ord. Syll. l. p. 100.)

Alnus fraxinea Gaertn. d. fr. et sem. 2. p. 54 Hort. fl. austr. 2. p. 612 Michx. fl. car. p. 175

O. filij subornatis basi uniuersis apice saepe repandis - 10000 glabris filiformi, axillis uniuersis utroque villosis, perianthii nonnullis quadrifido.

Syn. Betula alnus L. l. p. 2. t. p. 1394

Nov. C. F. New York. p. 1. n. 13

Engl. bot. t. 1508

Gaertn. d. fr. et sem. 2. t. 90. p. 2

Jub. brit. dy. astr. 1. t. 15

Michx. fl. car. l. c. p.

Myr. betula

Hab. in insulari circa ~~Nova~~ ubi hucusque Alnus invenitur. H. Mart. Cap. Fr.



*S. fruticosa* L. 1 p. p. 1643 ~~Hort flor. austr. 2 p. 634~~ Quirb. flor. 172  
<sup>172</sup>  
*S. abrotanica*, <sup>foliis</sup> ~~foliis~~ <sup>late lanceolatis acuminatis comulatis glandulosis inflexis serratis</sup>  
<sup>marginibus</sup> ~~marginibus~~ <sup>obtusis serratis, acuminatis in ramulis floris apice triangulis</sup>  
<sup>stylis</sup> ~~stylis~~ <sup>ovatis lanceolatis pedicellatis glabris, stylis ovatis in propriis axillis</sup>

Jap. *S. viciaria* Hoffm. h. p. 101. tab. 2. p. 9. Sm. engl. bot.

Non. Hort sal. tab. 14  
Hoffm. l. c. tab. 31.  
Engl. bot. t. 1807 et 1937

Hb. ad agr. Leipz. comitibus harenis circa Devins. Pl. Agri. Magi. ¶

*S. amygdalina* L. 1 p. p. 1643 Hort flor. austr. 2 p. 634

*S. abrotanica*, <sup>foliis</sup> ~~foliis~~ <sup>semicordatis, plicis lanceolatis acuminatis utraque glaucis glandulosis</sup>  
<sup>marginibus</sup> ~~marginibus~~ <sup>acuminatis in ramulis floris apice triangulis, capsulis ovatis conis</sup>  
<sup>pedicellatis glabris, stylis brevibus, nigrescentibus dimorphis conis</sup>

Syn. *S. amygdalina* p. *Xifala* Quirb. flor. austr. p. 171  
*S. hiemalis* Alsch. fl. jard. p. 8 nec L.

Non. Hort sal. t. 1. tab. 13. 14.

Hb. Leipz. ~~prop. Zan.~~ <sup>ad agr. long. Baumg.</sup> et in inst. Latina. Pl. Magi. ¶  
~~Hort flor. austr. 2 p. 634~~

*S. purpurea* L. 1 p. p. 1644 Quirb. flor. austr. p. 171

*S. fruticosa* ~~Differ.~~ <sup>caespitosa</sup> ~~foliis~~ <sup>foliis</sup> ~~obtusis lanceolatis glaucis, capsulis ovatis conis pedicellatis glabris, stylis brevibus, nigrescentibus dimorphis conis~~  
<sup>obtusis lanceolatis glaucis, capsulis ovatis conis pedicellatis glabris, stylis brevibus, nigrescentibus dimorphis conis</sup>  
<sup>obtusis lanceolatis glaucis, capsulis ovatis conis pedicellatis glabris, stylis brevibus, nigrescentibus dimorphis conis</sup>

Syn. *S. monarda* Arduin. Memo. d'agr. p. 67 Hoffm. h. p. 101. t. 1 p. 18  
*S. caprea* Pott. bot. wip. n. 502 et herb. vall. L.

Non. Hoffm. loc. cit. tab. 1. f. 1. 2. tab. 4. f. 1. tab. XXIII p. 8.  
Arduin. loc. cit. pro II.

Ulyss. Rehete.

Hb. ad agr. et in agroribus circa Devins, ad Casak d. Hög, et ad Sardon, et advia Solona.

Obs. hae ~~foliis~~ <sup>foliis</sup> ~~obtusis lanceolatis glaucis, capsulis ovatis conis pedicellatis glabris, stylis brevibus, nigrescentibus dimorphis conis~~  
<sup>obtusis lanceolatis glaucis, capsulis ovatis conis pedicellatis glabris, stylis brevibus, nigrescentibus dimorphis conis</sup>  
<sup>obtusis lanceolatis glaucis, capsulis ovatis conis pedicellatis glabris, stylis brevibus, nigrescentibus dimorphis conis</sup>

*S. inermis* Schrank. Stir. fl. 1 p. 230 Quirb. flor. austr. p. 150

*S. fruticosa* ~~Differ.~~ <sup>foliis</sup> ~~foliis~~ <sup>linearibus lanceolatis acuminatis glandulosis mucronatis</sup>  
<sup>marginibus</sup> ~~marginibus~~ <sup>obtusis serratis, acuminatis in ramulis floris apice triangulis, capsulis ovatis</sup>  
<sup>pedicellatis glabris, stylis elongatis, nigrescentibus dimorphis conis</sup>

Syn. *S. viciaria* W. 1 p. p. 4 p. 598 Hort flor. austr. 2 p. 634

Non. Hort sal. tab. 58. 59.

Hb. ad agr. floris Buttschitz ad Solubitz prope Linn, et in agroribus prope Schunig.  
Pl. Agr. Non. ¶





Flor. monici: masculi amaranthi lobuloy, pinnanthi irregulariter bi-nerviati bracte  
 sphaeriformi ducto unifloro sessile, staminibus 3-pluribus saepebus inflexis, filamentis  
 brevibus liberis, antheris erectis, connectivo incomposito et ultra lobos longioribus  
 deliquenti, productis: perianthio in unilobum apicem, calycis ovario ducto limbo saepe  
 4-lobato ducido, petalis 4 calycis limbo inflexi basi cohaerentibus, nullis, ovario imbr  
 rugoso unilobato hyp. bi-quadribulato, ovulo unico in chlamyde apicem sessile erecto, stig  
 m. 1-2 brevibus, stygm. lobis clavatis pinnatis, vel unico sessile peltato quadrilob. Nunc  
 usque ovario aut hinc, rarius ex alio, videtur in limbo saepe.

Abony rari: Nymphaeae imperii-gymnosae coriophyllae, foliis integerrimis serratis  
 graveolentibus.

Nymphaea L.

gen. pl. d. II. p. 354 (encl. p. 11)

Ch. Flor. monici: perianthio 5-6-lobato, staminibus 14-36: perianthio solitario aut gemis  
 glomeratis, corollae, stylis duobus brevibus, stygm. lobis clavatis recurvis pinnatis. Nunc  
 epianthio impubescens saepe, perianthio saepe soluto, ovario saepe in limbo, <sup>lobato</sup> <sup>4-lobato</sup>.

- Cl. Monoclin. ord. Polyand. Syst. 122. -

Nymphaea regia L. p. pl. d. II. p. 1415 Bot. Koenig. Detm. p. 38 Bot. fl. austr. 2. p. 623  
 Reichb. fl. exc. p. 489 Alch. fl. jav. p. 168 Bot. wagn. -

J. Flor. submersis ovalibus <sup>sub</sup>aequalibus subrotatis glabris, pinnatis subfimbriatis.

C. 2. Nunc gen. pl. p. 3. p. 2

Fl. elem. d. I. p. 215

Reichb. herb. t. 247

Method. Xp. ed. Vag. 1568 p.

Mey. Orch.

Herb. <sup>ex oriente</sup> <sup>paucis</sup>

ab. <sup>ab oriente</sup> <sup>paucis</sup> culta. Flor. April. 4

Uf. Nunc doli, lignum densitate firmi ac etiam praestant.

Flor. hermaphroditi: uniuersalis aut polygami, capitati <sup>cyana</sup> spicati <sup>aut</sup> geminati, vel saepe  
 utriusque uniuersalis, saepe in uniuersum. Perianthium inferum 3-6-lobum  
 tunc, laciniis acedens imbricatis. Stamina separata imo pinnatis inflexis, gyp  
 laciniis saepe saepe ad opposita, filamentis liberis, antheris saepe  
 Antheris 1-2 lobulatis, ovulo in lobulo unico, raro binio. Stylis 2-1. Nunc ind.  
 saepe simplex vel compositus, saepe, saepe vel gyp, ovario erecto aut saepe  
 Hinc aut abony, foliis simplicibus peltatis abony, saepe oppositis, stylis liberis saepe  
 2-lobatis

Clavis analytica generum

- |   |  |
|---|--|
| Fructus siccus vid. N. 1.                   | Fructus polygami vid. N. 5                                   |
| 1. fructus athers — 2                       | fructus circumscissus  |
| 2. fructibus spicatis capitatis cymosis - 3 | <u>Ulmus</u><br>fructibus in strabellum membranaceum diffusi |
| 3. seminibus erectis — 4                    | <u>Hamulus</u><br>seminibus pediculis                        |
| 4. perianthiis tetraphyllis                 | <u>Cornula</u><br>perianthiis perianthio monophyllo lobatis  |
| <u>Urtica</u>                               | <u>Pantonia</u>  |
| 5. fructus compressis — 6                   | fructus simplicibus  |
| 6. fructus apice perforatis intus cavo      | <u>Urtica</u><br>fructus imperforatis                        |
| <u>Rhus</u>                                 | <u>Morus</u>   |

Index I Urticacearum Reichb. l.c. p. 181

flor. polygami aut uniaxulari, Monimulibus <sup>ante aut post induritionem</sup> fructu simplici: siccis, seminibus erectis albuminis, embryonibus erectis

Urtica Tournef.

inst. v. herb. p. 536 tab. 308 dicitur fl. perianthio lobato, antheris <sup>perianthio perianthio</sup> ~~perianthio~~ <sup>perianthio</sup> lobatis & lobis <sup>perianthio</sup> ~~perianthio~~ <sup>perianthio</sup> lobatis

Ch. flor. monici aut bicis spicatis aut capitatis, <sup>perianthio perianthio</sup> ~~perianthio~~ <sup>perianthio</sup> lobatis & lobis <sup>perianthio</sup> ~~perianthio~~ <sup>perianthio</sup> lobatis  
<sup>perianthio</sup> ~~perianthio~~ <sup>perianthio</sup> lobatis & lobis <sup>perianthio</sup> ~~perianthio~~ <sup>perianthio</sup> lobatis  
 fructibus compressis perianthio persistente tubo. — U. Mousier. ord. Rhodod. Syst. 185.

Urtica urens L. sp. pl. ed. 2. p. 1396 Hist. pl. austr. 2. p. 613

Reichb. flor. p. 181 Alch. fl. jard. p. 223 Bot. bot. voy. n. 985.

U. fl. <sup>ovata</sup> ~~ovata~~ <sup>ovata</sup> lobatis & lobis <sup>perianthio</sup> ~~perianthio~~ <sup>perianthio</sup> lobatis  
 fructibus compressis, seminibus erectis, embryonibus erectis

Icon. flor. dan. tab. 839

Engl. bot. t. 1236

Mathth. l. 1. ad. Mg. p. 1185

Urtica urens L. sp. pl. ed. 2. p. 1396 Hist. pl. austr. 2. p. 613 Reichb. flor. p. 181 Bot. bot. voy. n. 985.

U. fl. <sup>ovata</sup> ~~ovata~~ <sup>ovata</sup> lobatis & lobis <sup>perianthio</sup> ~~perianthio~~ <sup>perianthio</sup> lobatis  
 fructibus compressis, seminibus erectis, embryonibus erectis

U. fl. <sup>ovata</sup> ~~ovata~~ <sup>ovata</sup> lobatis & lobis <sup>perianthio</sup> ~~perianthio~~ <sup>perianthio</sup> lobatis  
 fructibus compressis, seminibus erectis, embryonibus erectis

Syn. Urtica dioica L. sp. pl. ed. 2. p. 1396 Hist. pl. austr. 2. p. 613 Reichb. flor. p. 181 Bot. bot. voy. n. 985.







Uf. Necty in dependent gradibus caudales form fulvum <sup>extremis aut rubrom</sup> ~~et ducit~~ in unguibus  
sorsipulchris, hyemalibus, rubiculis, sorsipulchris, cubensis. Ductum unum, edent,  
conducentis, sorsipulchris, unum, unum, unum. Praeterea sorsipulchris, sine, hordens  
sive, conuident, conuident, ingenuent.

Tab. III. Moresas

flor. uniuersales, sorsipulchris 3-4, fruct. conuident, sorsipulchris aut, conuident, sorsipulchris  
dele. albumino, conuident, sorsipulchris.

Moresas Noua.

inst. v. herb. p. 589 tab. 562

Ch. Flor. sorsipulchris dente, sorsipulchris: perianthii quadrifidii, sorsipulchris <sup>ovis abster</sup>  
perianthii perianthii 4, sorsipulchris, sorsipulchris, sorsipulchris, sorsipulchris, sorsipulchris  
tamulis, sorsipulchris, sorsipulchris, sorsipulchris, sorsipulchris, sorsipulchris  
sorsipulchris, sorsipulchris, sorsipulchris, sorsipulchris, sorsipulchris, sorsipulchris  
— Cl. Moresas, sorsipulchris, sorsipulchris, sorsipulchris, sorsipulchris, sorsipulchris

Moresas alba L. p. fl. d. II p. 1398 N. sorsipulchris p. 40 Herb. fl. austr. 2. p. 614  
Fuchs. fl. austr. p. 179 Albin. fl. austr. p. 35

M. sorsipulchris alba sorsipulchris sorsipulchris sorsipulchris sorsipulchris <sup>ovis abster</sup>  
sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris  
sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris

Nov. C. sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris  
sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris

Moresas. Sorsipulchris.

Hab. in China et Persia, sorsipulchris in Europa, sorsipulchris. Necty sorsipulchris. H. Moresas. <sup>†</sup>  
Uf. Sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris

M. nigra L. p. fl. d. II p. 1398 Uf. l. c. Herb. fl. austr. 2. p. 615 Fuchs. l. c. Albin.  
l. c.

M. sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris  
sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris

Nov. C. sorsipulchris l. c. p. 2-25.

Lam. enc. bot. fl. 562. t. 1.

Dub. herb. sorsipulchris. 2. tab. 1.

Blaschke. herb. t. 126

Moresas. Moresas caerulea.

Hab. circa Lam, Sorsipulchris, Sorsipulchris in parte? H. Moresas. Necty sorsipulchris. <sup>†</sup>

Nova Noua.

inst. v. herb. p. 562 tab. 420

Ch. Flor. sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris  
sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris sorsipulchris

culi (Coccarthia Nesi jun.) carosae folijs. aut piperis folijs. bantus, <sup>spinae pediculi</sup> agni piperis,   
on spinae de piperis. m. culi piperis hujus, hinc in piperis: piperis piperis 5-6  
gravi pediculi, stylus lobatus, hinc in piperis hujus. In piperis piperis e venteris pediculi  
hujus utriusque piperis in piperis in piperis. — Cl. Pappus ord. piperis hujus.

Flora L. sp. pl. v. 2. p. 1513 Voj. Agr. del. p. 35 Herb. fl. ungu. 2. p. 555. Scit.  
Herc. p. 179 Alch. fl. jud. p. 219 Pet. bot. ungu. 2. 403

Fl. pl. hujus 3-5 lobis subcordatis venteris venteris, venteris hinc in piperis.

Don. C. P. Nesi jun. pl. piperis 2: 6 p. 1-15

Lam. enc. bot. pl. 861

Meib. herb. 1. 225

Meth. Agr. d. Voj. 1568 p.

Myo. ~~hinc in piperis~~, et piperis hinc in piperis.

Hab. spinae ad venteris in piperis et venteris venteris in piperis in piperis hinc in piperis

Uf. Quae hinc in piperis  
venteris hinc in piperis  
venteris hinc in piperis  
venteris hinc in piperis  
venteris hinc in piperis hinc in piperis

Tab. IV. Celticae July bot. pl. 1. p. 221.

~~hinc in piperis~~

Flora pl. hujus, hinc in piperis 4-5, piperis hinc in piperis, venteris hinc in piperis, albuminis  
venteris, embryonis piperis.

Celticae Nesi.

inst. v. herb. p. 612 tab. 383

Ch. Flora hinc in piperis aut venteris venteris hinc in piperis, piperis hinc in piperis, venteris hinc in piperis  
venteris, stylus lobatus, hinc in piperis  
venteris, omnia hinc in piperis  
hinc in piperis hinc in piperis hinc in piperis hinc in piperis hinc in piperis hinc in piperis hinc in piperis  
venteris hinc in piperis hinc in piperis

Celticae Nesi L. sp. pl. v. 2. p. 1478 Voj. Agr. del. p. 30 Herb. fl. ungu. 1. p. 220 Scit.  
fl. enc. p. 179 Alch. fl. jud. p. 65 Pet. bot. ungu. 2. 419.

C. pl. hinc in piperis  
venteris hinc in piperis hinc in piperis

Don. C. P. Nesi jun. pl. piperis 2: 6 p. 1-15

Lam. Enc. bot. pl. 844 p. 1.

Scop. delc. in piperis 2. 44. 18

Canari. egit. pl. p. 155

Meth. Agr. d. Voj. 1568 p.

Uly. Populka. Kostka

Hb. kraj saxony celkij Uly. Delonkie pruzny. Flor. mag. Fl. pers. 4

Uf. Maly Delaj sava, lignum, propter in - per unit.

Obi. In plant delonkie flory ud. kuzny pruzny tetrastyles et humilitas pruzny Dancy.

Uly. v. Ulmeae (Aitl. fl. exc. p. 180

flory hermaphrodit, staminibus 4-5, fructu sarcoisico, seminibus exalbuminis, calycem vultu.

Ulmus Saxon.

inst. v. herb. p. 601 hb. 372

U. Piceanthem <sup>perstans</sup> ~~perstans~~ 4-5 pedum, staminibus ut plurimum 5, quibus superius in pediculis  
antheris supra basin operis, ~~sepalis~~ ~~calycibus~~ ~~dehiscentibus~~. staminibus, stylis brevibus inter  
staminibus utriusque. Laminis corollae circumscissis membranaceis, ab his membranaceis. - U.  
Piceanthem ord. Dignis Syst. 100.

Ulmus campestris L. p. gl. d. II p. 327 Vir. sax. delm. p. 50 Hort. fl. mag. 1. p.  
350 (Aitl. fl. exc. p. 180 Aitl. fl. jud. p. 68 Bot. bot. mag. 1. p. 982

U. flory duplici-ovatis lapi. irregularibus, floris subventris fructibus globis hinc lym.  
florum et hinc. p. 637

a. uirt. : cortex ramorum laevi

Syn. U. campestris Sm. engl. fl. 2. p. 20  
Edm. C. F. Nuss. p. 11. p. 15, 10-16. Engl. bot. t. 1886. Flor. dan. t. 682. Lem. enc. bot. p. 115  
p. 115.

Syn. U. subrosa, Edm. Beitr. 6. p. 87 Vir. Sax. delm. p. 50 Hort. fl. mag. 1. p. 328  
Aitl. fl. exc. p. 180 Aitl. fl. jud. p. 68

Var. Engl. bot. t. 2461

Uly. Brit. et lignum Britanica

Hb. pinnis ad pappi et in montibus utriusque orientis, alibi in praevalentibus ad  
lignum in delonkie hinc in Zan, Schure, Vojvoda. Flor. pruzny.

H. Mart. ligni. 4

Uf. Cortex interior ulmi in asphragmatis lignis et in utroque praevalentibus in montibus  
nisi in Slavoniam ex aqua calida communiter adhibetur, lignum demum pruzny  
stey a floris in usum floris adhibetur.





Ordo XXX Hymelaceae

Juss. gen. pl. p. 26 (excl. Quinquaria)

Floris hermaphrodita, raro dioica, bracteata aut involucre. Perianthium tubulosum, lobis  
liberis, 4-5 perigonis cavosa basi intus striatum, limbo, 1-2 f. lobis aestivatione alternata in  
4-5 staminibus definitis perianthii fere tubosum insertis, eorum laciniis numeris 2-4, non fere  
— in aut copiosioribus, antheris bilobulatis, lobis parvis longius longitudinaliter subpeltatis  
Antherum superum unilobulatum, ovulo unico a loculo apice pendulo, stylo simplex, trigonate in  
pitato. Fructus nucamentaceus aut drupaceus, nudus, aut perianthii basi truncatus non  
— spermiis, semine apice affixo, albumina nullo aut tenui, embryone orthotrope, ~~radice~~  
superiore, cotyledonibus planis convexis, radice supra.  
Fructus, non tubosus, rarius lentibus striatis, foliis parvis oppositis simplicibus indivisis integris  
aut saepe nervosis exsertatis

Clavis analytica generum

Fructus baccatus nudus  
Daphnea

Fructus sine perianthio gemispermus intus  
Passerina

Daphnea L.

Ch. Floris hermaphrodita, perianthio infundibuliformi limbo quadrigono deciduo, staminibus  
octo bifurcatis, stylo brevissimo terminali. Drupe baccata nudus. — Cl. Beckeri ad. Hagen  
1871. p. 106.

Daphnea Meunieri L. V. pl. d. 2. p. 509. B. Bot. Belg. p. 33. Hort. Fl. Mont. p. 46.

Fruct. bacc. p. 164. Bot. Belg. p. 91.

D. floribus lobulatis ~~sublongis~~ sublongis pubescentibus, laciniis perianthii orbis ovatis, foliis  
lanceolatis, petalis perianthii deciduis.  
Linn. Sp. Pl. ed. 2. p. 1284. n. 324. var. L.  
Lam. C. de Nees gen. pl. fasc. VII. n. 2. f. 1-19

Engl. bot. 66. 1851.

Lam. in. bot. p. 290 f. 1.

Flor. dan. 1. 264.

Math. hist. Bot. d. Belg. 1868 p.

Mys. Mözlinia.

Herb. in nemorosis regionibus belgicae montium Velleb. et Nothae. H. Hebraeus, Mont.  
Flor. vici pyrenae. Hic gemmae terminaliter squarrosae, caetera hysterothia.  $\frac{1}{2}$ .

Uf. Cortex trunci et praecipue arboris in aëre, respirante gaudet. Folii ejusdem ore  
decidua duo generis parvula in nobis intant, seropulchra, siphylotica, althuckeque  
conceduntur.

D. alpinus L. V. pl. d. 2. p. 510. Hort. Fl. Mont. p. 476. Fruct. bacc. p. 164. Bot. Belg. p. 91.

D. floribus terminalibus, gemmis nullis, laciniis perianthii baccatis succulentis, foliis lanceolatis  
obovatisque pubescentibus, semine foliis squarrosis deciduis

Don Gesm. of bot. p. 24. t. 33 n. 7.  
Don. Barvel. in fl. 234.

Del. icon. reg. t. 270 p. 1. et aduif. p. 158

Myr. Molinaja  
Hb. in vappatis regni subelgias antius D. M. H. H. *Opavio* p. 158, et horkon

Hoif could upon p. 158. *Fl. reg. reg.* §

D. Fridum L. p. 2. t. p. 511 *Fl. reg. reg.* p. 164 *Alch. fl. reg. p. 91*  
D. fl. by *Conicidib* *agryth* *conicid*, *lavinis* *perianth* *obtus*, *flig* *capit* *lavan* *obtus*  
*lavinis* *perianth* *obtus*, *flig* *capit* *lavan* *obtus*

Don. Sitth. et Sm. fl. gr. cent. IV. t. 66. 356  
Don. Ch. hist. fl. 1. p. 81

Amor. epit. fl. p. 974

Hb. inter fructus long. collig. p. 158 *Callon*, *ind* *concomitant* *Ind. Nov.* *Fl.*  
a *veve* *in* *autumnum*. *H. alb.* *ind.* §

H. Et hujus cortex *apertus*.

*Passina* L. gen. fl. 2. t. p. 134

Ch. Hujus *hemisphaer*, *ovis* *hori*, *perianth* *infundibuliforme* *limbo* *quadripetalo* *persistente*,  
*shimul* *ob* *trifido*, *style* *filiforme* *lobato*. *Uterulo* *monogyne* *perianth*  
*persistente* *indurata*. — *Cl. Alch. in* *ord.* *Manzoni* *Syst. na.*

*Passina* *anura* *Willd.* *in* *act.* *botan.* *1820* *p.* *320* *Alch. fl. reg. p. 91*

P. caule *supra* *strich* *unif*, *flig* *quasi* *lobata* *lineari* *flig*, *florib* *exillit*  
*substant* *persistens*

Jyn. *Willd.* *Passina* L. p. 2. t. p. 511 *Fl. reg. reg.* p. 164 *Alch. fl. reg. p. 91*  
*St. galenus* *Summ.* *fl. in* *ord.* *1. p. 466* *bt. reg. p. 94*

Don. C. 2. *Reg. gen.* *fl. p. 7. a.* f. 1-11

*Ray.* *icon.* *fl. rar.* *1. t. 68*

*Column.* *aphor.* *1. pag.* *82* *f. 1.*

Hb. in *arvis* *agrestib*, *et* *inter* *septis* *in* *h. de* *Lomb.* *Hoif* *g. l. d.* *Fl.* *a* *reg.* *in* *autum*

Hb. *Paros* *caule* *unif* *ans* *subulpiu*, *florib* *ans* *regum* *persist* *et* *flig* *florib*  
*fructu* *agrestib*, *et* *hirculib*, *et* *regum* *longitudo*, *quod* *obtus* *in* *h. de* *in* *una* *cordage*  
*regum*, *et* *veve* *unius* *et* *Becht.* *in* *fl. rar.* *1. t. 68*.

P. *hirsuta* L. p. 2. t. p. 511 *Fl. reg. reg.* p. 164 *Alch. fl. reg. p. 91*  
*Fl. rar.* *p. 91* *Ped.* *bt.* *regum* *a.* *682.*

P. *caule* *lobo* *unif* *ovis* *nutribus* *incans* *lombis*, *flig* *quadrifid* *indurata* *ovis*  
*arvis*, *florib* *in* *h. de* *florib* *sub* *h. de* *axillit* *regum* *agrestib* *agrestib*

Don. *Sitth.* *et* *Sm.* *fl. gr.* *cent.* *IV.* *t.* *66.* *356* *in* *alchem* *fl.* *reg.* *p.* *91*

*Ch. hist.* *fl.* *1. p.* *81*  
*Ray.* *icon.* *fl. rar.* *1. t.* *68*  
*Column.* *aphor.* *1. pag.* *82* *f. 1.*

Hb. in *arvis* *agrestib* *indurata* *Savoye* *in* *h. de* *Flavaria* (*Guarata*), *in* *indurata* *con*  
*Lan*, *et* *in* *una* *Melida*. *Hoif* *in* *fl.* *reg.* *p.* *91* *H. rar.* *et* *autum.* §

Venten. tab. de veg. veget. II p. 245

Flory hermaphroditic aut absteru unisexualy, bracteis aut involucri. Perianthium caly.  
inter 4-6-8-10. Stylis perigynis axillis involucris, lobis alteris stylis  
inter aestivationem imbricatis. Stamina definita, ad numerosa, bitriplicata, styli margini  
inserta, intima abortiva, filamentis interioribus lobis glandulosis, antheris evanescentibus  
lobis simplicibus aut bilobatis connatis lobis immixtis divisis, singulis ovula longioribus  
vix persistentibus a basi ad apicem veritabiliter delinquentibus. Ovium regionem unitam  
ovulis unice a loculis apice pendulis, stylis simplicibus, stigmatibus angulatis. Trochylus brevissimus  
aut displicens vixit vel perianthio insertus, monospermus, semine involso; albumine  
nulla, cotyledone alioquin, cotyledonibus planis concavis, radicle super.

Arbor frondosa foliis alternis simplicibus integris venosis vel penninerviis, saepe coriaceis,  
penninerviis, serratis.

Laurus C. J. Nees et Ef.

Laur. ind. orient. in Wall. pl. orient. II p. 61

Ch. Flory divisa in involucri simplicis involucri styli, perianthio quadrifido. Ovula:  
naves staminibus duobus biserialibus, filamentis omnibus medio bifidulis,  
lobis simplicibus: foeminae staminibus 2-4 staminibus bifidulis heterosporibus,  
ovis ovatis, stylis crispis, stigmatibus subangulatis. Baccis carnosis monospermas. — Cl. Laur.  
scandens ord. Monogynae Syst. sex. —

Laurus utilis L. gen. pl. II p. 329. 1. theop. del. p. 39. Hall. fl. austr. II p. 667. Pl. Ind. Ind.  
exc. p. 185. Alch. fl. jud. p. 93. Bot. univ. p. 538

L. foliis alternis lanceolatis coriaceis subundulatis ovatis, floribus axillaribus simplicibus  
quadriplicatis.

- Non. C. J. Nees gen. pl. fasc. VII n. 2
- Sibth. et Sm. fl. graec. cens. IV. tab. 365
- Lam. Enc. bot. pl. 321 f. 1.
- Math. d. p. 2. Ordo. 1868 p. 146.

Ulyss. Laurika. Looor. Looor.

Hab. ad regem circa Naxos, et Castelli, et insulis Brugg, et Lauria. Pl. a Thunberg in apoc.  
Pl. insid. II

Uf. Mta in pulcherrima parte contra flum. Thunbergii, et Dispersion Lauridensis. Oculum ex baccis  
propagari vixit nunquam contra vixit, ex quo innumera contra tenentur figit adhibetur.

















*R. subty* L. sp. pl. d. II p. 480 Herb. fl. austr. v. p. 488 Reichb. fl. austr. p. 568 Alch. fl. austr.  
 p. 85 Bot. bot. austr. v. 293  
*R. radia* <sup>tenk</sup> *repente* fl. austr. - *lyallii* glaucy, sicuti *divinum* *lancey*, *varia* *laxiflora*, *gibbiflora*  
*in* *hirsutis* *ellipticis* *subtus* *obtusis* *membranaceis* *radialibus* *venis* *callosis*, *floribus*  
*polygamis*

Non. *Zinnia* ~~ist. bot.~~  
*Michx.* *herb.* p. 296  
*Michx.* *herb.* *brit.* p. 200  
*Bot.* *herb.* *1847* p. 292  
*Botan.* *peruv.* p. 649 f. ext. (ead. ac Lab.)

Herb. ad ~~laxiflora~~ *laxiflora* *in* *montibus*, et *Almira*, *in* *un* *in* *montibus* *Moriv*, *Diokora*,  
 et *Orion*. *Flora* *in* *in* *septem*. *Flora* *flavido* *rubra*. &  
 Obs. *variet* *floribus* *divinis*, et *hinc* *prostrata* *R. glaucy* *Sac.* *coll.* *17* *54* et *Non* *var.* *v.* *6* *67*, *floribus*  
*laxiflora* *glandiformibus*, et *hinc* *est* *R. hirsutiflora* *Reichb.* *fl.* *austr.* *caus.* *v.* *290*. *Dem* *quasi*  
*in* *montibus* *ante* *montem* *in* *floribus* *vestra*, *sed* *non* *in* *aliam* *pro* *laxiflora* *drupacea*  
*hirsuta*, *in* *pro* *montibus* *quidem* *hirsuta* *est*.

xx *Lagochum* *Donn.* *ist.* p. 504  
*floribus* *trifloris* *attenuatis* *obtusis* *cordatis*, *fl.* *hirsutifloris*

*R. pulchra* L. sp. pl. d. II p. 477 *Vic.* *1847*. *Donn.* *p.* *46* *Herb.* *fl.* *austr.* *v.* *487* *Reichb.* *fl.* *austr.*  
 v. 569 *Alch.* *fl.* *austr.* *v.* 85 *Bot.* *bot.* *austr.* *v.* 292  
*R. fl.* *in* *pinis* *condensatis* *glandiformibus* *obtusis*, *variet* *divinis*, *floribus* *pinis* *obtusis*  
*intus* *subtus* *obtusis* *pubescentibus*, *obtusis* *callosis*

Non *Reichb.* *ist.* *bot.* *cent.* *v.* *f.* *689*  
*Engl.* *bot.* *t.* *1846*  
*Dillenk.* *ist.* *ist.* *ist.*

*Moriv.* *herb.* *pl.* *2.* *f.* *5* *t.* *22* *f.* *13* *var.* *in* *via* *Lesing*.  
 Herb. *in* *callosis* *in* *callosis* *pro* *floribus* *divinis* *pro* *floribus*. *Flora* *austr.* *Flora* *hirsuta*. &  
 Wj. *Folia* *in* *callosis* *pro* *floribus* *divinis* *pro* *floribus*  
~~*R. subty* L. sp. pl. d. II p. 480 Herb. fl. austr. v. p. 488 Reichb. fl. austr. p. 568~~  
~~*R. fl.* *in* *pinis* *condensatis* *glandiformibus* *obtusis*, *variet* *divinis*, *floribus* *pinis* *obtusis*~~

*R. fl.* *in* *pinis* *condensatis* *glandiformibus* *obtusis*, *variet* *divinis*, *floribus* *pinis* *obtusis*  
*condensatis*  
 β. *divinis*: *floribus* *condensatis* *glandiformibus* *obtusis* *inferius* *lancey*  
 Syn. *R. divinis* L. sp. pl. d. II p. 478  
 Non. *Dill.* *herb.* *peruv.* *t.* *32* *p.* *2*.  
*Herb.* *Schreb.* *Schreb.*  
 Obs. *Hinc* *floribus* *in* *callosis* *glandiformibus* *obtusis* *inferius* *lancey* *et* *floribus* *pinis*  
*hinc* *in* *callosis* *glandiformibus* *obtusis* *inferius* *lancey* *et* *floribus* *pinis*  
*in* *callosis* *glandiformibus* *obtusis* *inferius* *lancey* *et* *floribus* *pinis* *in* *callosis* *glandiformibus* *obtusis* *inferius* *lancey* *et* *floribus* *pinis*  
*in* *callosis* *glandiformibus* *obtusis* *inferius* *lancey* *et* *floribus* *pinis* *in* *callosis* *glandiformibus* *obtusis* *inferius* *lancey* *et* *floribus* *pinis*

N. bupleuifolium L. sp. pl. 2. to p. 429, qui a cl. Host fl. austr. 1. p. 458 sed cl. Purn:  
soblej in Slavaci vivit et satis viri dicitur, nec ibidem usque Thure videretur,  
nec in herbis Bohemolugis surmunt, et in ipso herbis Hathi specimen d. locum  
repperit, cum illuc Turcom excepit edit circa <sup>Misiam</sup> ~~Slaviam~~ <sup>Syriae</sup> ~~Misiam~~.

N. Pakistic L. sp. pl. 2. to p. 426 Host fl. austr. 1. p. 455 Reichb. fl. ex. p. 570  
N. plij <sup>in part</sup> ortho-lanceolatis, <sup>partibus lanceolatis</sup> infimij basi subintegrj, superij lanceolatis, <sup>marginibus</sup> <sup>venis</sup> serratis,  
gymnophylly indivisibily late cordatis <sup>obtusis</sup> integrj <sup>bracte</sup> <sup>convallatis</sup>, <sup>venis</sup> <sup>collatis</sup>

Don. Faorb. 2. p. 1. et sem. 2. ab. 119 p. 2  
Blackw. herb. 2. t. 489.  
Dodon. jussq. p. 648 f. 2.

Illyr. Setavelj. Scholjadh pitomi.

Herb. ad fossos et agrorum marginem circa Duvnj. Host austr. Flor. herbeni. 2.

N. crispif L. sp. pl. 2. to p. 426 Host fl. austr. 1. p. 456 Reichb. fl. ex. p. 570  
Alsch. fl. jud. p. 95. Pet. bot. weyer. 1. 790

N. plij ortho-lanceolatis, <sup>marginibus</sup> <sup>venis</sup> <sup>serratis</sup> <sup>obtusis</sup> <sup>integrj</sup> <sup>bracte</sup> <sup>convallatis</sup>, <sup>venis</sup> <sup>collatis</sup>  
infimij lanceolatis, <sup>marginibus</sup> <sup>venis</sup> <sup>serratis</sup> <sup>obtusis</sup> <sup>integrj</sup> <sup>bracte</sup> <sup>convallatis</sup>, <sup>venis</sup> <sup>collatis</sup>

Don. Reichb. icon. bot. cent. 10. f. 788  
Engl. bot. t. 1998  
Lam. enc. bot. pl. 221 f. h  
Ment. 2. herb. brit. pl. 104.

~~Illyr~~ <sup>Illyr</sup> Setavelj. Scholjadh. ~~Setavelj~~ Konjski setavelj.

Herb. iijdem locis ac procedent, nec cum circa Lam. Schud. Salon. Il. austr. Fl. herbeni 2.  
Obs. Intodon, cum apud nos propter, colleit <sup>venis</sup> <sup>serratis</sup> <sup>obtusis</sup> <sup>integrj</sup> <sup>bracte</sup> <sup>convallatis</sup>, <sup>venis</sup> <sup>collatis</sup>  
cum abothis: Pet. jussq. tendent angustior, et glanior ac in specie.

N. angustif L. sp. pl. 2. to p. 429 Host fl. austr. 1. p. 457 Reichb. fl. ex. p. 569  
N. plij <sup>in part</sup> cordatis <sup>obtusis</sup> <sup>integrj</sup> <sup>bracte</sup> <sup>convallatis</sup>, <sup>venis</sup> <sup>collatis</sup> <sup>subintegrj</sup> <sup>venis</sup> <sup>collatis</sup>

Don. Reichb. icon. bot. cent. 10. f. 555 ost.  
Blackw. herb. 2. t. 490  
Camer. epit. p. 232 <sup>subulata</sup>

Herb. in agris (prope <sup>Magyo</sup> <sup>Magyos</sup> <sup>Magyos</sup> <sup>Magyos</sup> <sup>Magyos</sup>) und nist Subrijus. Host austr. Flor. herbeni. 2.

Obs. Plant. Linum hinc certe spectat, quamvis Linum sub hoc titulum synonyma Tabernemont:  
tenet et Menthae ab <sup>illo</sup> <sup>prope</sup> <sup>alio</sup>, et plij N. Hydrocotyle thes. p. 2.  
angustif. Lam. repentin. Popponij hinc a prof. Alsch. in fl. jud. cum omnibus herbis  
ubi in Slavaci vivit.



*Phytolacca decandra* L. p. pl. 2. II p. 631.

*P. floribunda* Decandif deagguj L. loc. cit. (Hort. fl. austr. 1. p. 563. Reichb. fl. exar. p. 586  
Alten. fl. ped. p. 105

Don. C. P. Nees. gen. pl. p. 344 n. 2

Lein. enc. bot. fl. 593 p. 1.

Dillen. h. elth. 2. t. 239 f. 309-310

Hb. in arenis ad mare circa Alajira, et Lajira, nec non in insula Leima. Flora  
Alajira mihi, bene ab Agave reus condon. Hort. Galii 1. sept. 2

W. Banca nunc Kustorum gestat.

Ordo XXXIV Chenopodeae

Vent. tab. veg. II p. 255

Flora hermograndis aut unicavata ditarsi vel florant. Perianthium calycinum  
2-5 foliolum vel partium, vel 4-5 fidei partium, plerumque partibus aestivatione  
imbricatis, post anthesin saepe allthera<sup>va</sup> metalum, nigricans vel durum carnosum.  
Stamina perianthii partibus minus aequalia et opposita, vel paucior, receptacula aut  
disci hujusmodi pendente inserta, antheris interspersis bilobatis longitudinaliter  
schizocentris. Ovium regimen umbellatum, ovulo unico placenta trilobari inserto,  
aut a periclyo brevi pendulo. Stigma 2-4 stylis filiformibus distinctis, vel basi  
in unum connatis. Ductus uterini perianthii ante aut postato ovulo imbric  
volvitur monospermus, semine reniformi, embryone vix annulari perigyni,  
vix albumen primum ingenti, vix coalescente et albumine equivo,  
radice umbilico proxima.

Herbae aut frutices caulis foliatis, interdum articulatis aphyllis, foliis in vete Thelyp.  
no stipulatis.

Clavis analytica generum

Caule continuis vid. N. 1.

1. perianthii compositis — 2

2. floribus bracteatis — 3

3. perianthii laciniis demum alatis — 4

4. ramibus sub glandulosis insertis

5. perianthii sub glandulosis insertis — 5

6. perianthii sub glandulosis insertis — 6

Salicoria

Caule articulato

Salicoria

perianthii sub glandulosis insertis — 6

floribus abrotinatis — 7

perianthii laciniis demum alatis

ramibus sub glandulosis insertis

perianthii sub glandulosis insertis — 6

Halimolobos

Campylorrhiza

perianthii sub glandulosis insertis — 6

Beta

- |    |   |  |
|----|---|--|
| 8. | <i>Stigmatium longisulcatum</i><br><i>ovoides</i> <i>styligera</i>                          | <i>Stigmatium crispum</i><br><i>ovoides</i> <i>styligera</i>   |
|    | <i>Agathophyton</i>   | <i>Chenopodium</i>   |
| 8. | <i>Stigmatium</i> <del><i>styligera</i></del> <i>speciosum</i> - 8<br><i>styligera</i> 2-4. | <i>Stigmatium</i> <del><i>styligera</i></del> <i>speciosum</i> |
|    |   | <i>Phyllogonium</i>  |
| 9. | <i>Stigmatium</i> <i>brinj</i><br><i>atrylex</i>  | <i>Stigmatium</i> <i>protuberans</i><br><i>speciosum</i>       |

Subord. I Cyclobaea C. A. Mey. in Ledeb. fl. alt. 1. p. 320

seminibus albuminis centrali episperma embryonem geostrophica annulatione cincta

Tribo I. Salicorniceae id. l. c. p. 321

caule articulo, floribus spicatis hermaphroditis

*Salicornia* Tourne.

Coroll. inst. v. herb. p. 51 n. 6. 485

Ch. Long in spec. Arabid immixti, quibus articulo tereti, decussatione oppositi, foliis floralibus cuneatis-vaginantibus bracteatis. Perianthium squamiforme galberrimum carnosum rimae hians, staminibus 1-2, strobiliis pediculis insertis, pericarpis unice, stigmatibus bifidis ac unice, fido, stylo crasso, ovula bivalvi. Fructus articulus conopseus, semina verticalia. — Cl. Monandria Ord. Monogynia Syst. sex. —

*Salicornia herbacea* L. p. pl. 2. t. 5. Hort. fl. austra. 1. p. 32. Reichb. fl. austr. p. 576.

Ch. ch. fl. j. d. p. 1. Pitt. bot. voy. n. 800

l. radice cauma, caule herbaceo erecto rarij appendiculis spicatis, spicis terminalibus pedunculatis, apicibus attenuatis articulis conopseis conopseis, internodijs obconicis, spicis parvis articulis tenuibus apice attenuatis.

Tour. C. 7. Nour gen. pl. p. 110 n. 17 p. 1-5

Lam. Enc. bot. pl. IV. f. 1.

Flor. den. t. 303

Engl. bot. t. 415

Herb. in montibus circa Zao, ad ~~Mariniam~~ *Mariniam*, *Salina*, et ad salinas Lago et Bogno. Flores viridula. Hort. Aug. Oct. ©

*S. frutescens* L. p. pl. 2. t. 5. Hort. fl. austra. 1. p. 32. Reichb. fl. austr. p. 576. ~~Ch. ch.~~

Fl. j. d. p. 1. Pitt. bot. voy. n. 899

l. radice parva, caule suffruticoso rarij erecto, articulis teretibus, spicis <sup>terminalibus</sup> pedunculatis

syn. *Arthrocnemum subrotundum* Moq-Tand. Chenop. enum. p. 111.

Flon Engl. bot. t. 2407

Lam. enc. bot. pl. h. f. 2.

Comer. epit. t. 246.



*S. globa* Mill. gard. Dict. n. 2. Vif. Stip. del. p. 48  
*S. filij* Moench. antip. fructibus immixtis. Feichtb. fl. ex. p. 578  
 Syn. *S. oleacea* var.  $\beta$ . L. p. pl. 2. II p. 1456  
*S. inermis* Moench. meth. pl. p. 318. Feichtb. l. c.  
 Don. C. F. Nees gen. pl. fasc. VII n. 15. p. 22-29  
 Moench. Vif. Stip. n. 5. t. 30 p. 2.  
 Hab. et citat. ut praecedens. Hort. cad. hagen <sup>et illius</sup> 1677 forte immixta. Conspicua prop. Alch. in fe.  
 p. Nr. p. 229. faciente d. Sprengel. © ut 5.

*Ariflex* L.

gen. pl. 2. II p. 385

Ch. Flores polygami chalcidati: masculi et hermaphroditi perianthis aequaliter  
 3-5-partitis, staminibus totidem receptaculis insertis, ovum semper 19 pinis abortivis,  
 seminibus horizontalibus: perianthis distichis oviculis compressis, stylo subulato,  
 segmentibus binis filiformibus. Mucos. uterinis perianthis aequalibus inclusis adnatis,  
 seminibus verticalibus. — Cl. Polygami ord. Monocotyl. l. 1. c. —

+ foliis <sup>utrinisque</sup> concoloribus.

*Ariflex potulacoides* L. p. pl. 2. II p. 1493 Vif. Stip. del. p. 27 1705 fl. austr. l. p. 319 Alch.  
 fl. jard. p. 232 124. bot. reg. n. 163

A. caule fruticoso. <sup>oppositifolius</sup> filij ~~obtusis~~ - densis integris potulacoides utrinque argenteis - 19 pinibus, gemmatis  
 fasciatis, bivalvis dorso uncinatis longipedunculatis <sup>ovoidibus</sup>, perianthis <sup>obtusis</sup>

Syn. *Ariflex potulacoides* Moench. Gard. Champ. cauc. p. 75  
*Helenium potulacoides* Waller. sched. ent. p. 112. Feichtb. fl. ex. p. 576.

Don. C. F. Nees gen. pl. fasc. VII n. 16. p. 16-17.  
 Engl. bot. t. 261  
 Donat. herb. 2. 1. cap. p. 74

Hab. ad littora maris hinc in continente, tunc in insulis praecipue. Hort. aequal. Hort. potul.  
 flav. ~~2. 1. cap. p. 74~~  
 Obs. W. Hallman a prop. Alch. in fl. jard. indicatione hucusque non vid. <sup>var</sup>  
 A. hortensis d. p. pl. 2. II p. 1493 1705 fl. austr. l. p. 319 Feichtb. fl. ex. p. 577 Alch. fl.

A. caule herbaceo erecto filij <sup>obtusis</sup> cordatis - triangularibus <sup>dentatis</sup> - densis <sup>acutis</sup> uncinatis, <sup>gemmatis</sup> <sup>ovoidibus</sup> <sup>perianthis</sup> <sup>obtusis</sup> <sup>19 pinibus</sup> <sup>integro</sup> <sup>dorso</sup> <sup>Gustav</sup>, <sup>inter</sup>

Don. C. F. Nees gen. pl. fasc. VII n. 15. p. 18-21  
 Blackw. herb. t. 99  
 Matth. Nyc. in Diss. ed. Vlyp. 1568 p. 487

Myr. Lobelia.  
 Citatur et omnino ad fructus, et horti in h. Diluvii. Hort. aequal. Fl. hibernica. ©

Uf. Herb. ovium oleacea.  
 Obs. Nomen plantae communiter verbum est, sed utriusque <sup>partem</sup> <sup>etiam</sup> <sup>quam</sup> <sup>propria</sup> <sup>et</sup> <sup>fasciata</sup>.

*Chrysan libanif* L. sp. pl. ed. II p. 1494 Hort. fl. austr. 1. p. 321 Reichb. fl. ex. p. 577

*A. caule herbaceo erecto, foliis oblongis pectatis* <sup>lycei lanceolatis</sup> ~~caulis lanuginos~~ integris indivisib., infimij  
<sup>interjunc</sup> ~~lycei~~ <sup>una</sup> ~~duobus~~ <sup>duobus</sup> ~~perianthiis~~ <sup>perianthiis</sup> ~~fructibus~~ <sup>fructibus</sup> ~~shombis~~ <sup>shombis</sup> ~~orbis~~ <sup>orbis</sup> ~~dentatis~~ <sup>dentatis</sup> ~~longis~~ <sup>longis</sup> ~~minimis~~ <sup>minimis</sup>  
Syn. A. angustifolia Ten. fl. bot. n. p. 258 Reichb. fl. ex. p. 577 P. B. bot. wagen. n. 140  
Ten. 2 Engl. bot. t. 308  
2 Flor. den. t. 1287

1. Bauh. hisp. pl. 2. p. 977 f. 4.  
2. P. Nees 207 p. 11. n. 12 f. 12-3  
Hab. in arenisj montium hincj Alouitj pinguis. Nov. aestate. H. vinty - 0

xx *foliis subij argenteo splanis*  
*A. laevior* L. sp. pl. ed. II p. 1494 Hort. fl. austr. 1. p. 320 Reichb. fl. ex. p. 578 Alou.  
fl. p. 252

*A. caule herbaceo patula mox p., foliis oblongis, infimij oppositj, pectatis, Alouitj*  
*innata densisj, superioribj* ~~hirsutis~~ <sup>hirsutis</sup> ~~oblongis~~ <sup>oblongis</sup> ~~perianthiis~~ <sup>perianthiis</sup> ~~fructibus~~ <sup>fructibus</sup> ~~shombis~~ <sup>shombis</sup> ~~orbis~~ <sup>orbis</sup> ~~dentatis~~ <sup>dentatis</sup> ~~longis~~ <sup>longis</sup> ~~minimis~~ <sup>minimis</sup>  
~~dentatis~~ <sup>dentatis</sup> ~~longis~~ <sup>longis</sup> ~~minimis~~ <sup>minimis</sup> ~~latentibus~~ <sup>latentibus</sup> ~~hirsutis~~ <sup>hirsutis</sup> ~~oblongis~~ <sup>oblongis</sup>

Syn. Engl. bot. t. 165  
Mon. hisp. pl. sec. 5 t. 32. p. 17  
Math. d'ic. in Dion. ed. Valg. 1588 p. 291

β. *diffusa* = *foliis oblongis* ~~in~~ <sup>in</sup> ~~caulis~~ <sup>caulis</sup> ~~perianthiis~~ <sup>perianthiis</sup> ~~obis~~ <sup>obis</sup> ~~latentibus~~ <sup>latentibus</sup> ~~subdenticatis~~ <sup>subdenticatis</sup>  
Syn. A. *diffusum* Tenor. fl. neop. p. 52  
Ten. Gen. fl. nap. t. 251. p. 2.  
Hab. in montium circa Lava: var. lycea hincj circa Pinguis. Flor. aestate. 0

### *Thelygonum* L.

gen. pl. ed. II p. 350

Ch. Flores monocii axillares diujs subterni: ~~maxij~~ <sup>maxij</sup> ~~erectis~~ <sup>erectis</sup> ~~perianthiis~~ <sup>perianthiis</sup> ~~diffusis~~ <sup>diffusis</sup>  
~~compressis~~ <sup>compressis</sup> ~~demum~~ <sup>demum</sup> ~~revelatis~~ <sup>revelatis</sup> ~~staminibus~~ <sup>staminibus</sup> ~~8-16~~ <sup>8-16</sup> ~~imo~~ <sup>imo</sup> ~~perianthiis~~ <sup>perianthiis</sup> ~~imp. orbis~~ <sup>imp. orbis</sup>:  
~~perianthiis~~ <sup>perianthiis</sup> ~~superis~~ <sup>superis</sup> ~~oppositis~~ <sup>oppositis</sup> ~~bracteis~~ <sup>bracteis</sup> ~~bracteis~~ <sup>bracteis</sup> ~~repulis~~ <sup>repulis</sup> ~~perianthiis~~ <sup>perianthiis</sup> ~~clavatis~~ <sup>clavatis</sup>  
~~apice~~ <sup>apice</sup> ~~a~~ <sup>a</sup> ~~stipite~~ <sup>stipite</sup> ~~perforatis~~ <sup>perforatis</sup> ~~sub~~ <sup>sub</sup> ~~antheris~~ <sup>antheris</sup> ~~lateralis~~ <sup>lateralis</sup> ~~emergente~~ <sup>emergente</sup> ~~ovis~~ <sup>ovis</sup> ~~comitis~~ <sup>comitis</sup>  
~~ovis~~ <sup>ovis</sup> ~~longioris~~ <sup>longioris</sup> ~~stipite~~ <sup>stipite</sup> ~~latenti~~ <sup>latenti</sup> ~~stipite~~ <sup>stipite</sup> ~~cyliindris~~ <sup>cyliindris</sup> ~~obtusis~~ <sup>obtusis</sup> ~~multis~~ <sup>multis</sup> ~~draga~~ <sup>draga</sup> ~~monij~~ <sup>monij</sup>  
~~na~~ <sup>na</sup> ~~aeris~~ <sup>aeris</sup> ~~ovis~~ <sup>ovis</sup> ~~verticalis~~ <sup>verticalis</sup>. — *Ch. Monocii* <sup>Ord. Polgandii</sup> <sup>Syst. nat.</sup>

*Thelygonum Cyperanthi* L. sp. pl. ed. II p. 1411 Vif. Kir. Detm. p. 49 Hort. fl. austr. 2. p. 620 Reichb. fl. ex. p. 588 Alou. fl. p. 226 P. B. bot. wagen. n. 974

*Hetero succulenta* ~~obovata~~ <sup>obovata</sup> ~~perianthiis~~ <sup>perianthiis</sup> ~~foliis~~ <sup>foliis</sup> ~~pectatis~~ <sup>pectatis</sup> ~~oppositis~~ <sup>oppositis</sup> ~~shombis~~ <sup>shombis</sup> ~~orbis~~ <sup>orbis</sup> ~~5-6~~ <sup>5-6</sup> ~~pluribus~~ <sup>pluribus</sup> ~~stipitibus~~ <sup>stipitibus</sup> ~~obtusis~~ <sup>obtusis</sup>.

Non. C. F. Nees gen. pl. fasc. VIII p. 1.

Delic. det. du Thel. in Ann. des sc. nat. avr. 1830 p. 370

Lam. enc. bot. pl. 228

Barcl. ic. pl. t. 335

C. Bauh. pinx. p. 59

Hab. ad rivum saxorum, et bca lapid p. tenui cernuntur hinc involucrem proferunt. Paris

Ag. Herg. alt. d. 0

Obs. Novum generum, usque quod a *Triu frumosa* Duchon, scribi debet *Thelyp.*  
inve nec *Thelygonum* uti scripsit ~~invep.~~ Planta ab *Dracosis* ad *Santaluca*, a  
*Jussieu* ad *Waltley*, a *Law* ad *Thomson* ~~relata~~, modo et singularum generum et stylis  
structura a d. ~~Triu~~ ~~frumosa~~ *Cynocaulaceae* recte.

Tab III Anserineae Mey-Tand. char. ~~enum.~~ p. 2

caule continens, floribus hermaphroditis polygamis, persistentibus inaequalibus.

*Chenopodium* L.

gen. pl. ed. II p. 80

Ch. florum hermaphroditis florum chartulata, perianthis aequaliter 5-partitis, tenuibus  
quinque una perianthis infatis, ovario deorsum a stylo, stigmatibus brevibus  
filiformibus. Ductus uterinus perianthis connatis pentagonis obliquis, seminibus  
horizontalibus. — C. Pentaria ad. *Elymus* Syst. sax. —

\* floris integrissimis

*Chenopodium Valeriana* L. sp. pl. ed. II p. 321 Hist. bot. p. 30 Pet. bot. univ. v. 250

Ch. cano-prinifera, floris chartulis ovatis ~~integrissimis~~, floribus conformibus spicatis.

Syn. Ch. solidum Curt. fl. Lond. fasc. III. no. 20. Bull. fl. univ. v. 1 p. 323 Fisch. fl. ex. p. 528 Alch. fl. jard. p. 66

Non. Curt. fl. Lond. l. c.

Flor. dan. t. 1152

Bull. herb. de la p. t. 323

Bleeker. herb. t. 100

Mey. Lobelia smaragdina. Lobelia smaragdina

Hab. ad rivum et videretur vulgarijssimum. Herg. ut hae planta cano-prinifera <sup>facta</sup> floris aequalibus. ~~Paris~~ . 0

Ch. polygamum L. sp. pl. ed. II p. 321 Hist. fl. austr. v. 2 323 Fisch. fl. ex. p. 580  
~~Ch. cano-prinifera~~ Pet. bot. univ. v. 248

Ch. viride, floris ovatis mucronatis nudis, floribus conformibus spicatis, perianthis fructibus persistentibus.

Syn. C. amphiplex Kit. in Schult. oest. fl. v. 458 Fisch. l. c. Hist. fl. austr. v. 324

Non. Engl. bot. t. 1480-81

Flor. dan. t. 1155



- Ch. mundula* L. p. fl. d. II. p. 318 Herb. fl. austr. I. p. 322 Reichb. fl. em. p. 579  
*Ch. vivida* <sup>ov. rhombic</sup> <sup>ov. irregular</sup>  
 fl. viv. in v. <sup>ov. rhombic</sup> <sup>ov. irregular</sup>  
 subcylindric, laminae dimidiatae, seminibus ovatis maxime caeruleis  
 Syn. *Ch. hystrix* Hoffm. p. 30 *Ch. fl. austr.* p. 66? *Pitt. bot. voyer.* n. 256.  
*Engl. bot.* t. 122  
*Mois. hist.* p. 566 31. f. 1. *Ch. vivida*  
*Math. Nyc. in Diss.* I. p. 490  
*Zalozum. ic. stop.* f. 122.  
 Hab. in v. <sup>ov. rhombic</sup> <sup>ov. irregular</sup>  
 661. *Mois. hist.* *Math. Nyc.* *Zalozum. ic.* *Ch. vivida* *Ch. hystrix* *Ch. vivida*  
*Ch. vivida* *Ch. hystrix*  
 5. *Ch. caerulea* L. p. fl. d. II. p. 318 Herb. fl. austr. I. p. 321 Reichb. fl. em. p. 580  
 fl. viv. fl. <sup>ov. rhombic</sup> <sup>ov. irregular</sup>  
<sup>particulari</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup>  
*Engl. bot.* t. 122  
*Flor. dan.* t. 1148  
*Duxb. en. pl. agr.* *Hall.* p. 69  
 Hab. in v. <sup>ov. rhombic</sup> <sup>ov. irregular</sup>  
 1. *Ch. ambrosioides* L. p. fl. d. II. p. 320 Herb. fl. austr. I. p. 323 Reichb. fl. em. p. 579  
*Pitt. bot. voyer.* n. 248  
*Ch. vivida* <sup>ov. rhombic</sup> <sup>ov. irregular</sup>  
 fl. viv. fl. <sup>ov. rhombic</sup> <sup>ov. irregular</sup>  
<sup>whorled</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup>  
 Syn. *Ch. ambrosioides* *Spach* *Ag. phaner.* 5. p. 291.  
*Icon Bav. ic. fl.* 1185.  
*Lancon. hist.* tab 30  
*Mois. hist.* *sect. V. t. 6. 31. f. 8.*  
 Hab. *prope in v. <sup>ov. rhombic</sup> <sup>ov. irregular</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup>*  
*Hf. <sup>ov. rhombic</sup> <sup>ov. irregular</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup> <sup>ov. rhombic</sup> <sup>ov. irregular</sup>*

*Agathophyton* Moq-Tand.

*nov. Ann. des scienc. nat.* *nov. 1834* I. p. 291

- Ch. fl. viv. hermaphrodit. ant. prostratae abrotae ptygami, chartulati, pinnatis aequaliter*  
*oppositis, staminibus quinguis imo pinnatis infertis, ovulis ovatis, style brevi crasso,*  
*involucris 2-4 long. pinnatis, subulatis. Nervi staminibus pinnatis commixtae abrotae, seminibus*  
*viv. — Cl. Pucciniae ord. Agrostis sp. n.*  
*Agathophyton Bonny Heavily Moq-Tand. l. c.*



Syn. Bd. Cista Pet. bot. magw. n. 152 an L. (Exell. syna.)

Non Libb. et Sm. fl. gr. cent. III hb. 254 opt.

Hab. in montibus circa Schier, Prus, palata, Lesina, Mojos: etiam in <sup>montibus</sup> herbis inter  
Velika et Syn. Flor. aed. H. v. v. 24.

Bot.

*Caryophorus* L.

gen. pl. 2. fl. p. 62

Ch. Flor. hermaphroditi abscissati, perianthio quadrifido, laciniis dentibus majoribus curvatis,  
staminibus quatuor sine perianthio insertis, ovario ovato, stylo tenui b. bifido, laciniis  
vix striatis inter stigmatibus. Fructus utriculus compressus perianthio immutato  
incluso, semine utriculo. — Cl. Tetradia O. D. Jussieu Monog. p. 121.

*Caryophorus* ~~montana~~ <sup>mongolica</sup> L. 17. pl. ed. II p. 178 Vif. Kir. Del. p. 29 Herb. fl. austr. 1.  
p. 213 Persb. fl. enc. p. 581 Alsch. fl. ind. p. 34 Pet. bot. magw. n. 190.

L. Jussieu <sup>caulibus floribusque</sup> herbis, foliis <sup>caulibus</sup> brevibus subulatis, floribus axillaribus glomeratis spicatis subsili-  
silibus, perianthio glaberrimo tri-lobato breviter truncato.

Non. C. D. Nees gen. pl. fr. VII n. 9

Call. illustr. pl. t. 57

Lam. enc. bot. pl. 86

Duxb. cent. pl. I. tab. XXVIII f. 1.

β. glaberrimus: subglobosus, immixtus. Mojos-Tand. chem. enc. p. 99

Syn. C. glabra L. 17. pl. ed. II p. 178?

Hab. in montibus, agris, saxosis in h. Delonchi horti: ab insula Veglia ad Mojos-Tand.  
Non. <sup>ab insula Veglia in</sup> Flor. aed. H. v. v. 24

bb. C. glabra L. glabra truncata = foliis <sup>caulibus</sup> subulatis glabris immixtis = ab ipsa truncata  
modo desinata, et involuta Delonchiensis sine h. fl. 1179 proinde videtur  
ad erigendam nossem cum L. Mojos Tand. teste supra posse. Plantae autem  
Delonchiensis haud glabrae sunt Delonchiensis ipsae testibus, <sup>non glabrae sunt</sup> ~~non glabrae sunt~~ fl. aff. <sup>non glabrae sunt</sup>  
Non. <sup>non glabrae sunt</sup> subulatis aligantibus villosa.

Subord. II Spirolobaceae (C. A. Mey. in Led. fl. alt. 1. p. 570  
eminis albumine carotinis parvis vel nullis, embryone parvis.)

*Suaeda* Forsk.

fl. aegypt. <sup>Arab.</sup> p. 69

Ch. Flor. hermaphroditi bracteati, perianthio quinquepartito, laciniis aequalibus cruribus, staminibus  
quinque sine perianthio insertis, ovario digynio, stylo tenui breviter bifido vix curvato. Fructus utriculus  
depresso perianthio laciniis cruribus impletis inclusis, semine horisovale. — Cl. Pentadria O. D. Jussieu  
144. 144.

~~Chenopodium~~ *Suaeda maritima* Mey-And. den. unum. p. 308

1. *herbacea* ~~rufa~~ *diffusa*, folij *caespitibus* *multis* *semicylindricis*, *floribus* *bracteatis*,  
foliis *subtus* *axillaribus*, *semine* *nigra* *tenispermis* *regulis*

Syn. *Chenopodium maritimum* L. sp. pl. 2. II. p. 321 Hort. fl. austr. 1. p. 324 <sup>Meth. fl. Ind. p. 67</sup> Pall. bot. voyag. n. 247  
*Schoberia maritima* C. A. Mey. in Led. fl. alt. 1. p. 400 <sup>Willd.</sup> Reichb. fl. ex. p. 582 C. Z. Nees  
gen. pl. fasc. VII. n. 11.

Icon C. Z. Nees gen. pl. l. c. p. 13-18

Silb. et Sm. fl. gr.  
Engl. bot. t. 1633  
Flor. den. t. 489  
Zamich. off. plant. austr. t. 16.

β. *salsa*: *maior* *caule* *erecto*, folij *oblongis* *ovatis*

Syn. *Chenopodium salinum* L. sp. pl. 2. I. p. 221  
*Salicora salsa* L. sp. pl. 2. II. p. 324 Hort. fl. austr. 1. p. 325  
*Suaeda salsa* Pall. ill. pl. reg. p. 16  
*Schoberia salsa* C. A. Mey. in Led. fl. alt. 1. p. 401 <sup>Willd.</sup> Reichb. fl. ex. p. 582

Icon. ~~Pall. loc. cit. t. 1633~~  
C. Z. Nees gen. pl. l. c. n. 11 p. 1-12  
Pall. loc. cit. t. 1639

Hab. in *paludibus* *maritimis* *in* <sup>Zamich.</sup> *Green*, *et* *Linn.*, *ad* *lacum* *salinum* *Marisje* *prope* *Schnee*  
*co.*, *et* *ad* *salinas* *prope* *Bayna*. *Flora* *aestiva*. *Plum* *indiv.* 0

Obs. *semine* *in* *aliquo* *planch* *tenispermis* *regulis*, *et* *viscibus* *bracteis* *minimis*. *His* *viscibus* *oblongis*,  
*aut* *acutiusculis*, *nee* *diffinitum* *constanter* *exhibent*.

1. *fruticosa* *Dorch.* *fl.* *egypt.* *arab.* p. 30

1. *ligata* *rufa* *erecta*, folij *caespitibus* *brevibus* *cyllindricis* *late* *umbellatis*, *floribus* *paucis*  
*bracteatis*, *floribus* *substituentibus* *axillaribus*, *semine* *nigra* *levis*.

Syn. *Salicora fruticosa* L. sp. pl. 2. II. p. 324  
*Schoberia fruticosa* C. A. Mey. l. c. Reichb. fl. ex. p. 582

Icon. Silb. et Sm. fl. gr. ex. cons. 3. t. 255  
Engl. bot. t. 655

Hab. *ad* *rupes* *maritimas* *in* *Bayna*. *Flora* *aestiva*. *Plum* *indiv.* 7

Obs. *Indurum* *epidermij* *caulicibus* *trijuna* *Aspil* *rimis* *igit* *et* *exterior* *probat* *non* *videtur*  
*speciei* *genus* *sphaeris*, *quam* *interim* *S. <sup>1618</sup> ~~seca~~ <sup>maritima</sup>* *et* *hanc* *exalte* *repat*,  
*et* *in* *no* *Archiv* *der* *Sumpeds.* *sub* *nomine* *S. <sup>1618</sup> ~~seca~~ <sup>maritima</sup>* *et* *hanc* *exalte* *repat*,  
*Habes* *specimen* *in* *Bayna*. *Sum* *legit* *Hort* *civis* *Zam.* *in* *fidis* *latibus* *caespitibus* *plan*  
*hanc* *delictum* *nigra* *in* *Vin* *bracteis* *communibus*, *quosdam* *addi* *in* *flor. den.*  
*spec.* *p.* *XIX.* *sed* *Indurum* *specimen* *descript* *justo* *specimen* *ad* *salinas* *prope*  
*collection.*

*Salsola L.*

gen. pl. 2. t. p. 81

Ch. *Noyl homogredita* bracteata, perianthis subaequaliter quinque-partita, laciniis  
dorso, nervis tenuibus alatis, stramineis quinque disco glandulosa insertis, ovario  
dorso, style brevi, stigmaeque subultraeque pubescentibus. Nectary utrinque longius  
perianthis sessile quinquealato inaequali, ramum horizontale - U. Pers. in Bd. Dygii  
Vis. 1879. Delm. 46. Syn. rex. -

*Salsola Kali* L. sp. pl. ed. II. p. 322 Hort. fl. austr. t. p. 325 Sicil. fl. ex. p. 583

Abch. fl. jadr. Pitt. bot. voy. n. 803

l. foliis sparsi subultra-tigridis quinquepartitis <sup>laciniis parvulis</sup>, floris axillaribus, laciniis  
albi perianthis amplis subultraeque partibus laciniis

a. hirta : Tour. vll. fl. near. p. 124  
Diffusa, hirta, vngary, albi perianthis brevibus

Leon. C. D. Des. gen. pl. fasc. VII n. 10

Enfl. bot. t. 634  
Flor. dan. t. 818  
Pall. illustr. pl. t. 28 f. 2. et t. 29  
Math. Nf. in Diss. 2. Valgr. 1568 p. 1089

β. glabra : Tour. l. c.

subultra, glabra, indivis, albi perianthis longioribus

Syn. l. Noyl L. sp. pl. ed. II. p. 322 Nip. Kirp. delm. p. 46 Sicil. fl. ex. p. 583 Abch.

fl. jadr. p. 67 Pitt. bot. voy. n. 804

~~Leon. Salsola Kali in fl. jadr. p. 67 (sub Kali)~~  
~~S. vngary et in vngary p. 124~~  
~~fl. hirta.~~

Leon. Casab. ~~tab. 266~~  
Delach. hupp. pl. 2. p. 1477

Abch. ic. 1679. p. 492  
Math. Nf. in Diss. 2. Valgr. 1568 p. 1089

Meyr. Smakalj.

Hab. ad ora maris in arenis circa ~~Leon~~ <sup>Craggus</sup> ~~Salsola~~ <sup>Castelli</sup> ~~Noyl~~ <sup>Noyl</sup>, ~~Nudon~~ <sup>Nudon</sup>, ~~Caput~~ <sup>Caput</sup> ~~Castro~~ <sup>Castro</sup>  
vix. circa ~~Noyl~~ <sup>Noyl</sup> et in vngary ~~Leon~~ <sup>Leon</sup>. Florae aestate. Noyl subarct. ☉

Obs. Albi perianthis forma et magnitudine variat, et quaedam in specie sunt perianthis  
longioribus, quaedam in vngary, et cetera, ut ex aliis quibus in vngary, et ex  
Descriptionibus auctorum qui eas deserviunt, congerunt, accipit.

l. Soda L. sp. pl. ed. II. p. 323 Nip. Kirp. delm. p. 46 Hort. fl. austr. t. p. 325 Sicil. fl. ex. p. 583 Abch. fl. jadr. p. 67

l. foliis alaribus cylindricis mucronulatis caudis velutis, floris alaribus geminis

alij perianthii <sup>briviter</sup> ~~sp. ...~~ <sup>glacispermis</sup>

Non Jacq. herb. ind. 1. t. 68

Pall. Enum. pl. t. 30

Labl. Icon. p. 394

Herb. presertim vix in meridij sine East, Salma, et Nyssa. Hort. aestiv. King  
alkh. 0

Uf. Ex hujus plantis invenitur, sed ex procedentibus quibus, abierunt florum matrem  
non solum abstrahitur.

Halozeton C. A. Mey.

in Led. fl. alt. 1. p. 387

Ch. Hort. hermaphroditis hereditatis, perianthii subpeltatis quinquepartitis, laciniis demum  
induratis tenuibus, alatis, immixtis subquibus vegetantibus impati. bracteis utrinque compositis  
semine univulvatis. Cetera ut in tabula. — U. Antandrii ord. Dignis 1828 var.

Halozeton sativum Mey-Tand. chen. monogr. p. 188

H. herbaceum ramis diffusis foliis incernis, foliis alteris teretibus immixtis  
carnosis glabris, floribus alteris <sup>glomeratis</sup> pentandris.

Syn. Tabula sibirica L. sp. pl. ed. 4. p. 323. Vif. Voy. Nov. p. 46 Hort. fl. austr. p. 325

Reichb. fl. roc. p. 583 Alch. fl. judr. p. 67

Don Cat. bot. pl. t. 291

Cotitur inter olea. Hort. aestiv. King vix. 0

Ordo XXXV Amaranthaceae

A. Br. prod. fl. nov. Holl. p. 413

Hort. hermaphroditis vel abortu univulvatis, fomentis, bracteis involutis. Perianthium calycis  
cucurbitaceum semispermis propter immixtos 3-5-phyllum, phyllis aestivatione alba in  
bracteis. Stamina hypogaea glomerata numero aequalia, lacinis vel basi monodelphis, semine peltatis  
vel alteris stentibus, antheris introrsis, raro affixis univulvatis longitudinaliter dehis-  
centibus. Ovarium superius univulvatum uni-loculariter, ovulis peltatis dyonitibus ex imo  
ovarii vix singillatim appressis. Stylus simplex, raro subulcus, stigma unice integro,  
aut lobato, aut trifido. Bracteis utrinque utriusque irregulariter vixis, aut pappi formam dehis-  
centis, semine lobulato-veniformi verticillate appressis, embryone arcuato aut angu-  
lari albumen primarium cingente, radicle umbilico proxima.

Herbae (nonnullae) ramosae, foliis alteris simplicibus integris penninerviis serratis  
vel breviter petiolatis, exstipulatis.

Clarior analytica generum

Hortis ~~polygones~~ tribracteis feminis  
Amaranthus

Hortis tribracteatis strobilatis  
Polygnum



Hab. ad vias, et in cultis circa Schinia, Trau, Caltan. Flory pallide viridis. Fl. asept.

Obs. Spadix hujus floris viscosus acribus, ut accedens omnes affricat, id penitus  
infra subglaberrima, patenter raris, quae uter ab omnibus amarantibus diffinitur

A. retroflexus L. sp. pl. 2. p. 1407 Vig. Ruy. Delm. 24 Herb. fl. austr. 2. p. 613 Reichb.  
fl. exc. p. 585 Alch. fl. jud. p. 68 Pet. bot. weyr. n. 51

A. annuus, caule erecto glabro, vix declinans - asperatus, foliis ovatis - acuminatis apice obtusis  
nerviatis, venis, floribus parvis, axillaribus terminalibus spicatis, spicis hemisphaericis  
longioribus ~~longioribus~~ styloideis congestis, stylis parvis obtusis - hookii, tuberculis  
acuminatis singulis terminalibus.

Icon. Willd. Reichb. Icon. bot. cent. v. p. 668

Willd. hisp. amarant. f. 21.

Myr. Schiv. Schiv.

Hab. utriusque in cultis vulgarissimum. Flory pallide viridis. Fl. asept. ☉

Uf. Hecus junius cetera inter olea viliora et communiter adhibetur.

Obs. Praeter hanc speciem frequentius videtur in Dolomiti montana A. gemulatus  
L. sp. pl. 2. p. 1406, praesertim circa Pokrovnik, Miovich, <sup>in una cum Caltan</sup> Vestlika, sub  
nomine illyrica Schiv castrensi, Drator, Rumini videtur, immo sponte vixit  
ajunt indigende in monte quodam prope Vestlika <sup>in vicinis</sup> floris nomine Schivoviz  
ajunt de hac ut de alia planta agitur explorari vultus  
Nichtus quod ~~hanc videtur~~ Colitur prope in insula Lerina A. hybridus  
L. loc. cit. in ~~hanc videtur~~ Dulis iust.

### Polycnemum L.

gen. pl. 2. p. 17

Ch. Flory hermaphroditis, perianthis pentaphyllo, staminibus 1-3 longius tenuis, basi in annulo  
caulibus, tuberculis caulis induratis. - Cl. Triandria Ord. Monogynia Syst. nat.

Polycnemum arvensis L. sp. pl. 2. p. 50 Vig. Ruy. Delm. p. 44 Herb. fl. austr. 1. p. 41  
Reichb. fl. exc. p. 583 Alch. fl. jud. p. 16 Pet. bot. weyr. n. 716

P. foliis rigidis triquetris - subulatis imbricatis averis Reichb. l. c.

Icon C. F. Nees gen. pl. fac. VII n. 19

Jay. fl. austr. tab. 365

Myr. Felica

Hab. hujus viridis, thalicti, averis, ad vias a Larn ad Ragusan. Flory viridis. Fl. asept. ☉

Scleranthaceae

Benth. Beitr. II p. 153

Ch. Pteris hexagonata var. distans Benth. Perianthium papillosum 4-5 folium tubo  
corollae, lobis aestivatione imbricatis. Stamina 1-10, alterna, supra filamentum, ovum  
in anthera perigynum fere perianthii adnatum basi connatis, anthera  
imbricatis deorsum subulatis, loculis non longitudinaliter dehiscentibus. Ova  
in ovum unitularem unilocularem, stylis sagittatis brevibus, aut stylis  
ovis apice emarginatis. Fructus utriculus monogermis (~~Scleranthus~~) curvis imbricatis  
super perianthii tubo induratis et fere univerte inclusis, libris seminibus  
liberis et apice perianthii a loculis fere appendicibus pendulis, embryone annulari  
abnormi primario ingente, radicula umbilico proxima.  
Herba caulis viridibus nodis articulatis, foliis oppositis basi connatis simplicibus integris  
nerviis.

Scleranthus L.

gen. pl. Det. p. 153

P. Willd.

Ch. Stamina decem, filamentis alterna, setaeae multae, ~~ovum~~ <sup>ovum</sup> bivalvatum  
bivalvatum, stylis duobus distinctis apice inter se junctis. Utriculus mono-  
germus. — Cl. Decand. Ord. Dypsis Syst. —

Scleranthus annuus L. gen. pl. II. p. 580 Vef. Sup. N. n. p. 47 Herb. fl. europ.  
I. p. 514. Nees. fl. europ. p. 565

S. perianthii laciniis antheris angustissimis <sup>membrae</sup> marginatis, tubum appendicibus, punctis prof  
pachylis

Nees C. F. Nees gen. pl. p. VIII n.

Engl. bot. t. 331

Flor. dan. t. 504

Herb. in ovum et unguis imbricatis, Chap. et propag. Sibirica, Syriaca, Caesarea, Cathaca. Pteris  
herbar. N. aestiva. ©

Obs. Caulis in nostris vix pubescenter ut in illo Engl. bot. fl. h. p. membrana  
marginata et ciliata. Ab hoc forma non differt Scl. h. p. p. 331. Bot. d. fl. sic.  
p. 65.