

88. Caulis ^{in 15} ~~in 15~~ foliis oppositis teretibus, in quadrangulos foliis teretibus involucriis parvis trijuncis. Petioli
aliquando rugulati vincti; lenticuli etiam spatulati, ovati ^{lobi} ~~lobi~~ ~~lobi~~ aut haud
aggravati pinnatis breviter lanceolatis aut ovato-lanceolatis instructis; petioli
assillae foliis sub longioribus. Petioli ovati-acuminati ovati-guttulati foliis
multis brevioribus. Ab hoc differt *S. Delphinium* L. petioli oppositi brevissimi
pinnatis et haud ovatis ovatis brevioribus.

Phalopus fl. inc. v. g. 5th.

Syn. *S. deltoideum* Ten. fl. veg. prov. p. 6. — Non. Little et Sm. fl. gr. v. t. 446. Column.
 phlegm. p. 42 fig. dexter. Camer. herb. t. 2.

Obs. Inflorascens variet nunc intra secundae racemiformis, runc cymoso-
tripita: color florum interdum albidus: capsulae duae mollior patentes rufae.

Sedum Caprea L. fl. pl. 2. II. p. 617. Herb. fl. a. p. 1. p. 548. Phil. fl. ex. p. 549.

Syn. S. galinsburyi Laburnet. ed. luyd. p. 12. All. fl. ped. 2. p. 120. *S. tosylifolium* Litch.
S. galinsburyi Laburnet. ed. luyd. p. 12. All. fl. ped. 2. p. 120. *S. galinsburyi* Laburnet. ed. luyd. p. 12. All. fl. ped. 2. p. 120. *S. galinsburyi* Laburnet. ed. luyd. p. 12. All. fl. ped. 2. p. 120.

Waldst. et Kit. l. c. p. 104. Bonann. synth. iii. t. 81. f. 2. ^{Cur. in f. p. 11. t. XVIII.} ^{Cur. capit. p. 623}
Hab. in vegetabilibus umbrosis Moraviae, Leptane et Pongau prope Catharo, nec non circa
Musch. Hort. acrole. H. albid., semis pablorum vellellif. ③
8. pedicelli longioribus autatis

Sedum olympicum Boiss. Diagn. pl. nov. orient. III. p. 16.

Hab in vegetabilis montis Oien ^{continuum, hucus} et iuvnet Fr. Neumayer. R. albis. H. Augusto 2.
Obs. Plantae parva, caules teretes ascendentes, tomentosius nigrescentes, apice ramorum, plus minusve

Cotyledon ascendens

C. - leaf inferiorly spatulate-obovate
flowing ^{lance} recurved, ~~long pedicellate~~, ascendingly, pedately
bracteae integerrimae subequalibus, corolla tubulosa calyce triplici
laminis quereatis lobis ovatis apiculatis erectis.

Ab. con. Cephala. d. Catron. Visit Androm. Aug. 1851.

affinity C. Umbellata brief, a few diff. parts flow ascending
are pedately, corolla uncinata per tubulos. campanulata
in apice lobatis, ⁺ prope pedis quereatis.

Myriophyllum verticillatum L. p. pl. 2. II. p. 1410. Hart fl. austr. 2. p. 819. Fieb.
fl. ex. p. 633. Handel-Mazzini basenig. Linnaeus

N. filix masculina ~~capitata~~ peristoma-pastosa laevis capitulo, florum viridibus axillis
 sub breviter ~~pilosis~~ floribus pediculis

G. F. Steyerger: f. VIII n. 13 f. 20
 Germ. - Slov. - Engl. - Lat. 218. Clus. hist. pl. p. 252.
 med. ind.

G. p. leafy pet. VIII n. 137 20
clav. hisp. pl. p. 252.
Syn. M. pectinatum DC. fl. fr. v. g. 59.
Vern. S. pectinatum: folia floribus floribus paulis
Des. H. d. t. 1046

S. pectinatum is *glab. flor. pale* ~~long~~ —
H. viridulum Vahl in *Ord. fl. den. p. 1806* 5 — *Syn. Ord. fl. den. 1. 1806*

M. crystallinum Vahl in Ord. fr. den. p. 180 p. 3 - Syn. 1797
 Hab. in aquis lente florentibus ad ~~lucum~~ lacum: *Diagnosis* page 220: var. *var.*
 Nannh. fl. aestate. 7.
 C. G. C. 194.

trib. II. Ingeae Coll. et Germ. ~~1944~~ ^{1. c. g. 1944.}

~~New York. Aug 19th~~

Phryx hemaphysoides *marcescens* *trichosetula*, *hemimachus* *quatuor*, *ovis* *montanus* *1876*
filipini, *stipitatus* *capitata*.

Traga a.

Gen. pl. 3. p. 10

Gen. pl. 2. II. p. 10. *Prunum blanda. Mu.*
 Char. Calyx tubo ~~lapp~~ adnato, limbo quadrijanito persistente. ~~Corolla tubulosa~~
 tubo corollae unilobato, monospermo, calycis limbo aucto indiviso
 quadrijanito. — Claps. *Retrandu* Ord. *Monggini* Gert. sex —

Dryas octata L. sp. pl. 2. II p. 175 (excl. syn. (Heed.) - Hort. fl. angl. 1. p. 21. Publ. fl. europ. 633

Alsch. H. jadv. p. 34.

Aletris fl. jod. p. 86.

D. plijf, vromerijf gezwakt-gerukt leuning capellancij, emerlijf vromelijf sterking dautelij,
gehoelj mitje iustelij, fruchtig vromelijf, sping gheelij dampgelyk vromen rubij.
L.F. Nies gen. pl. VIII n. 15
Don. Lam. enc. bot. pl. 78. Mattheis. in Dijk. p. 102. — Myr. Oras cadden.

Don. Lam. enc. bot. pl. 75. Matth. in Dr. p. 102. — Myr. Oak wood.

Hab. in laeue Nadin. Fl. alb. Fl. aegre - 4.

Upes. Semen *Fulicaria pondusum* edule.

Tab. III. Hippuridae DC. prod. III. p. 21.

~~Page 19th not. 3 of 74.~~

Platy hermaphroditi nanae, galei apertae, brachia unius, ovarii uniloculari, stylis plurimis perispermatis.

Higgins L.

Gen. pl. 2. p. 1. 4.

[illegible]

Hippuris vulgaris L. p. pl. ed. tt. p. 6. Kest. fl. austr. 1. p. 1. fideb. fl. ex. p. 632.

Her. foliis serrulatis 8-12 linearibus.

H. folii unstratifié 8-12 linéaire.
C.F. Neesgen. ^{N. 12.} Dec. f. 87. Low. enfl. bot. t. 763. Bull. herb. de la fr. pl. 365.
H. folii unstratifié 8-12 linéaire.

Hub. in ^{Salvador} Florif. Naveute. H. Majo, Naveute. 4.

Lythrum Salicaria L. sp. pl. ed. II. p. 640. Hist. fl. austr. I. p. 585, fl. flex. p. 640.

Bl. h. j. ad p. 104. Pet. bot. ungu. n. 583.

L. ~~crataegae~~ ^{subsp.} ~~fl. cordato-lanceolatis~~, ^{subsp.} ~~floribus~~ ^{subsp.} ~~dodecandris~~ ^{subsp.} ~~unibracteatis~~ - ^{subsp.} ~~speciosis~~, ^{subsp.} ~~calycibus~~ ^{subsp.} ~~obtusatis~~.

Icon. bot. angl. bot. t. 106. Des. fl. den. t. 67. Lam. enc. bot. pl. 408 f. 1. Method. in Dispo. p. 1000. Benth. n. 120.

Var. β . *canescens*: fl. ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~obtusatis~~. - ^{subsp.} ~~Lythrum~~ ^{subsp.} ~~canescens~~ ^{subsp.} ~~fl. austr. p. 332~~ ^{subsp.} ~~fl. austr. p. 332~~

Hab. ad flos et rivas lacuum palustris circa Zara, Velletra, Salona, Narona, Cattaro.

et in insula Pag. fl. ab aestate in autumnum. fl. violacei. ^{subsp.} ~~fl. austr. p. 332~~

Ob. fl. valent ovata lanceolata, linearis, lanceolata, opposita verticillata, are clavae: 18-20 uterque ut.

Usp. radix et folia viridia asperguntur pollent, hinc ex aqua ducta praesent in diarrhoea

profluensque mucus.

Lythrum ~~Thymifolia~~ ^{subsp.} ~~Thymifolia~~ L. sp. pl. ed. II. p. 642. fl. flex. p. 640.

L. ~~crataegae~~ ^{subsp.} ~~fl. oblongis~~ ^{subsp.} ~~lanceolatis~~ ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~floribus~~ ^{subsp.} ~~dodecandris~~ ^{subsp.} ~~unibracteatis~~ ^{subsp.} ~~calycibus~~ ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~speciosis~~.

~~fructibus~~ ^{subsp.} ~~Lythrum~~ ^{subsp.} ~~Thymifolia~~ ^{subsp.} ~~fl. austr. p. 332~~ ^{subsp.} ~~fl. austr. p. 332~~

~~L. Bauh. h. p. 3 p. 192 f. 3~~

~~Hab. Lythrum Thymifolia~~ L. sp. pl. ed. II. p. 642. fl. flex. p. 640

L. ^{subsp.} ~~fl. oblongis~~ ^{subsp.} ~~lanceolatis~~ ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~floribus~~ ^{subsp.} ~~dodecandris~~ ^{subsp.} ~~unibracteatis~~ ^{subsp.} ~~calycibus~~ ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~speciosis~~.

Icon. Benth. icon. 1849. 223. f. 2. L. Bauh. h. p. 3 p. 192 f. 3.

Var. β . *lysiptifolia*: fl. ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~lanceolatis~~ ^{subsp.} ~~linearibus~~ ^{subsp.} ~~floribus~~ ^{subsp.} ~~dodecandris~~ ^{subsp.} ~~unibracteatis~~ ^{subsp.} ~~calycibus~~ ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~speciosis~~.

L. *lysiptifolia* L. sp. pl. ed. II. p. 642 non herb. Hist. fl. austr. I. p. 586. fl. flex. p. 640.

- Icon. Bauh. fl. austr. II. t. 233. Bauh. h. p. 3 p. 192 f. 3.

Hab. in humidis et inundatis, nec in in palustribus salsis, circa Zara, Spalato, Narona.

et in insula Zara. fl. violacei exiguus. fl. aestivum.

Ob. Plant. hujus ptyonoghe variis caulis erectis vel ascendens, folia latiuscula

vel angustiora, floris ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~lanceolatis~~ ^{subsp.} ~~linearibus~~ ^{subsp.} ~~floribus~~ ^{subsp.} ~~dodecandris~~ ^{subsp.} ~~unibracteatis~~ ^{subsp.} ~~calycibus~~ ^{subsp.} ~~obtusatis~~ ^{subsp.} ~~speciosis~~.

quinta senaria in eodem L. *Thymifolia*. Hinc *Thymifolia* speciosa propter nullam

not. In herbis *Lythraea* pro L. *Thymifolia* sicut L. *Lythraea* *Lythraea* *Lythraea*

multa majoribus et floribus dodecandris *Lythraea*.

Ord. XII. Onagraceae

Russ. ann. du Mus. 3. p. 215 (Oxyl. gen.)

Flor. hermaphroditi subregulans. Calyx monophyllus ovatis adnatis, lobis 4-pidis, dentibus aspi-

ratione velatis. Corolla tetragynala calycis fere inserta, asperitudo imbricata-lanceolata

aut nullis. Stamina octo aut quatuor non fidele affere, antheris interis bilobulatis longis

tubulosis distinctis. Pistillum ex ovario parvis interis et uteris calycis connatis

unilobulatis multilobulatis, stylis plurimis, stylis quatuor parvis vel in clavae aut

capitulum coarctatis. Capsula quadrilobis quadrilobis, loculis pluribus distinctis, semine

bus parvis vel ascendens clavae unilobis affixis connatis aut calycis, aut clavae,

embryone recto, radícula ad latera vasa.

embryonic roots, rudiments of future leaf.
Hence filix singularis, castigulatus, oppositi, albae, floris, axillaris, solidus, et
necnon terminalis.

Epilobium L.

Gen. pl. d. II. p. 133.

Obs. Calyx tubo longioribus ovibus corollae aequante, limbo quadrijavito circumscissis deciduo. Corolla tubulosa. Stamina octo. Stylis filiformibus rhynchophyllis quatuor unguibus aut in lacum coactis. Capsula baccata, seminum coram sericea. — Alp. Oran, din ord. Monogyne Syst. sex. —

Sect. I. Champsignon Gauch hort. canol. p. 1.

Pisula vix conjugata ^{affinis} *Stylippus senum reflexo*

Epilobium angustifolium L. p. pl. v. II p. 99. Hist. fl. austr. v. I p. 462. Phleb. fl. ex. p. 632.

E. glabrum, fls. sessile, corollary parts ^{very} subequal, petals ^{very} spreading, fls. yellow.
 seeds - narrow, ^{very} strongly curved. Rt. brown. 50.

Syr. E. spicatum Lam. enc. bot. II. p. 189. — *Non. Non* engl. bot. t. 1947. Coll. et germ. l. c. alt. 16. 12 p. 1. Lam. enc. bot. pl. 278 f. 1. Ord. fl. dan. t. 280.

Var. β . - ~~lobatifolia~~ - plicifolia - lanceolata ~~pubescentia~~ - lync. & angustiplum β . Lycop. p. 694.

Hoof. in 14 lottig wettig Borkers. H. magni perquis. H. asperiti. 4.

Sect. II. *Lysimachion* Darsch loc. cit.

Petala biloba. Stamina stylisq[ue] erecta

Egglestem hispidum L. 1. p. 12. II. p. 194. Hall. fl. austr. 1. p. 468. Phil. fl. ex. p. 636. Alsch. fl.

E. Myricaria ^{Jard. p. 91.} ^{pubescens} ^{argentea} ^{officinalis} ^{Myrica-lanceolata} ^{argentea} ^{femina}

Scor. Col. et Germ. alb. et. hb. XII f. B. Low. engl. hb. t. 838. Des. fl. Don. t. 826. Minj. oxon. rec. 211.
hb. 11 f. 3. Puck. hsp. p. 491.

Feb. 11 p. 3. Nicht bes. p. 491.
Hb. Leip. edif. 19. leinisch et ad vivas citra Jan. Lanthier, Dornitz, Versteke, Irm. Naur.
Fl. magis purpurea. H. aequalis. 2.

Epilobium parviflorum Schreb. *Spit. fl. lif.* p. 146. *thit fl. austr. i. p.* 468. *Nob. fl. ex. p.* 636.

Epilobium ^{pubescens} ~~pubescens~~ ^{subulatus} ~~subulatus~~ ^{oblong-lanceolatus} ~~oblong-lanceolatus~~ ^{ramis} ~~ramis~~ ^{prostratis} ~~prostratis~~ ^{floribus} ~~floribus~~ ^{calycibus} ~~calycibus~~ ^{multis} ~~multis~~

Lynx. *E. leucurus* B. L. p. pl. p. 494. *E. melleo* Zimm. enc. bot. II. p. 371. Coll. et Germ. fl. garif. p. 189.
E. jubatus Both test. fl. germ. I. p. 607. — *E. pelusio* Vrijhof. Atlas Ind. E. p. 371. Coll. et Germ. atl. et. t. xii f. C. low.
E. jubatus Both test. fl. germ. I. p. 607. — *E. pelusio* Vrijhof. Atlas Ind. E. p. 371. Coll. et Germ. atl. et. t. xii f. C. low.
 conf. bot. t. 495. Dec. fl. den. t. 547. Ben. p. 189. f. 22. f. Minif. oxon. rec. t. 11. f. 11.

Hub. in humilis et umbrosis circa Salona et in insula Pago. H. d'Arbo. viciat. H. asper. 74.
 1849. 1850. 1851. 1852. 1853. 1854. 1855. 1856. 1857. 1858. 1859. 1860. 1861. 1862. 1863. 1864. 1865. 1866. 1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. 1875. 1876. 1877. 1878. 1879. 1880. 1881. 1882. 1883. 1884. 1885. 1886. 1887. 1888. 1889. 1890. 1891. 1892. 1893. 1894. 1895. 1896. 1897. 1898. 1899. 1900. 1901. 1902. 1903. 1904. 1905. 1906. 1907. 1908. 1909. 1910. 1911. 1912. 1913. 1914. 1915. 1916. 1917. 1918. 1919. 1920. 1921. 1922. 1923. 1924. 1925. 1926. 1927. 1928. 1929. 1930. 1931. 1932. 1933. 1934. 1935. 1936. 1937. 1938. 1939. 1940. 1941. 1942. 1943. 1944. 1945. 1946. 1947. 1948. 1949. 1950. 1951. 1952. 1953. 1954. 1955. 1956. 1957. 1958. 1959. 1960. 1961. 1962. 1963. 1964. 1965. 1966. 1967. 1968. 1969. 1970. 1971. 1972. 1973. 1974. 1975. 1976. 1977. 1978. 1979. 1980. 1981. 1982. 1983. 1984. 1985. 1986. 1987. 1988. 1989. 1990. 1991. 1992. 1993. 1994. 1995. 1996. 1997. 1998. 1999. 2000. 2001. 2002. 2003. 2004. 2005. 2006. 2007. 2008. 2009. 2010. 2011. 2012. 2013. 2014. 2015. 2016. 2017. 2018. 2019. 2020. 2021. 2022. 2023. 2024. 2025. 2026. 2027. 2028. 2029. 2030. 2031. 2032. 2033. 2034. 2035. 2036. 2037. 2038. 2039. 2040. 2041. 2042. 2043. 2044. 2045. 2046. 2047. 2048. 2049. 2050. 2051. 2052. 2053. 2054. 2055. 2056. 2057. 2058. 2059. 2060. 2061. 2062. 2063. 2064. 2065. 2066. 2067. 2068. 2069. 2070. 2071. 2072. 2073. 2074. 2075. 2076. 2077. 2078. 2079. 2080. 2081. 2082. 2083. 2084. 2085. 2086. 2087. 2088. 2089. 2090. 2091. 2092. 2093. 2094. 2095. 2096. 2097. 2098. 2099. 2100. 2101. 2102. 2103. 2104. 2105. 2106. 2107. 2108. 2109. 2110. 2111. 2112. 2113. 2114. 2115. 2116. 2117. 2118. 2119. 2120. 2121. 2122. 2123. 2124. 2125. 2126. 2127. 2128. 2129. 2130. 2131. 2132. 2133. 2134. 2135. 2136. 2137. 2138. 2139. 2140. 2141. 2142. 2143. 2144. 2145. 2146. 2147. 2148. 2149. 2150. 2151. 2152. 2153. 2154. 2155. 2156. 2157. 2158. 2159. 2160. 2161. 2162. 2163. 2164. 2165. 2166. 2167. 2168. 2169. 2170. 2171. 2172. 2173. 2174. 2175. 2176. 2177. 2178. 2179. 2180. 2181. 2182. 2183. 2184. 2185. 2186. 2187. 2188. 2189. 2190. 2191. 2192. 2193. 2194. 2195. 2196. 2197. 2198. 2199. 2200. 2201. 2202. 2203. 2204. 2205. 2206. 2207. 2208. 2209. 2210. 2211. 2212. 2213. 2214. 2215. 2216. 2217. 2218. 2219. 2220. 2221. 2222. 2223. 2224. 2225. 2226. 2227. 2228. 2229. 2230. 2231. 2232. 2233. 2234. 2235. 2236. 2237. 2238. 2239. 2240. 2241. 2242. 2243. 2244. 2245. 2246. 2247. 2248. 2249. 2250. 2251. 2252. 2253. 2254. 2255. 2256. 2257. 2258. 2259. 2260. 2261. 2262. 2263. 2264. 2265. 2266. 2267. 2268. 2269. 2270. 2271. 2272. 2273. 2274. 2275. 2276. 2277. 2278. 2279. 2280. 2281. 2282. 2283. 2284. 2285. 2286. 2287. 2288. 2289. 2290. 2291. 2292. 2293. 2294. 2295. 2296. 2297. 2298. 2299. 2300. 2301. 2302. 2303. 2304. 2305. 2306. 2307. 2308. 2309. 2310. 2311. 2312. 2313. 2314. 2315. 2316. 2317. 2318. 2319. 2320. 2321. 2322. 2323. 2324. 2325. 2326. 2327. 2328. 2329. 2330. 2331. 2332. 2333. 2334. 2335. 2336. 2337. 2338. 2339. 2340. 2341. 2342. 2343. 2344. 2345. 2346. 2347. 2348. 2349. 2350. 2351. 2352. 2353. 2354. 2355. 2356. 2357. 2358. 2359. 2360. 2361. 2362. 2363. 2364. 2365. 2366. 2367. 2368. 2369. 2370. 2371. 2372. 2373. 2374. 2375. 2376. 2377. 2378. 2379. 2380. 2381. 2382. 2383. 2384. 2385. 2386. 2387. 2388. 2389. 2390. 2391. 2392. 2393. 2394. 2395. 2396. 2397. 2398. 2399. 2400. 2401. 2402. 2403. 2404. 2405. 2406. 2407. 2408. 2409. 2410. 2411. 2412. 2413. 2414. 2415. 2416. 2417. 2418. 2419. 2420. 2421. 2422. 2423. 2424. 2425. 2426. 2427. 2428. 2429. 2430. 2431. 2432. 2433. 2434. 2435. 2436. 2437. 2438. 2439. 2440. 2441. 2442. 2443. 2444. 2445. 2446. 2447. 2448. 2449. 2450. 2451. 2452. 2453. 2454. 2455. 2456. 2457. 2458. 2459. 2460. 2461. 2462. 2463. 2464. 2465. 2466. 2467. 2468. 2469. 2470. 2471. 2472. 2473. 2474. 2475. 2476. 2477. 2478. 2479. 2480. 2481. 2482. 2483. 2484. 2485. 2486. 2487. 2488. 2489. 2490. 2491. 2492. 2493. 2494. 2495. 2496. 2497. 2498. 2499. 2500. 2501. 2502. 2503. 2504. 2505. 2506. 2507. 2508. 2509. 2510. 2511. 2512. 2513. 2514. 2515. 2516. 2517. 2518. 2519. 2520. 2521. 2522. 2523. 2

Epilobium montanum L. p. fl. d. Th. p. 494. Hoff. fl. austr. 1. p. 409. ^{var. p. 409. Hoff. p. 409. Hoff. p. 409.}
aut. inat. h. i. abund. in p. 409. Hoff. p. 409. Hoff. p. 409.

E. jubulum, bliz ordo ^{auch} ~~unvollständig~~ ^{bis} ~~schon~~ ^{vollständig} ~~in~~ ^{ausgeworfen} ~~von~~ ^{demselben} ~~als~~ ^{ca.}
auf ~~unvollständig~~, signifikant mutation paternity
Neb. ic. br. B. f. 328.

low. esp. et form. att. fl. par. XII f. D. low. enfl. bot. t. III. Ord. fl. den. t. 922.

Instit. vei herb. p. 6u. t. 381.

Char. Calyx 5-sepalus. Stamina filamentis liberis. Antheris quinquelocularibus, locis 2 locumque
— Clasi. Polyandria ord. Monogynia 1981. 1818.

Attilia Journ.
Macht. v. h. b. p. 611. t. 381.

Pilea ^{subseri} ~~intermedia~~ ^{parvifolia} Ehrh. beiss. v. p. 159. Hort fl. austr. 2. p. 62. fide. fl. ex. p. 828.

9. folijs oblique subrotundo-cordatis ^{acuminatis} pinnis utrinque glabris rubris glaucis et in axillis
nervorum brevibus congestis multifloris, fructu obsolete apato.

Syn. *D. europaea* J. L. 19. fl. ed. 10. p. 253. ~~*D. europaea* Led. fl. ed. 10. p. 253.~~ *D. alba* Muhl.
Willd. en. l. br. 1. p. 565. *D. ulmifolia* Less. fl. car. 1. p. 374. — Don. Quid. in fl.
germ. VI. tab. 311. 312. Led. fl. den. t. 553. Will. cat. h. p. 1. 449. f. 3. Matth. in
Bot. p. 191.

Diagn. p. 191.
Hab. in nemor. marium Vellebith ad Paklenitz alt. 2500 ad 3000 ped. us.
legit Jos. Kargl. H. austri. ♀

~~Titia~~ *grandifolia* *Eschsch. baird. v. p. 158*

Delia platyphyllos Scop. fl. carn. v. II. p. 373. *Delia* fl. ex. p. 828.

7. slief oblique notstand - cordatly uterque flatus subty pectentibus et in axillis nervi
non habet, ingens b-typer.

2. Var. *B. sphaerocarpa* Webb. fl. sax. p. 442: fructu subgloboso ciliolato - Pon. fl. b. v.
fl. gem. VI. t. 316 f. a.

1. Var. σ : *foetata*: fructu ovato acut. turbinate, capsula apert. — fr. *T. europaea* p. 5. S. L.
Bull. herb. v. 11. n. 942. *T. grandifolia* Edh. bot. v. 1858
Sp. pl. ed. 11. p. 233. *T. europaea* Pers. bot. ungu. n. 942. *T. grandifolia* Edh. bot. v. 1858
Sp. pl. ed. 11. p. 233. *T. europaea* Pers. bot. ungu. n. 942. *T. grandifolia* Edh. bot. v. 1858

[illegible]

Ufus. Lignum - gressu toxiaco praesens: flos ex aqua parente i. p. p. benigne adhaerens;
aqua ex eodem distillatione parata inter remota nimis et indolis botulin.

Ordo. XVI Malvaceae
R. Brown in Zuercher lang. p. 428
~~Benth. ex. nat. cl. p. 346~~

Ordo. XVI Malvaceae
A. Brown in Turkey cong. p. 428
~~Barth et al. nat. pl. p. 344.~~

[illegible]

Syn. *G. umbrosum* Walldt. et Kt. fl. vrb. hunc. 2 p. 131. *G. villosum* Desv. fl. vrb. ^{L. XI. LXII.} ~~jacq. p. 166.~~ — Icon. fl. ic. fl. germ. v. ~~p. 166.~~ ^{tab. exch. Den.} fl. vrb. p. 166. Coll. et Germ. fl. par. atl. v. 1. Walldt. et Kt. l. c. tab. 124. ^{Gen. Off. gallego p. 434 tab. 16 f. 6.} Hab. in arboribus et saxosis circa Ima, Spales, Pygus et in rupibus Lemnae. Fl. villosi, aut caeruleo-purp. Fl. vrb. 2. ^{grandiflorus}

Obs. Simile *G. pusillo*, a quo potius quam petalis quadratis, per lacinias calycis vix de-
ant hinc, vix hinc dimidia majore. Effert dentatione perenni nec annua, car-
tilleis nec pubescenti-glanduloso, et petalis vix angusticulis atropurpure longe barbatis, nec
admodum ciliatis. — *G. villosum* Desv. — ^{carpella sunt vix angusticulis atropurpure} carpella sunt vix angusticulis atropurpure
adpressis imbricatis oblonga nec longissimis glandulis seminatibus ^{oblongis} ^{mollis} ^{quales exhibet icon fl.}
neg. t. 166 litt. D. per petalis vix per hunc ~~G. pusillum~~.

xx Annuum

Geranium pusillum L. sp. pl. d. II. p. 957. Hott fl. austr. 2 p. 293. Fl. ic. ex. p. 228.
G. pubescens, foliis orbiculato-venosis 5-7-partitis, segmentis tri-quingulatis, pedunculis bifloris
petalis oblongo-obovatis calycis subaequalibus basi ciliatis, carpellis pubescentibus seminibusque latis
Icon. Fl. ic. fl. germ. v. f. 4877. Coll. et Germ. fl. par. atl. pl. VI. f. D. Vaill. bot. par. p. 15. f. 1. pub.
bist. p. 205.

Hab. ad sepes circa Nagusa. Fl. parvi isolati aut caeruleo-purp. Fl. vrb. in aequalum. 2

Geranium dissectum L. sp. pl. d. II. p. 956. Hott fl. austr. 2 p. 294. Fl. ic. ex. p. 227.
Alsch. fl. jacq. p. 156. Pet. bot. vrb. n. 442. ^{linearibus tri-angulatis}

G. pubescens, foliis orbiculatis 5-7-partitis, segmentis linearibus tri-angulatis, pedunculis bifloris
foliis subaequalibus, petalis oblongo-obovatis calycem aequantibus ^{oblongis} ^{basi barbatis}, carpellis
caudis ^{pubescentibus} ^{seminibus} ^{abundantibus}.

Icon. Fl. ic. fl. germ. v. f. 4876. Coll. et Germ. fl. par. atl. pl. VI. f. C. ^{l. 166. angl.}
bot. t. 753. Cav. monod. typ. tab. 58. f. 2. Vaill. bot. par. p. 15. f. 2.

Hab. in agris et cultis circa Ima, Schinde, Ima, Spales, Pygus. Fl. parvi purpurei.
Fl. vrb. 2

Geranium columbinum L. sp. pl. d. II. p. 956. Hott fl. austr. 2 p. 294. Fl. ic. ex. p. 227. Alsch. fl. jacq. p. 156. Pet. bot. vrb. n. 441.

G. glaberrimum foliis orbiculatis 5-7-partitis, segmentis linearibus tri-angulatis, pedunculis
bifloris filiformibus foliis longioribus, petalis ovatis emarginatis calycem longius
austum aequantibus ^{oblongis} ^{basi barbatis}, carpellis glabris latis, caudis pilosis, seminibus
abundantibus.

Icon. Fl. ic. fl. germ. v. f. 4875. Coll. et Germ. fl. par. atl. pl. VI. f. B. ^{l. 166. angl.} bot. t. 259. Cav.
monod. typ. t. 82. f. 1. Vaill. bot. par. p. 15. f. 4. — ^{Allyr. Karuvar.}

Hab. in agris, saxosis, demum tibiis dolosis. Fl. vrb. in aequalum. Fl. villosi-vrb. 2

[illegible]

L. annuum, foliis linearibus lanceolatis, utrinque scabris, axillis unguiculatis, marginibus serratis, nervis floribusque apice corymbosis, spiciferis, pediculis praecipue brevibus, vix, singulis lanceolatis attenuatis, vix glandulosis, vix capitulo subrotundo acuto saepe longioribus.

Syn. *Cotyledonum strictum* ^{var. a.} Phil. ic. fl. germ. VI. p. 62. — Non. Phil. ic. fl. germ. VI. p. 5170. a. J. Bauh. hist. pl. III. p. 455. L'Her. icon. t. 411. f. 2.

Var. β . *nicotianum*: florum corymbosis ^{axillis} spiciferis. — Syn. *L. inaequalis* Pers. del. prog. p. 53. ^{Phil. ic. fl. germ. VI. p. 5170. b.}
Cotyledonum strictum β . c. Phil. ic. fl. germ. VI. p. 62. — Non. Phil. ic. fl. germ. VI. p. 5170. c. Sibth. et Sm. fl. gr. IV. t. 304.

Hab. locis riuiculisque spicis circa Rhodum, Spalato, Ragusa, et in insulis Sansego, Cherso, Pago, Brijuni, Lesina. Fl. aegypt. 3.

Ordo XCIX Oxalidaceae

St. prod. 1798. art. 1. p. 689.

Floris hermaphroditis regularibus. Calyx quinquepartitus persistens aestivatione imbricatus. Corolla pentapetala hypogaea, petalis ~~longis~~ ^{imbricatis} et immixtis, aestivatione contorta. Stamina decem hypogaea basi connata, quinque petalis opposita breviora, aestivatione libera, longius longis immixtis, longitudiniter deliscentibus. Pistillum gemmae liliaceae ex omni ~~partibus~~ ^{partibus} connatis bi-pluriovulatis quinquevulvatis, placentalibus, ovulis pendulis, stylis ~~longis~~ ^{longis} quinque, stygmatis capitulis filiformibus. Capsula quinquevulvata, loculis pluribus dorso longitudiniter deliscentibus, seminibus pendulis, ~~multis~~ ^{multis} embryone rectiflexo in axi albuminis tenui, radicata ad hilum vix.

Herba acida, foliis ovatisque pinnatis cuneatis, petiolatis, albis, triplicatis, stipulis membranaceis nullis, pedunculis axillariis pluribus aut indivisis unis.

Oxalis L.

Syn. plant. ed. II. p. 160.

Char. Idem ac Oxidif.

Oxalis corniculata L. sp. pl. ed. II. p. 623. Hist. fl. aegypt. I. p. 553. Phil. fl. ex. p. 781. Alsch. fl. jud. p. 102. Pet. bot. weig. n. 676. ^{excl. 190. Reg.}

O. caulis ^{ramulosis} prostratis basi radicantibus, foliis basi irregularibus, pedunculatis subulonis filis subaequalibus, pediculis praecipue vix

Syn. O. villosa Bieb. fl. tur. cauc. I. p. 555. Phil. fl. ex. p. 781. — Phil. ic. fl. germ. V. f. 4896. 4897. Sibth. et Sm. fl. gr. V. t. 451. Bow. engl. bot. t. 1726. Lag. oxal. t. 5. Char. hist. pl. VI. p. 249 f. 1.

Hab. vulgaris in vix horti et cultis vix Salomonis litoribus et insularibus. Fl. flori. H. Magi. Rhod. 3.

Orb Cl. *Meliaceae*

Teil. in dem. der Mus. III. p. 436.

[illegible]

Melin L.

Gen. pl. 2. II. p. 148.

Char. Nov. ac. ording. — Class. Secundaria. Rd. Monogynia. Syst. lat. —
 excl. var. p. 26

Meli Azidavach L. 1p. pl. 2. II. 558. ^{excl. var. β.} *Cavend. 2p. monod. 7. p. 365.*

20. pro N. 145. not. 1. p. 62. Alah. fl. jad. p. 155.

M. polij biznansij laevitij, flakij arakij, cyprij subnity, panitij
aylij.

Icon. Curv. t. c. t. 207. Lam. enc. bot. pl. 372. Commel. herb. 1. t. 70. Dodon.
pempt. p. 848. — Myr. Persoon o³ jede varste.

Hab. ~~fertile~~ sponte ad fructus circa p^ro, et p^ro Catus. Koy 64.
columnae thimuli ^{pallide} ~~longissimi~~,
civis ^{experiment} ~~fragantif~~. H. acuta. §

Obs. Pulpa fructus dulcissimae, ex cl. Persoon vincta, niti regit^{experimento} et immola.

Ordo CII. Acetivores

DC. théor. élém. d. II. p. 246.

Hyssopus officinalis, and above unisexually, regular. Calyx lobes 4-8-partite, leaving
 aestivation imbricatif. Corollae partite, leaving calyx narrow acuminate at apex
 margin of the hypogynous insert, like, very small. Stamina octo filamentis
 liberis filis medi insertis, antheris bilobatis, dorso apice insertis, longioribus
 latis reliquantibus. Pistillum brevius liberis filis insertis, sessile, bibe, bilobis, ex
 omni breviter breviter receptum prope staminem infra adnatis, placenta
 centralibus, ovulis adscendentibus, stylis brevibus rhynchophyllis liberis brevibus. Fructus
 in capsule bina monosperma indehiscentia exsert in aliam oblongam partem a staminem
 persistentem breviter breviter, breviter adscendentibus exaltantibus, indehiscentibus.

Velleboth
Hab. in sylvestribus montibus Romaniae, Bulgariae, Polyst, et circa Axiarum. *Velleboth, lign.*
H. virens. $\frac{1}{2}$

Ufus. Omnia ac praesentia.

Acer campestre L. sp. pl. ed. II. p. 1497. Dif. 189. Delm. p. 23. Hist. fl. austr. I. p. 480. Pl.
fl. ex. p. 769. Alsch. fl. jedr. p. 90. Pelt. bot. weng. n. 3.

A. plis bps cordatis palmatis quinquefoliis laciniis oblongis obtusis integris lobatis, compatis
erectis regulariter palmatis illis, alij fructuum dimidiatis.

Don. fl. h. II. fl. germ. v. f. 4825. Don. angl. bot. t. 304. Ced. fl. Don. t. 1288. Duh.
trait. de arb. t. 16. 10. f. 7. — Myr. Maklen. Makljen.

Hab. in sylvestribus montibus Velleboth, nec non circa Norum, Zann, Lign, Lign, Lign, Lign et
alibi. H. virens. $\frac{1}{2}$.

Ufus. Lignum ipsum usibus praestit ac in praesentibus. Cortex desuper virens et la-
nem inspicit color lateris, gorgonium lignum corolla carnea.

Acer monspeliacum L. sp. pl. ed. II. p. 1497. Dif. 189. Delm. p. 23. Hist. fl. austr. I.
p. 481. Pl. fl. ex. p. 769. Alsch. fl. jedr. p. 90. Pelt. bot. weng. n. 4.

A. plis bps cordatis palmatis quinquefoliis laciniis oblongis integris, ^{hinc} ~~compatis~~ ^{aut pubescentibus}
erectis regulariter palmatis illis, alij fructuum erectis

Don. fl. h. II. fl. germ. v. f. 4826. Duh. trait. de arb. t. 16. 10. f. 8. Platan. aln.
t. 251 f. 3. J. Bauh. hist. pl. t. p. 168. — Myr. Klen.

Hab. in sylvestribus circa Zann, Norja, Lign, Polyst et alibi. H. buten-
virens. $\frac{1}{2}$.

Ufus. Lignum ipsum usibus adhibetur ac praesentibus. Cortex desuper ^{virens} ~~virens~~
et lacnem inspicit color ruber. aurantiaco.

Bbs. Variat lris plerumque obtusis, ~~lris~~ ^{et} aut, et hoc forma posterior exhi-
et illyricum Jany. fl. $\frac{1}{2}$ utrumque videtur in videri iconibus Fischerianis.

Ordo III. Euphorbiaceae

Don. gen. pl. p. 384 (nom. em. in fl. Br. Flind. vry. 2. p. 555)

Herz monili aut d'op, virens nuda et cartilaginea comuni inclusa, virens ob calycem
virens et corolla instructa. Calyx liber 4-5-fidus asperitudo instructa multigen.
Corolla pentagloba hypogaea 5-fidus nuda. Herz monili, staminibus corollae
sunt sub ovum rudimentis instructis, alibi oblongis longitudinaliter debi-
tibus. Herz femineae geminae libro ex ovum tribus vix debis receptivis.
prospice clauoni adnexis ^{in tubulosis} ~~in tubulosis~~, latis uni-bivalvis, placenta centris
libis, ovulis pendulis, et stylis tribus, vix debis integris bps 1/2. Capsula tri-bis

Icon. Phleb. ic. fl. grum. V. f. 4750. Sibth. et Sm. fl. gr. V. t. 461. Dodon. pempt. p. 377. Clus. hist. pl. VI. p. 187 f. 149.

Var. β . canescens: caulibus stipulis villosis canescentibus & sylv. E. canescens L. sp. pl. ed. II. p. 652. Jacq. collect. suppl. p. 115. Hort fl. austr. 2. p. 555. Phleb. fl. exc. p. 755. Pet. bot. weing. n. 382. Icon. Phleb. ic. fl. grum. V. f. 4751. Jacq. l. c. t. 2. f. 3. Cav. icon. et descr. pl. t. 1. t. 63.

Hab. in arvis et iugis circa Lann, Schuier, Popul, et in insulis Vexin, Lepine et Lina, vixitque propinquus ac speciosus. Glandulae florum rubrae appendicibus albidis aut canis. Fl. ab aestate in autumnum. $\odot$

Pls. Stipulae lineares vixitque integre aut pariter. Pubi modo vixitque, modo florum caulem, modo caulem et folia, interdum caulem folia et corymbos induit. Hinc vixitque, vixitque ellipso-oblonga, oblonga lenticulae immixta, ut dicitur dicitque glandulae acuminatae. Interdum hinc vixitque vixitque, et hinc vixitque in S. maculata L. den hinc in Ang. den p. 35. Euphorbia hinc L. sp. pl. ed. II. p. 652. Vif. hinc. Delm. p. 35. Hort fl. austr. 2. p. 556.

Phleb. fl. exc. p. 755. Alch. fl. jadr. p. 220. Pet. bot. weing. n. 393. E. caulem pariter, folia pariter oblongis bry. semicordatis subvixitque integris, florum axillariis solitariis, corymbis semivixitque laevibus

Icon. Phleb. ic. fl. grum. V. f. 4753. Lav. engl. bot. t. 2002. Clus. hist. pl. VI. p. 187 fig. 1. Maab. in Diss. p. 1322.

Hab. in littoribus, tum continenter tum insalubrem propinquus. Glandulae vixitque appendicibus albidis. Fl. ab aestate in autumnum. $\odot$

Pls. Stipulae vixitque integre ut pariter. Corymbae angule carinatae.

Sect. II. Galorhoecy Haw. syn. rusc. p. 157. Glandulae hinc exappendiculatae bry. vixitque in vixitque. Interdum glandulae vixitque vixitque. Phleb. ic. fl. grum. V. f. 4754. Lav. engl. bot. t. 2002. Clus. hist. pl. VI. p. 187 fig. 1. Maab. in Diss. p. 1322.

Euphorbia helioscopia L. sp. pl. ed. II. p. 658. Hort fl. austr. 2. p. 563. Phleb. fl. exc. p. 755. Alch. fl. jadr. p. 220. Pet. bot. weing. n. 390.

E. caulem pariter, folia pariter ovatis serratis, vixitque emarginatis, umbellae quinquefide vixitque vixitque, interdum vixitque vixitque, subvixitque, corymbis hinc, semivixitque vixitque.

Icon. Phleb. ic. fl. grum. V. f. 4754. Lav. engl. bot. t. 883. Hed. fl. den. t. 725. Sachs. hist. stng. p. 311. Hb. in cultis ubique vixitque. Fl. a hinc in D. hinc.

Uly. Hic et quoniam plurimas vixitque acri vixitque, vixitque ad vixitque vixitque. Hic hinc vixitque vixitque.

Aug. 10. var. 1. t. 86. - *Barb. i. c.* ~~197~~ 197. — *Myr. Binkneyi*. *Peraga*, *H. a. Marki*
Lamm. h. p. f. ed. m. t. f. t.
in meju. H. l. t. i. t.

Obi. Plante culte plurilobée, obtusif, ^{et} sans non éminemment prostrée. Vénix
ramelle 1-2-3-4-5-6-7-8-9-10-11-12-13-14-15-16-17-18-19-20-21-22-23-24-25-26-27-28-29-30-31-32-33-34-35-36-37-38-39-40-41-42-43-44-45-46-47-48-49-50-51-52-53-54-55-56-57-58-59-60-61-62-63-64-65-66-67-68-69-70-71-72-73-74-75-76-77-78-79-80-81-82-83-84-85-86-87-88-89-90-91-92-93-94-95-96-97-98-99-100-101-102-103-104-105-106-107-108-109-110-111-112-113-114-115-116-117-118-119-120-121-122-123-124-125-126-127-128-129-130-131-132-133-134-135-136-137-138-139-140-141-142-143-144-145-146-147-148-149-150-151-152-153-154-155-156-157-158-159-160-161-162-163-164-165-166-167-168-169-170-171-172-173-174-175-176-177-178-179-180-181-182-183-184-185-186-187-188-189-190-191-192-193-194-195-196-197-198-199-200-201-202-203-204-205-206-207-208-209-210-211-212-213-214-215-216-217-218-219-220-221-222-223-224-225-226-227-228-229-230-231-232-233-234-235-236-237-238-239-240-241-242-243-244-245-246-247-248-249-250-251-252-253-254-255-256-257-258-259-260-261-262-263-264-265-266-267-268-269-270-271-272-273-274-275-276-277-278-279-280-281-282-283-284-285-286-287-288-289-290-291-292-293-294-295-296-297-298-299-300-301-302-303-304-305-306-307-308-309-310-311-312-313-314-315-316-317-318-319-320-321-322-323-324-325-326-327-328-329-330-331-332-333-334-335-336-337-338-339-340-341-342-343-344-345-346-347-348-349-350-351-352-353-354-355-356-357-358-359-360-361-362-363-364-365-366-367-368-369-370-371-372-373-374-375-376-377-378-379-380-381-382-383-384-385-386-387-388-389-390-391-392-393-394-395-396-397-398-399-400-401-402-403-404-405-406-407-408-409-410-411-412-413-414-415-416-417-418-419-420-421-422-423-424-425-426-427-428-429-430-431-432-433-434-435-436-437-438-439-440-441-442-443-444-445-446-447-448-449-450-451-452-453-454-455-456-457-458-459-460-461-462-463-464-465-466-467-468-469-470-471-472-473-474-475-476-477-478-479-480-481-482-483-484-485-486-487-488-489-490-491-492-493-494-495-496-497-498-499-500-501-502-503-504-505-506-507-508-509-510-511-512-513-514-515-516-517-518-519-520-521-522-523-524-525-526-527-528-529-530-531-532-533-534-535-536-537-538-539-540-541-542-543-544-545-546-547-548-549-550-551-552-553-554-555-556-557-558-559-560-561-562-563-564-565-566-567-568-569-570-571-572-573-574-575-576-577-578-579-580-581-582-583-584-585-586-587-588-589-590-591-592-593-594-595-596-597-598-599-600-601-602-603-604-605-606-607-608-609-610-611-612-613-614-615-616-617-618-619-620-621-622-623-624-625-626-627-628-629-630-631-632-633-634-635-636-637-638-639-640-641-642-643-644-645-646-647-648-649-650-651-652-653-654-655-656-657-658-659-660-661-662-663-664-665-666-667-668-669-670-671-672-673-674-675-676-677-678-679-680-681-682-683-684-685-686-687-688-689-690-691-692-693-694-695-696-697-698-699-700-701-702-703-704-705-706-707-708-709-710-711-712-713-714-715-716-717-718-719-720-721-722-723-724-725-726-727-728-729-730-731-732-733-734-735-736-737-738-739-740-741-742-743-744-745-746-747-748-749-750-751-752-753-754-755-756-757-758-759-760-761-762-763-764-765-766-767-768-769-770-771-772-773-774-775-776-777-778-779-780-781-782-783-784-785-786-787-788-789-790-791-792-793-794-795-796-797-798-799-800-801-802-803-804-805-806-807-808-809-810-811-812-813-814-815-816-817-818-819-820-821-822-823-824-825-826-827-828-829-830-831-832-833-834-835-836-837-838-839-840-841-842-843-844-845-846-847-848-849-850-851-852-853-854-855-856-857-858-859-860-861-862-863-864-865-866-867-868-869-870-871-872-873-874-875-876-877-878-879-880-881-882-883-884-885-886-887-888-889-890-891-892-893-894-895-896-897-898-899-900-901-902-903-904-905-906-907-908-909-910-911-912-913-914-915-916-917-918-919-920-921-922-923-924-925-926-927-928-929-930-931-932-933-934-935-936-937-938-939-940-941-942-943-944-945-946-947-948-949-950-951-952-953-954-955-956-957-958-959-960-961-962-963-964-965-966-967-968-969-970-971-972-973-974-975-976-977-978-979-980-981-982-983-984-985-986-987-988-989-990-991-992-993-994-995-996-997-998-999-1000-1001-1002-1003-1004-1005-1006-1007-1008-1009-1010-1011-1012-1013-1014-1015-1016-1017-1018-1019-1020-1021-1022-1023-1024-1025-1026-1027-1028-1029-1030-1031-1032-10

8 *Euphorbia dendroidea* L. p. pl. ed. II p. 662. *Lehb. fl. exc.* p. 758. *Pest. bot. univ.* n. 384.
var. glabra

8. ~~caerulea~~ ^{Pichanthus} fruticosa ramis dichotomis, folijs lanceolatis oblongis in petiolum attenuatis
integris, florum umbellulis tri- octopletis, radijs bi-tripletis, ^{Pichanthus n.} corollae longe oblongae,
^{transversae} apice laevibus.

Sp. E. *divicata* Jacq. ic. rar. t. p. 9. — Pan. *Plhb.* ic. fl. germ. v. f. 4772 opt. Nth.
et Sm. fl. gr. v. t. 470. Jacq. ic. rar. t. 58.

Hab. in saxo et rupibus maritimis circa Ragusa. Fl. lutei. Fl. jam a
 Romanis in Aprilis.

9 *Euphorbia pilosa* L. p. pl. 2. 11. p. 659. (Chab. fl. ex. p. 258. (excl. syn. Portenridt.)

8. *perennij, pubescent, folij oblongo-lanceolatis* h^{ij}. *costis ampliusculis* apice
rematis, *florum umbellae quinquedecim, radij trifidj subterminales, involucralis*
rhombico-ovatis *auris*, *capsula uncinata* *hirsuta*, *seminibus laevigatis*.

~~Syn. E. fictiva Portenuth. en fl. dalm. p. 107. E. Illyria Lam. enc. d. de Pad. 2. p. 432.~~

E. puberulum Desf. fl. atl. i. p. 386 (non Wahl) Nichb. fl. exo. p. 258 — *non Wahl.*
fl. crit. cent. II. p. 269 et ic. germ. v. p. 4169 et 4170 d. 3. *fl. pub. d. 3. p. 15*
fl. pub. d. 3. p. 15

Ab. ad paludes maritimas et rivulos circa Stagno, ubi legit Neumayer. 7.
~~nam legit Potermodon in summis jugis montis Bithoni.~~
luti. H. Myr. Junis. 4.

Obs. Huc plant. cuiusdam succent. *E. plurimum* Potensill. ex pl. Dalton p. 15 t. 3,
et Hart p. 49 p. 20 p. 566
Jus tamen ex iure et ratione, nec non ex analogia plant. in hibern. Potensill.
genu. scilicet hanc pariem d'fuit folij repensib. squar. pinnatis subsessib.
auris, ~~hinc tamen~~ ^{hinc tamen} plurim. annulatis, unde nomen, ~~et~~ ^{et} folij glandulis imbric.
emarginatis nec integris ut in *E. filif.*: Plank. et hinc saxatile subalgis.
Plank. ides dubit. nec ad ull. post Potensillagium inveni.

Sect. III Epula Hb. l. c. p. 168.

Platichthys grandis *exagynulata* *lunata* aut *biorny*. *Colydium*
lunum. *exagynulata*

10 *Euphorbia cyathophora* L. 1^o fl. v. p. 651. W. 1^o fl. v. p. 35. Hist. fl. v. p. 2. p. 565. Schb.

fl. ex. 1862.
E. flg linearly lanceolatis angustioribus, floribus umbellatis multifloris, subsessilibus
costis-incurvatis, glandulis brevibus, capsulis parvis, immixtis, laevibus.

* * * Biennes et annuaire

18 *Euphorbia Lathyris* L. p. pl. ed. II. p. 655. Hort. fl. austr. 2. p. 558. Rhod. fl. exc. p. 860.
Alsch. fl. ped. p. 220.

[illegible]

Icon. Rhb. ic fl. germ. v. f. 4283 opt. - *Nov. engl. bot. t. 2235. Bull. herb. franc. t. 403.*
Engl. Indist. Poist.

fl. green. V. p. 1105. — Myr. *Luchiniera vella*. Smilges. Irolit. Poist. Cist.

Cisto. verruc.
Hab. in ~~humiditate~~ et in cultis
~~humiditate~~ are Spitz-pape Schiew, Zagued prop Moschi, Bezato
prop Ragusa et in insula Virginia, Cherso. Fl. luteo. H. var. ♂.
est plant. diem ex ipso possum et dei guttorem

Ufus. Semina drachma virgute solent: oleum ex ipsorum pressum et sibi guttatum
6-12 exhibitum anthelminticum optatum, et olei Crotonis hylum
per succedaneum.

Euphorbia alypsa L. p. pl. ed. II. p. 657. Phil. Fl. ex. p. 760. Deshm. Fl. Ind. v. p. 80

[illegible]

Syn. E. juncea Hall fl. austr. 2. p. 562. Rhiz. fl. exerg. 2.60. quoad loc. nat. non dicit et
excl. ab omni ~~descriptio~~ ~~et synonymis~~ — Non Rhiz. ic fl. germ. v. f. 486. opt.
Sibth. et Sm. fl. gr. v. t. 462. ~~opt. Rhiz. exerg. 2.60.~~
Hall et Robinson, Poisson, et J. Stephens

Silbth. et Sm. fl. gr. v. t. 462. *Chrysom. canalic.*
Hab. ad muros agrorum et in agris aggeribus ad Bottrichella, Poissan, et S. Stephens
urae
frons apicalium. H. fusticuli. H. acuta. ☉
Nix. Males en. ob. ing. aug. N. p. 322.

Euphorbia *palustris* M. Maly en fl. ing. a. 1832.
in *Quintessence* All. fl. p. 257.
in *Quintessence* All. fl. p. 257. *anthelm.* *schol.*

[illegible]

gularis, flandis, hirsutus, capitis, ...
 syn. E. terracina Portm. in Holt fl. austr. 2. p. 558 nec alior. ex herb. Portensis!
 Hab. in ... inter spicas et Alniflo, nec non ...
 ...

[illegible]

[illegible]

8. filij oblongi lanceolati, inflexi limbo-
 indurati, per aperturam angustam, florum
 corollam tri-petalem, radiis
 longis, induratis ovatis ellipticis mucronatis,
 glandulis limbo-
 affixis laevibus, immixtis transversis sulcatis

Syn-E. acuminata Lam. enc. bot. d. n. Pad. II. p. 424. *E. oblongum* Lott. notk. p. 76 et
fl. gall. d. II. t. 1. p. 359. *E. -* *Stem* *fib.* ii. fl. germ. v. p. 4255 et 4276. Lott.
fl. gall. tab. 29. Ray. fl. austr. II. tab. 121. Boiss. fl. sic. t. 13 f. 37.
1841

Euphorbia exigua L. 1^a fl. 2^a fl. 654. Herb. fl. austr. 2^a fl. 557. Poth. fl. ex. p. 759.
Poth. bot. wenz. n. 386.

E. filix linearibus vel linearibus - cuneatis, acutis, vix, non floribus umbellae tri-pedunculatis, vix bifidis, subternatis, involucribus involucribus et pappi subternatis linearibus acutis, glandulis setaceo - bracteatis, capsula laevi, seminibus albo - subulatis.

Var. α . *acuta*: foliis ovatis, acutis, ~~longioribus~~ — Icon. Rhod. 16. fl. germ. v. p. 4777.
Oed. fl. Dan. t. 592. Dalech. horti 1656. p. 2.
emarginatis δ +

Var. β . *retusa* = *Phil. Grayii* ^{emarginatus} *obtusif. truncat.* — Syn. *P. retusa* Cav. ic. et descr. pl.
 1. f. 21. — *E. diffusa* Jacq. miscell. 2. p. 311. Herb. pl. austr. 2. f. 558 — Non. *Phil. ic.*
Regerm. - f. 1178 Cav. l. c. t. 34. f. 3. Jacq. ic. rar. 1. t. 88.

Var. *y. heterophylla*: folij inprimis obliq., recurvis acutis. — Syn. *E. corymb. y. L. p. gl.*
D. W. p. 654. — Bern. Magnol. bot. monog. p. 258.

Hab. in agris, laevis cultis et herbis etiam collibus totius Dalmatiae. H. lutea. H. a. var.
in artemisium. ☉

21 *Euphorbia Popley* L. f. pl. 2. II. p. 653. Vif. 187. Helm. p. 35. Holt fl. 20. p. 2. p. 556.
 f. alb. fl. ex. p. 759. Alch. fl. jed. p. 220. Pott. bot. weing. n. 392.

8. plant. perfoliat. obovato-subcordatis, floribus umbellae ligatis, radiis brevibus, tubulosis, involucris thymalis, ovibus ovatis, glandulis sessilibus, capsula coriacea, laevigata, baccata, bidentata, seminis ovibus perfoliatis, fere longiusculis, baccatis.

Don. Tab. ic. fl. grm. v. f. 423. Sow. engl. bot. t. 29. Bull. herb. de la Fr. t. 79. Des fl. des t. 110.

Don. Tab. ic. fl. grm. v. f. 423. Sow. engl. bot. t. 29. Bull. herb. de la Fr. t. 79. Des fl. des t. 110.

Don. Tab. ic. fl. grm. v. f. 423. Sow. engl. bot. t. 29. Bull. herb. de la Fr. t. 79. Des fl. des t. 110.

Buxus Tourn.

Instit. rei herb. p. 578. t. 345.

[illegible]

Buxus sempervirens L. f. pl. ed. II. p. 1394. Vir. 189. Sem. p. 28. Hort. fl. agra 2. p. 613.
 Rhob. fl. ex. p. 765. Alth. fl. jad. p. 224.
lucida, ...

B. filix ovato-elliptica, petiolis ~~in~~ pilis, antheris ovato-lanceolatis obtusis. ^{lucida,}
 Jon. Rechb. ic. fl. germ. v. f. 4808. Th. F. Nees gen. pl. III. n. 16. Bull. herb. de la Fr.
 t. 263. Low. engl. bot. t. 1341. — Rhyn. Zelenika. Bux. Schimsch. Merija ^{pitoma.}

Hab. in havis, sponte in insulis et propinque in riv. Arabe. Fl. luteo. H. ven. $\frac{1}{2}$

Sect. I. *Alaternus*. ~~Tarn. inst. p. 1. 566.~~

~~Plang polygonum dicitur ruscari 4-5 foliis 4-5 andis, stylis bi-hispidis. Stems rufis profundis, angulis exiguis et ruscariis.~~

1. *Phanery Alaternus* L. p. pl. ed. II. p. 281. Vif. 1879. dat. p. 45. Hort. fl. austr. 1. p. 305. f. ed. fl. ex. p. 488. Alch. fl. jedr. p. 55.

Ph. arborescens erectus, ^{marginis serrulatis} foliis ^{lanceolatis} ovatis ellipticis, remote serrulatis, cuneatis, ^{marginis serrulatis} basi ^{marginis serrulatis} connatis.

Don. Duk. tract. des arb. 1. p. 40. Cher. hift. 1879. 1. p. 50. J. Bauh. hift. fl. 1. p. 542.

— Mlyr. Konarka

Hab. in collibus sylvestribus circa Schin, et in viderum Vergata et Paga. Fl. virideli.

Pl. var. $\frac{1}{2}$.

Sect. II. *Cervipina* ~~Dillen. var. gen. p. 1. 8.~~

~~Plang polygonum dicitur pinnatis tetrapetala, stylis bi-hispidis. Semina rufis profundis, angulis exiguis et ruscariis.~~

2. *Phanery catharticus* L. p. pl. ed. II. p. 279. Hort. fl. austr. 1. p. 302. f. ed. fl. ex. p. 487.

Ph. arborescens erectus, ^{marginis serrulatis} foliis ^{marginis serrulatis} ovatis ellipticis, subterminalibus serratis, ^{marginis serrulatis} basi ^{marginis serrulatis} connatis.

Don. Low. engl. bot. t. 1629. Oed. fl. den. t. 850. Lam. enc. bot. fl. t. 128 f. 2. Des. genest. p. 156.

Method. in Drie. p. 175.

Hab. in dumetis circa Sign et Roqua. ~~ad hunc agri Caputini, unde nuncit Roumayer.~~ Fl. virideli. Baccae maturo nigrae. Pl. var. $\frac{1}{2}$.

Ufus. Baccarum recentium 10-20 ingestae, vel candorem recens expressae a frigido decoctione ad quatuor, vel decoctionem ex 20-30 brevibus, vel syrupum ad usum ad 2 uncias si dragma et peremptoriam etiam diuresis potest. Baccarum ignis immaturum si cum alamine et aqua stillata colorem lacteam, maturumque lactem videri, maturumque et caducum peremptoriam praebent. Symplicium totum.

2. *Phanery* ~~Alaternus~~ L. p. pl. ed. II. p. 281. Hort. fl. austr. 1. p. 303. f. ed. fl. ex. p. 488.

Ph. arborescens erectus, ^{marginis serrulatis} foliis ^{marginis serrulatis} ovatis ellipticis, subterminalibus serratis, ^{marginis serrulatis} basi ^{marginis serrulatis} connatis.

NB. Syn. Ph. vaxanthus Vif. 1879. dat. p. 45. Alch. fl. jedr. 55 non f. — Don. Vif. fl. ed. t. XXXVII. A. fl. cum stipul. B. fl. masc. C. fl. fem. 2 bacc. Duk. tract. des arb. II. p. 51. J. Bauh. hift. fl. 1. lib. VI. p. 50 f. 1. 2. (ed. 3.) — Mlyr. Tarn. Tarn. exar. Grana. Mast. Bugiokka. Tarnovina.

Hab. in saxosis agris ^{marginis serrulatis} hinc ^{marginis serrulatis} Delmatia. Fl. virideli. Baccae nigrae. Fl. var. $\frac{1}{2}$.

Ufus. Tissem usum ^{marginis serrulatis} in ^{marginis serrulatis} murt, sed baccarum colorem praestantior praebent idcirco murt ipsum antipruritus.

Obs. Spines plerumq. ad aurantibus ~~seem~~ forming capitula the maximally immature
 & qua statim differt puberis brachy renger Canine una quod vixit et ultra
 breviter the quid ion Arden. mem. Dioneo p. 11 quod planta ex herb ex
^{Pellaea} in colibis veronensis obscure, see the infectum ibidem quid quid definitum.
the lineary Walton et Kit. pl. var. hug. 3. p. 283, uni huc quod ion Clay.
hug. pl. p. 111, folia herb ovato acuta, sed quod hanc ut certa longi.
non diffini requit.
Chemney spines L. 18. pl. ed. II. p. 280 et max. alt. p. 341. Hort. pl. cap. p. 12. Vol. pl. pl.
 11. exc. p. 458.

8 fl. ex. 458.
Rh. abovens ^{abovensis} var. alberr incunctus, folijs ellipticis basi rhomboidis emarginatis
^{puberulis}
ginebrae crassimulari ~~puberulis~~, calycibus ~~tubis~~ quadripartitis lobis, parvis,
lamina multo brevioribus.

Lamina multorum *multorum* *multorum*.
Non. Hall. hist. stirg. helv. II. p. 43. J. Banch. hist. pl. I. lib. V. p. 562.
Vallisbella ad Vella Pothlyngia

Buccae regina H.
Habitat in regione
Majo in aestate.
Flores brevipedunculatae, tubulosae, liliaceae, corollae lobis
Ranunculus Ranuncula

Ph. abnormis erectis ramis alternis immixtis, foliis ovatis bps obliquis connatis
 subintegris, ~~calycibus~~ ^{quadr-} ~~quinque~~ ^{lobis} lobis, petalis laniatis nud.
 lobis brevioribus.

Loc. enst. bot. p. 250. Sed. fl. den. p.

Urtica brevifolia
Syn. *Frangula vulgaris* Rehb. fl. ex. p. 488. — Don. Low. angl. bot. t. 250. Pad. fl. den. p. 1.
— Boiss. med. di. omer. tab. 15. Matth. hist. in Drac. p. 1334. — Meyer. Lexikon.

[illegible]

var β . *penicillata*: putrescent ascending and prostrate, fls. solitary or subsessile, 1/8-1/4 in. diam. - 6-8. Host fl. Aug. - 10. 304
 5-6 mm. — Syn. *ph. penicillata* L. mant. l. p. 49. *penicillata* L.

[illegible]

268. *Myrica Myrtina* (Schreb. fl. ex. p. 488) — Don. Cat. ex. p. 111. —
var. 2. fl. 181. *Myrica fl. europ. p. 10. X. Don. fl. v. p. 123. fl. carn. t. 6. 5.*
var. 2. fl. 181. *Myrica fl. europ. p. 10. X. Don. fl. v. p. 123. fl. carn. t. 6. 5.*

no. 2. f. 181. Poem fl. europ. Jacq. x. Sen fl. p. 67.
Hb. spec. præsertim in sylvestri videretur Vellut., forma minor in collibus
et raptis circa Schio, Vini, Salona, Ragusa, varietas in vespertino all-
Zonitella circa Spalato, ^{vini} Ragusa, et in monte d'Opere. H. alb. Sæpe viridis, dein
viride aut caeruleo-nigra. H. var. f.

Obs. Antherum ovarium qui de Ph. perula agens in ejusdem utroque
 ut utriusque notum hanc variabilitatem testantur. Hinc Antherodis ex
 Linneo Cavendishii Poenae Melfinis et Segalis ab alijs duntaxat monon et
 poggiani: quod p. d. ex ~~homo~~ Cavendishii Nervos et Bostonia, poggiani sunt
 Melfinis et nunc quod p. d. nunc poggiani Segalis et Poenae. Stylus triplo juxta
 Poenae, Cavendishii Nervos et Bostonia, indistinctus dicitur a Melfinis et p. d.
 batis, a Kothis ~~ab alijs~~ Vni Ph. perulorum a Ph. virentis repertus indistinctus in hoc b. d.
 p. d. in illo agnoscitur. Hinc clar. signatur utriusque non obrem habere com.
^{reperit} ut hunc p. d. ad speciem virentis. Quoad alia diffinitioni

Var. β . *sepium*: *padanensis* *frutescens* *inermis* *fabry*. — *Syn. N. sepium* *Thunb.*
fl. pers. ed. II. p. 282. *fl. rubra* *sepium* *Savi*. *fl. ital. I. p. 103.* *fl. agrostis* *Savi*
fl. pis. I. p. 455. *fl. oxy. 618.* — *Non.* *Savi* *fl. ital. I. tab. 35. p. 2.* *Sow. conf. bot.*
t. 2650. *Pollen* *fl. uron. II. tab. 2. p. 4.*

Hub. ad rep. et demet. circa Zam, Galat, in insula Loxia, in montibus
Dithori et Sabas: var. in sylvestribus ad Thymali super Libanum. H. pulcherr.
mici. H. Major, Taurus etc.

Now collum lay. fl. anfr. 2. 58. Hoit fl. anfr. 2. p. 22. *fl. ex. 620.*

1. aculeis longis - 2. aculeis ad medium, foliis ellipticis ovatis pubescentibus
 utrinque glandulosis, stipulis 2, aculeis ovatis ~~longis~~
 4. aculeis longis, foliis ^{ind.} ellipticis, laciniis calycis albis, bracteis punctatis,
 calyce punctato obliquo aut ovato.

Syn. *Q. dumetorum* Deso. Journ. bot. 1815 p. 115. ~~f~~. From Paq. p. 1. 197.
 Now conf. bot. t. 2579.

Hub. in demselben 2 Minuten nach Beginn. H. f. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840. 841. 842. 84

~~*Q. macrophylla* Hart. fl. angl. p. 192. L. minor. fl. alb. p. 245 excl. in fr. Dan. Hart. fl. angl. p. 26. fl. ex. p. 605. Alb. fl. p. 113. Hart. bot. voy. n. 778.?~~

N. praeumbra, semis flegmatizans, aurescens valde congesto - Nodulis denticatis, parvis
orbibus et ovatis-lanceolatis

Rosa rugosissima L. fr. pl. d. t. p. 404. Felt. fr. ex. y. 685. Alut. fr. i. j. 113. P. b.
H. wuz. u. 781.

[illegible]

syn. *P. balnearia* syn. *P. II. p. 49. P. abnormis* *W. fl. h. t. p. 1. p. 4. — R.*
W. fl. h. t. p. VI. Dill. h. elth. II. t. 246. p. 218.

North. ch. in fl. gr. V. t. 585. Dio. l. c. t. VI. Dill. n. etn. II. r. c. 116.
Nas. p. glutinosa: per. anther. calyx. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828

Hob. in density at reg. by long distance from it and the total mass
it is ~~the~~ the insom. H. asph. 7.

Polio pumila L. fl. reg. p. 262. Botk. H. Tr. v. p. 220. Hist. fl. ang. p. 27. 23. fl. c.
H. c. p. 622 exal. p.

[illegible]

Syn. R. yellow a. pinnil D. H. R. pubescens & pubescens
IV. n. 3709. - *Ulm. Ray. fl. austr. 2. p. 1980. Vill. fl.*
No. 11. 22. *Chus. Hoff.*

Herb. in Mly. circa Zar, Kruin, Marne et Jyga. H. rose. alba. H. magis. 5.

Sect. II. Malay Journ. inst. p. t. 406.

Very umbellate; styles 4, connate, minimally exserted convergent divergency. Præputia
4, umbellately endocarpis cartilagineæ.

Syring vulgaris L. p. pl. II. p. 686. a. Diff. King. Schim. spec. p. 14. Hort fl. austr. 2. p. 13. Pluk.

fl. exl. p. 631

fl. ex. p. 631. *arctis, obliq. duplicata - semitip*
P. folij arctis obliquis, ovata, semit. petio. longioribz, fl. arctis, semit. petio.
subrotund - strobil. fructibz globosis, bafi umbellatis.
alt. Hook. p. 631. Sem. cont. 66. t. 120. Sid. H. den t. 1101.

Syn. *P. acerba* Bl. *prodr.* syst. nat. 2. p. 635. — *non*. Sw. *enfl.* bot. t. 179. *ped. fl.* den t. 1101.
Pickw. herb. t. 178. *Method.* in *Diep.* p. 239. — *Wlgr.* *Tsukuba*.

Hab. in collibus saxosis ad Vercelli, in monte Mosor, et in insula Veglia ad Capriam.
H. alb. eximie nuda, odor. H. Agrosti, Major. f.

Pl. alt. calyx rose, stem - H. April, May 4.
 Pl. variat foliis junioribus calycibus glaberrimis, et hoc est P. aerea H., nulli alii
 nota a specie diverg.

Sect-III. Aria H. prod. syst. nat. 2. p. 635

Flora nuncup. composita, glandulis mucosis, stylis 2-4, ^{strobili} fructu mucos. globosis enduratis
nuncupaco.

membranaceo.
Pyraea Esh. litt. IV. p. 40. Sitth. et m. p. gr. v. p. 63. Alsh. p. j. d. p. 110
biji uenuey uenuey
biji uenuey uenuey

Pipera and *Ech.* oct. IV. p. 40. *Silph.* chm. p. 97.
P. folij ellipticis dyalictis - seminis subglobosis, pericarpio multo longioribus rubris can-
 tonensis. 681. *Sobes* and *Croton* 1872 aug. p. 11 p. 46

Syn. Contaeus *ana* L. *W. p. ed. II p. 681.* *Solus ana* *Curtz* *W. p. p. II p. 46.*
ana *W. p. ed. II p. 681.* *Solus ana* *Curtz* *W. p. p. II p. 46.*

Arisaema *flexifol.* 628. *Arisaema* *nivea* Mill. fr. augr. 2 p.
fl. gr. v. t. 479 Low. engl. bot. t. 1858. Oct. fl. den. t. 302. Crumf. l.c. t. II p. 2
fl. alb. fr. asper. t.

Hub. in yelshy monum Vallibith et Brokov. fl. ann. p. 1. f. 1.
Ob. Ab hoc non differt lobus quereae Loddig. see S. ania var. quereae Griseb. spec. fl.
virens 1. p. 90 nisi ~~perantrum~~ jugis paniculatis (5-6) et ramulosis.

X - X. nat. rub. 2 p. 836.

Pyraea terminata Sect. IV. *Dominica* L. prod. 14th ed. 2 p. 636
 2-5 lvs. connatis. Nubby tw.

Flory corymbosi, pedunculis ramosis, stylis 2-5 bris connatis. Fructus testaceus
subterreus limbo calycis deciduo minutus, endocarpiis membranaceis.

Pygmy terminalis Ehrh. herb. vi p. 92. Alsch. fl. rar. p. 109
~~is not~~ ^{demum fl. rar.} leafy ^{subterminal} ^{acuminate} ^{long} ^{imbr.}

Pyrus terminalis Ehrh. bot. VI p. 92. Alida ^{remota yellowish} ~~pinnatifida~~ lobes acuminate, serrate, ~~imperfectly~~
P. filix late ovary ~~dissected~~ lobes bifurcate subovate [?] lobes acuminate, serrate, imperfectly
 patently ^{indistinct} ~~indistinct~~ l. - 10. st. 2-II p. 681. Host ft. augr. 2-p. 9. Lobes terminally bran-
 ching? [?] ~~l.~~ gen. and bot. t. 248. Pers.

Syn. Cortaeus terminelis L. *sp. pl.* 2. II p. 681. Host *fl. austr.* 2. p. 9. *Schiz. terminelis* (Gmel.)
fl. austr. pro II 45. *Phid. fl. ex.* 625. — *Don. low. engl. bot.* t. 298. *Pag.*
Phid. hisp. pl. I p. 10 f. 3. — *Myr. Ostrop.* *Di.*

fl. inf. v. t. 443. Sed. fl. den. t. 798. Claf. n. fl. t. 104.
Hab. in sylvis ad Kravica di. obkrova, in montibus Vellebshi dist. et Brkova. fl.
ab. fl. mayo t.

Tabl. veg. III. p. 349.

Flor hermaphroditi regulari. Calyx tubo exsertato est vix exsertato inter boni laevium an
 nulis perigynis limitatis vestitis, limbo quinquepartito laevij asperatione vix
 Corolla petalis quinque annatis perigynis inferij asperatione imbricata. Stamina
 cum petalis opposita ipsa vix minus aequalia aut plura antheris introij. Obcordatis
 Pappum ovarij induratum receptaculo demum exserto longius unguiculatij, ovum
 pendulo aut ascendente, stylis stygmatisque singulisque lobis. Fructus (^{ovariis})
) ex acheniis plerumque vix ut Dryadeis indehiscens, rarius adpendente
 aut intro, embryone exalbuminoso recto, radiculari ad hilum posita.
 Arbores aut frutices interdum acaules, folij parvis pinnatis aut digitatis rarij
 stipulatis serratis, floribusque corymbosis.

Classis analogica generum

Fructus acheniis exsertis W. N. 1.		Fructus acheniis Dryadeis	
#	Achenia plura	W. N. 1.	Achenia Fructus 1-4
1.	Achenia exserta	2	Achenia succulenta
2.	Receptaculum exsertum	3	Receptaculum canoso-succulentum
3.	Calyx bracteolatus	4	Calyx bracteolatus
4.	Receptaculum cylindricum continue		Dryas
	Genus		Receptaculum vix concavum
5.	Flor corollati	6.	Potentilla
			Flor apetalis W. N. 7.
6.	Calyx calyculatus		Calyx basi nudus
	Aromonia		Agrimonia
7.	Stamina 1-4		Stamina 20-30
	Achillea		Potentilla

Dryas L.

Char. Calyx patens ^{Fruct. ed. p. 135} 8-9 partibus. Corolla 8-9 partibus. Stamina et Pappi
 plurima achenia fere receptaculo convexo imposita, stylis subterminalibus glomeris
 persistenteribus caudatis. — Cl. Rosaceae Ord. Polygones Syst. 115

Dryas octopetala L. p. 11. Hall fl. europ. 29. 20. Pl. fr. europ. 39.

D. folij ovato-oblongis ovuliferis oblique serratis hys uteris, subrotundis, subuliferis, nuda, tomentosa.
 Non Low. engl. 64. t. 451. Ord. fl. den. t. 31. Lam. enc. 66. fl. 443. Clus. hys. fl. 1. p. 351. fig. cat.
 Hb. in officij ad Monte Sank montium Velleb. fl. ab. Engl. Ros. hys. fl. 1. p. 351. fl. 1. p. 351.
 aetate. t.

internus vel albidus

Hab. in montibus revivitatibz totius Dramatis. Fl. alb. fructus rubri. Haec Margi in Anglorum
off. Fructus pinguis edulis.
Tromia collina Vahl. bat. J. p. 26. Host fl. austr. 2, 3. h. f. l. ex. 597. Alud. fl. j. d.

p. 111.

Iron Bed. fl. Dec. 1. 1389. — Myr. Tijde.

Hb. in herbis alium humiliterum vix Boueygo, Leruy, Vezony et in impet. H.
alb. fructus ^{interius} rubri ut albid. rubi. H. Mayo, Nervo &

Trigonotis latior Ehrh. *bot.* 7. p. 23. *fl. austr.* 2. p. 33. *ph. fl. exc.* 9. 597.

7. gaudios, blanchy elongaty. filis caulis potidem perennantibus fortissim., capitis fructus reflex.

Syr. F. vesca β . *potensif* L. p. pl. 2. II. p. 709. — Non Low-engl. bot. t. 2197. Briton. p. 672. Lobel. ii. 1497. p. 697.

Herb. in monthly electricity, and holds firm to its peculiar nature a Pos. Magl. 76.
alt. fl. rubr. Flor. ab aestate in Septembrem 4.

— Potentilla L.

Gen. pl. d. II. p. 174.

Chas. Calyx plenus limbo 4-5-partito, calyc. 4-5-bracteolatis. Corolla 4-5-fidula. Stamina
et Pistilla plurima. Cochlearia fere receptacula univexa aut oviculi externi pilis im-
positi. — Clap. Repandria Ord. Myzogenia type sex. —

Seet. I. *Potentillastrum* Linn. in *DL. pndr.* 2. p. 571.

Receptaculum breviter pilosum. Atherium glabrum.

Potentilla hirta L. sp. pl. ed. It., 712. Rep. stirp. belm. sp. p. 44. Hort. fl. austr. 2, p. 37.
 Tab. fl. ex. p. 394. 7-5-fol. lobis

[illegible]

Var. a ^{in liq. superantibus strobil.} ~~pinis: strobis pilulis dentatis, ad apicem subulatis~~ ^{subulatis subulatis}
~~subulatis pilulis dentatis, ad apicem subulatis~~ ^{subulatis subulatis} - *Var. b* *Laurina* *Chlecht.* in *bot. mag.* VII p. 291 ex
P. reticulata Mill. *in* *bot. mag.* p. 38. *Publ. fl. exs.* 593. *P. reticulata* *Publ. fl. exs.* in *bot. mag.* p. 291
Lehm. mon. phtat. p. 90. *in* *herb. alt.* fl. ped. 2. p. 55 ~~in~~ *herb. alt.* - *Mon. mon.*
Lehm. l. c. tab. IX. alt. fl. ped. 1. p. 1. - *Mys. Mas. Dips.*
^{subulatis subulatis}

Var. β . *angustula*: ^{meant} ~~slight~~ ^{slightly} cuneato-conicoides, ^{thruout} ~~margin~~ ^{margin} ~~of~~ ^{of} ~~floridif.~~ ^{floridif.} — *lyn* P.
lecta *Lehm.* l. c. p. 88. *P. lecta* *Nob!* *Proc.* p. 595. — *Don.* *Lehm.* l. c. t. VII.
Lehm. l. c. p. 88. *P. lecta* *Nob!* *Proc.* p. 595. — *Don.* *Lehm.* l. c. t. VII.

Var. *p.* *levis* - *clavus*, *fl.* *longo-lanceolatis* a *br.* ad apicem *pinnae* *fid.*, *18* *rad.* *1* *emargin.*
n *fid.* - *syn.* *P. levis* *Waldst. et Kit. med.* *1801* *encl. 1* *rad.* *1* *emargin.* *1801* *encl. 1* *rad.* *1* *emargin.* *1801* *encl. 1* *rad.* *1* *emargin.*
Lehm. *mon. posth.* *t. VII.*

19. Var. *D. infusa*: ^{colleg. majoribus} *pusilla*, caule decumbente apiculato. — Syn. *P. hirta* β . Nees. mon. ² p. 56 (excl. syn. Nees.) Lehm. mon. bot. p. 88. *P. hirta* β . *infusa* Pers. 190 pl. 2, p. 54 (excl. syn. Ventr.).

(excl. syn. vult.)
Hab. in agros, praevalens collibus totius Dalmatiae: var. p. ad Grador et ad Metkings;
prope Daring. H. auri. fl. apul. 4.
H. d. p. a. vult. ad septem, pro

Sbs. Hark admodum variis numero phloemum diculis et pectus ad septem, per
jam utarum lignae, conundin totitudin, forma, et hacten usque ac
fuerunt, pectus ulgen ut parum ut decto longioribus. P. pilosum W. sp. pl. 2. p. 100
horum
quidquam differt folij rubro post folio clauso tamen arene rubrans, pectus

Polenilla inclinata. Willd. *Leist.* des pl de Grueph. 3. p. 567. *Rechb. fl. ex. p. 594.*
7-5. *foliolary*

P. canescens, caulibus adpendentibus erectis, folijs septembris primis nigrescentibus
rubris uncinatis, foliolis oblongo lanceolatis aut ovato lanceolatis, incisis - serratis
tota plantae basi cernitibus integris, calycis laciniis auratis, petalis calyce sublongioribus,
adhaerentibus tenuissimis arcuatis - rugulis marginatis pilis raris multo superantibus.

Syn. *P. lanerius* ^{pmv.} *Beg. fl. galic.* 1. p. 330. Hort fl. austr. 2. p. 37. Fuchs fl. ex. p. 394. Pr. v. 1.
Dag. fl. austr. IV. p. 43. — Non. Dag. f. c. t. 383. Will. t. c. t. XLV. (male).
colle *acid. # sinensis* (Kunze)

Hub. in collibus saxosis apertis circa Gradus, et in colle arif. # lunensis (Quen).

Polentilla argentea L. fr. pl. ed. II. p. 772. Hort. fl. austr. 2. p. 38. fl. ed. p. 594.

Pett-bot-wang. n. 725

[illegible]

Iron. Oct. fl. den. t. 865. Sw. engl. bot. t. 289. Carn. epit. pl. p. 260. Matth. in Drog. p. 1074.

Non. Ced. fl. den. t. 805. var. inf.
fl. : ... et Vrijesima. fl. lutea. fl. a. Major in asplenem. 4.

Polentilla reptans L. fl. p. 257 p. 714. Vif. 1 kg. Sem. spec. 46 Herb. R. n. 2 p. 42.
 Pub. fl. exc. p. 591. Alb. fl. p. 111. Pet. bot. wing n. 728
 5-3-petals indur.

P. puberula, caudicibus prostratis, floridiformibus, folijs quinatis lanceolatis, peltatis
oblongo-ovatis serratis, calycis lacinijs acutis, parvis, strobilij calyce rubiginosius, achenij
laeviusculis immixtis pilis receptaculis superimixtis.

Don. Low. engl bot. t. 862. Ged. fl. dan. t. 156. Salbat. hort. vom. v. t. 45. Blackw. t. 454. ^{heib.}

— Myr- Petoparst. Petolist. Pitparstiza -

Hab. ad vici ^{famine} ~~fructus~~, agrorum marginibus, pascuis, etc. *Silvestris*. *H. lutea*. *H. aspera* &c.

Obs. Aranea tenebrifera et araneus imperforatus punctulatus, sed nec granulatus nec scabrus, dorso vix carinatus
et omni margini excepto ut in precedenti.

Alchemilla vulgaris L. 18. pl. ed. II. p. 178. Hott fl. austr. p. 219. Nöb fl. ex. p. 609.

Q. foliis intralobis cordatis-obovatis 5-9 lobis, ^{longiusculis} ~~oblongis~~ nervis oblongis, caulibus 5-lobis,
caulibus decumbentibus, floribus cymosis terminalibus

Don. Low. engl. bot. t. 392. Sed. fl. dan. t. 993. Lam. enc. bot. pl. 58. f. 1. Blackw. herb. t. 32
M. et al. in Linn. p. 1236. — Wly. Zeydenjak.

Ver. S. montana: foliis subterre- hispatis. - Syn. A. alpine p. hybrid L. popl. d. 13.

A. hybrid Hoffm. fl. germ. 1. p. 99. *A. vulgaris* v. *hybrida* W. fl. pl. 1. p. 598. A.
montana W. en fl. h. herb. p. 170. *Prilag.*
Zürich. *Montenapoli*: var. in monte ~~Prilag.~~

montana W. & A. ^{Siberia}
 Hab. in parvis montibus ^{Siberia} et in Montenegro: var. in monte ~~Siberia~~
 fl. viridi: fl. aeth. 4. (ex var.)
 1. 219. Herb. fl. ex p. 610.

Alchemilla alpina L. p. pl. ed. II p. 129. Hort fl. aet. p. 1 p. 219. Pith. fl. ex. p. 610.
 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828.

A. folij radicalibus cordatis-obovatis, denticatis, plicatis ~~obtusis~~, sessilibus
stip. brevibus, caulibus sub-3-florescentibus, caulis erectis, floribus cymosis terminalibus.

B. *inimica* 386. Clav. h.p. pl. v. p. 108.

Shif. Hering, ^{unpub.} sub-3-fl. very, calving in 1911. ¹⁹¹¹ ¹⁹¹² ¹⁹¹³ ¹⁹¹⁴ ¹⁹¹⁵ ¹⁹¹⁶ ¹⁹¹⁷ ¹⁹¹⁸ ¹⁹¹⁹ ¹⁹²⁰ ¹⁹²¹ ¹⁹²² ¹⁹²³ ¹⁹²⁴ ¹⁹²⁵ ¹⁹²⁶ ¹⁹²⁷ ¹⁹²⁸ ¹⁹²⁹ ¹⁹³⁰ ¹⁹³¹ ¹⁹³² ¹⁹³³ ¹⁹³⁴ ¹⁹³⁵ ¹⁹³⁶ ¹⁹³⁷ ¹⁹³⁸ ¹⁹³⁹ ¹⁹⁴⁰ ¹⁹⁴¹ ¹⁹⁴² ¹⁹⁴³ ¹⁹⁴⁴ ¹⁹⁴⁵ ¹⁹⁴⁶ ¹⁹⁴⁷ ¹⁹⁴⁸ ¹⁹⁴⁹ ¹⁹⁵⁰ ¹⁹⁵¹ ¹⁹⁵² ¹⁹⁵³ ¹⁹⁵⁴ ¹⁹⁵⁵ ¹⁹⁵⁶ ¹⁹⁵⁷ ¹⁹⁵⁸ ¹⁹⁵⁹ ¹⁹⁶⁰ ¹⁹⁶¹ ¹⁹⁶² ¹⁹⁶³ ¹⁹⁶⁴ ¹⁹⁶⁵ ¹⁹⁶⁶ ¹⁹⁶⁷ ¹⁹⁶⁸ ¹⁹⁶⁹ ¹⁹⁷⁰ ¹⁹⁷¹ ¹⁹⁷² ¹⁹⁷³ ¹⁹⁷⁴ ¹⁹⁷⁵ ¹⁹⁷⁶ ¹⁹⁷⁷ ¹⁹⁷⁸ ¹⁹⁷⁹ ¹⁹⁸⁰ ¹⁹⁸¹ ¹⁹⁸² ¹⁹⁸³ ¹⁹⁸⁴ ¹⁹⁸⁵ ¹⁹⁸⁶ ¹⁹⁸⁷ ¹⁹⁸⁸ ¹⁹⁸⁹ ¹⁹⁹⁰ ¹⁹⁹¹ ¹⁹⁹² ¹⁹⁹³ ¹⁹⁹⁴ ¹⁹⁹⁵ ¹⁹⁹⁶ ¹⁹⁹⁷ ¹⁹⁹⁸ ¹⁹⁹⁹ ²⁰⁰⁰ ²⁰⁰¹ ²⁰⁰² ²⁰⁰³ ²⁰⁰⁴ ²⁰⁰⁵ ²⁰⁰⁶ ²⁰⁰⁷ ²⁰⁰⁸ ²⁰⁰⁹ ²⁰¹⁰ ²⁰¹¹ ²⁰¹² ²⁰¹³ ²⁰¹⁴ ²⁰¹⁵ ²⁰¹⁶ ²⁰¹⁷ ²⁰¹⁸ ²⁰¹⁹ ²⁰²⁰ ²⁰²¹ ²⁰²² ²⁰²³ ²⁰²⁴ ²⁰²⁵ ²⁰²⁶ ²⁰²⁷ ²⁰²⁸ ²⁰²⁹ ²⁰³⁰ ²⁰³¹ ²⁰³² ²⁰³³ ²⁰³⁴ ²⁰³⁵ ²⁰³⁶ ²⁰³⁷ ²⁰³⁸ ²⁰³⁹ ²⁰⁴⁰ ²⁰⁴¹ ²⁰⁴² ²⁰⁴³ ²⁰⁴⁴ ²⁰⁴⁵ ²⁰⁴⁶ ²⁰⁴⁷ ²⁰⁴⁸ ²⁰⁴⁹ ²⁰⁵⁰ ²⁰⁵¹ ²⁰⁵² ²⁰⁵³ ²⁰⁵⁴ ²⁰⁵⁵ ²⁰⁵⁶ ²⁰⁵⁷ ²⁰⁵⁸ ²⁰⁵⁹ ²⁰⁶⁰ ²⁰⁶¹ ²⁰⁶² ²⁰⁶³ ²⁰⁶⁴ ²⁰⁶⁵ ²⁰⁶⁶ ²⁰⁶⁷ ²⁰⁶⁸ ²⁰⁶⁹ ²⁰⁷⁰ ²⁰⁷¹ ²⁰⁷² ²⁰⁷³ ²⁰⁷⁴ ²⁰⁷⁵ ²⁰⁷⁶ ²⁰⁷⁷ ²⁰⁷⁸ ²⁰⁷⁹ ²⁰⁸⁰ ²⁰⁸¹ ²⁰⁸² ²⁰⁸³ ²⁰⁸⁴ ²⁰⁸⁵ ²⁰⁸⁶ ²⁰⁸⁷ ²⁰⁸⁸ ²⁰⁸⁹ ²⁰⁹⁰ ²⁰⁹¹ ²⁰⁹² ²⁰⁹³ ²⁰⁹⁴ ²⁰⁹⁵ ²⁰⁹⁶ ²⁰⁹⁷ ²⁰⁹⁸ ²⁰⁹⁹ ²¹⁰⁰ ²¹⁰¹ ²¹⁰² ²¹⁰³ ²¹⁰⁴ ²¹⁰⁵ ²¹⁰⁶ ²¹⁰⁷ ²¹⁰⁸ ²¹⁰⁹ ²¹¹⁰ ²¹¹¹ ²¹¹² ²¹¹³ ²¹¹⁴ ²¹¹⁵ ²¹¹⁶ ²¹¹⁷ ²¹¹⁸ ²¹¹⁹ ²¹²⁰ ²¹²¹ ²¹²² ²¹²³ ²¹²⁴ ²¹²⁵ ²¹²⁶ ²¹²⁷ ²¹²⁸ ²¹²⁹ ²¹³⁰ ²¹³¹ ²¹³² ²¹³³ ²¹³⁴ ²¹³⁵ ²¹³⁶ ²¹³⁷ ²¹³⁸ ²¹³⁹ ²¹⁴⁰ ²¹⁴¹ ²¹⁴² ²¹⁴³ ²¹⁴⁴ ²¹⁴⁵ ²¹⁴⁶ ²¹⁴⁷ ²¹⁴⁸ ²¹⁴⁹ ²¹⁵⁰ ²¹⁵¹ ²¹⁵² ²¹⁵³ ²¹⁵⁴ ²¹⁵⁵ ²¹⁵⁶ ²¹⁵⁷ ²¹⁵⁸ ²¹⁵⁹ ²¹⁶⁰ ²¹⁶¹ ²¹⁶² ²¹⁶³ ²¹⁶⁴ ²¹⁶⁵ ²¹⁶⁶ ²¹⁶⁷ ²¹⁶⁸ ²¹⁶⁹ ²¹⁷⁰ ²¹⁷¹ ²¹⁷² ²¹⁷³ ²¹⁷⁴ ²¹⁷⁵ ²¹⁷⁶ ²¹⁷⁷ ²¹⁷⁸ ²¹⁷⁹ ²¹⁸⁰ ²¹⁸¹ ²¹⁸² ²¹⁸³ ²¹⁸⁴ ²¹⁸⁵ ²¹⁸⁶ ²¹⁸⁷ ²¹⁸⁸ ²¹⁸⁹ ²¹⁹⁰ ²¹⁹¹ ²¹⁹² ²¹⁹³ ²¹⁹⁴ ²¹⁹⁵ ²¹⁹⁶ ²¹⁹⁷ ²¹⁹⁸ ²¹⁹⁹ ²²⁰⁰ ²²⁰¹ ²²⁰² ²²⁰³ ²²⁰⁴ ²²⁰⁵ ²²⁰⁶ ²²⁰⁷ ²²⁰⁸ ²²⁰⁹ ²²¹⁰ ²²¹¹ ²²¹² ²²¹³ ²²¹⁴ ²²¹⁵ ²²¹⁶ ²²¹⁷ ²²¹⁸ ²²¹⁹ ²²²⁰ ²²²¹ ²²²² ²²²³ ²²²⁴ ²²²⁵ ²²²⁶ ²²²⁷ ²²²⁸ ²²²⁹ ²²³⁰ ²²³¹ ²²³² ²²³³ ²²³⁴ ²²³⁵ ²²³⁶ ²²³⁷ ²²³⁸ ²²³⁹ ²²⁴⁰ ²²⁴¹ ²²⁴² ²²⁴³ ²²⁴⁴ ²²⁴⁵ ²²⁴⁶ ²²⁴⁷ ²²⁴⁸ ²²⁴⁹ ²²⁵⁰ ²²⁵¹ ²²⁵² ²²⁵³ ²²⁵⁴ ²²⁵⁵ ²²⁵⁶ ²²⁵⁷ ²²⁵⁸ ²²⁵⁹ ²²⁶⁰ ²²⁶¹ ²²⁶² ²²⁶³ ²²⁶⁴ ²²⁶⁵ ²²⁶⁶ ²²⁶⁷ ²²⁶⁸ ²²⁶⁹ ²²⁷⁰ ²²⁷¹ ²²⁷² ²²⁷³ ²²⁷⁴ ²²⁷⁵ ²²⁷⁶ ²²⁷⁷ ²²⁷⁸ ²²⁷⁹ ²²⁸⁰ ²²⁸¹ ²²⁸² ²²⁸³ ²²⁸⁴ ²²⁸⁵ ²²⁸⁶ ²²⁸⁷ ²²⁸⁸ ²²⁸⁹ ²²⁹⁰ ²²⁹¹ ²²⁹² ²²⁹³ ²²⁹⁴ ²²⁹⁵ ²²⁹⁶ ²²⁹⁷ ²²⁹⁸ ²²⁹⁹ ²³⁰⁰ ²³⁰¹ ²³⁰² ²³⁰³ ²³⁰⁴ ²³⁰⁵ ²³⁰⁶ ²³⁰⁷ ²³⁰⁸ ²³⁰⁹ ²³¹⁰ ²³¹¹ ²³¹² ²³¹³ ²³¹⁴ ²³¹⁵ ²³¹⁶ ²

Hab. in vegetabilis edentibus montium Vellebita, Gungat et Nogh. p. uellu. p. nagh.
avenit h. h. ven. i. o. us. Noh. p. exc. bog.

Alchimiella ^{arvenis} ~~arvenis~~ ^{leg. fl. carn. i. p. 115. pub. fl. exc. 609.}

A. filix ovata, palmate - infid. having 3-5 pty, petiole of. *auriculata*, *argentea*,
 fleshy axillary glomerate. Herb. Aug. 18. 220. *Plat. Mich. 34. A. ovata*.

florib. axillary glomerat. Herb. fl. exsicc. p. 220.
Syn. A. Aphanes Less. fl. herb. p. 55. Alch. fl. jud. p. 34. Aphanes arvensis L.
sp. pl. ed. II p. 179 - Rem. Sw. encl. bot. t. 1011. Lam. enc. bot. pl. 87. Gaertn. 32
p. 145. t. 53. Bed. fl. Rem. t. 995. Col. exsicc. 1. p. 145.

H. p. et *p.m.* 1. 183. Ver. p. 1. 183.
Habitat in arvis et herbis circa Zara, Verlicia, Nittich, et in montibz Dordos, Lescina.
H. mündelii. *H. ver.* ©

Potterium L.

[illegible]

~~Potentilla sanguinea L. 118. fl. 111. Inf. 117. Inm. 118. fl. 119. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840. 841. 842. 843. 844. 845. 846.~~

Sect. I. *Rytidopterium* Ernst gen. sp. p. 1244

Calycis tubus frutiger induratus nigerus. Sepala florum rubiginea. Embryo

Pterium Sanguinalis L. p. pl. ed. II. p. 1411 V. f. 187. delm. p. 44. Hort fl. austr. 2. p. 620. f. 186.
fl. ex. p. 610. Alsch. fl. p. 118. l. 8. bot. weng. n. 730.

P. herbaceum, glaucifolium, ^{involubrum} angulatif., folij imbricatis, capsulis inferius
^{oblongis} subrotundis ovatis, superioribus oblongis achenio ^{tetrangulo-} ~~triangulo-~~ rugoso.
Syn. *Laguridula minor* Sieg. fl. carm. t. 1. p. 110. — Pers. nov. engl. bot. t. 860. Partea l. c. t. 32. Cam
 epit. p. 777. Column. sept. 1. p. 124. — *Nyct. Oskorofriza*. Karwinskaja. Stanovitak.
 Var. *P. polygamum* achenijs stato-2-gonij. — *Syn.* *P. polygamum* Walst. et Kit. fl. hucy. 2. p. 247. Hoff fl.
 austr. 2. p. 620. Hochst. fl. exl. p. 610. — Pers. Walst. et Kit. 2. t. 198.

frequentior unitas ac species

[illegible]

Sect. II - *Leiotenium* DC. prod. l. c.

Calyx tubus prope carnosus brev. Spicae florum cylindricae.

Potium spinosum L. p. ph. 2. II. j. 1412. Hart p. aug. 2. p. 521. ph. p. ex. p. 510. Alud.

fl. jade g. 118. Pet. bot. wing. n. 731.
inset on top

P. frutescens, var. ^{superioris} ~~frutescens~~ ^{subterminalis} ~~frutescens~~, folij pinnatifidis, ovatis
acutis ~~subterminalis~~.

arbenis ~~*fulvipes*~~.
Syn. *Lagenorhynchus spinosa* Bestal. *fl. ital.* 2. p. 171. *Pinguicula spinosa* Gaertn. & p. 1. p. 1. 162.

— Non. Paethen de p. et p. m. t. 32. Salbat. h. v. m. 1. t. 13. Samt. ii. pt. 631. Mox. oxon.

sect. 8. sub. 18 f. 5.

Herb. in maritimis circa Spalato, et secus Pironem de Castelli. H. v. m. 7.

Ordo CXIV. *Spiraeaceae* Benth.

Art. nat. pl. p. 403.

Pl. nat. fl. p. 40
lib. 10
Huf. hemisphaer. regulat. calyx ~~sub~~ ^{non unisulcat} ~~sub~~ ^{marginatus} intus tri. lacinia annulo
perigynio limitata ~~sub~~ ^{libre} 5-partita, laciniis ~~expansis~~ ^{reflexis} - corolla petalis quinque
annulo perigynio insertis patentibus. ~~Thalamus~~ cum petalis ~~affixis~~ ^{infixis}, antheris introrsum
oblongatis. Stylus a summo quina, ovario a calyce libere 2-4-lobis, ovulis per
lobos ut ascenduntibus, stylis ~~totidem~~ ^{stylis} stigmatibusque innumeris totidem. Fructus ex filis
calicis capsulae lobis ~~intus~~ ^{apice} ~~oblongis~~ ^{orbiculis}, seminibus drupis pluribus pendulis, ex
humore coarcturatis recto radicle ad hilum positis.

~~Plants~~ ^{perennial} Plants herbaceous, aut lignosae, ~~many~~ ^{inermis} ~~terrestrial~~, fls. spars. singly or, rarely cymes, sepals, petals, multiple, fleshy, corolla lobes at anthesis

Spiraea L.

Gen. pl. ed. VI. p. 253.

~~Char. Calyx laciniis perispermatis asperatione remotis.~~

Char. Nem ac ordinij.

Sect. I. Chamædion Series. in G.C. prod. II p. 542

Flora hermaphroditi: ovaria libera pluriloculata: caules frutici: stipulif. ^{multif.} ~~stipulif.~~?

[illegible]

9. plijf ook - chrijp overblijp op inco - tenzij ingewinnig, namelijk tenzij, flinby meeng-
winniglyk gelyk, ^{uutlyk} zelyf lating reflexij.

Triby II. Lotese St. podr. yst. nat. II. p. 115.

Stamina didyga aut monadyga. Legumen biser. nec arborescens.
Embryo curvatus, cotyledonibus in germinatione plicatis apertis.

Conspicui Genus

Legumen annulari setting planis VII. N. 1.

Legumen spin. bilobum
aut setosum apertum ^{in b. p.}

1. Alae corollinae margin plicatae - rugosae — 2.

Alae margin planae — 6

2. Filamenta monadyga — 3

Filamenta didyga

3. Filamenta ovarina cuneata — 4

Chamaecyrtis

4. Carina monopetala — 5

Filamenta unum rugosa lobum

5. Alae corollinae libere

Borbo

Cyrtis

Carina didyga

6. Filamenta apice dilatata — 7.

Alae rugosae corollae

7. Filamenta monadyga — 8

Legumen

8. Alae corollinae libere

Filamenta attenuata aut apertae

ovaria

Filamenta didyga — 12

9. Alae cum carina connatae — 10

Alae cum carina connatae

10. Legumen bilobum — 11.

Antyphylis

11. Calyx ^{inflatus} ~~inflatus~~ clausus

Alae corollinae libere

Physanthus

Lotis

12. Petala inter se connata — 13

Legumen unilobum

13. Ovarium rectum — 14

Triplisium

14. Legumen subbaccare

Calyx campanulatus apertus

Myrica

Thymus carpes

15. Corolla tetrapetala — 16

Petala libera — 15

16. Legumen lineare

Ovarium arcuatum

Scilla

Phedra

17. Carina infra apicem mucronata

Legumen subglobosum oblongum

Oxytropis

Melilotus

Corolla pentapetala

Glycyrrhiza

Legum. apertum - artem

Colletia

Carina mucronata

Antyphylis

Subtrib. I. Genistaceae St. l. c. p. 115.

Alae corollinae in margin corollae rugae bifen plicatae - aut plicatae - rugae.

Stamina plurimorum apice attenuata apertis. Petala triplicata, uniplicata aut uniplicata.

[illegible][illegible][illegible][illegible][illegible]

stetit, cum C. praevalens promissis
ann. 1830 die 14 Aprilis natus fecit, nunc dicitur moribus suis ac ego typis
C. Milleri nuncupatione et descriptione in eadem Bot. Zeit. die 28 Junii
- Planchon adque in botanica (Paris 1838) naturae Librum report, et hoc nunc dicitur anni
- cum dicitur natus esse et tunc tunc

Cytisus monspeliensis L. sp. pl. ed. I. p. 240

~~Sect. II. Lemnispenton~~

~~*Cytisus candidus* L. sp. pl. ed. I. p. 240. *Rehb. fl. ex. p. 524.*~~

C. erectus adnase pubescens, foliis caudiculatis, ^{foliis hispido} foliis obovato-ellipticis mucronulatis, vix
vixatis ~~caudis~~ ^{lobis subsymetricis} subtus angulatis & serratis, subcapitis erectis, calycis breviter campanulati
persistens ~~lobis~~ ^{lobis} segmentis bifidis, inferius tridentatis, canna pubescente, legumibus ^{callosis} margine
flavis. vix.

Syn. *Penisa candidus* L. amoen. 4. p. 284 et sp. pl. ed. II. p. 997. Gaertn. de fr. & sem. 2.
p. 329. *Cytisus candidus* Lam. enc. bot. 2. d. 2. p. 246. *Rehb. fl. ex. p. 524* - non
Gaertn. de fr. & sem. 2. t. 151 f. 7.

Hab. in sylvestribus circa Castelnovo di Cattaro. Fl. flavo. inodori. Fl. vix. $\frac{1}{2}$
Obs. ^{terminale hinc} *Stigma capitatum* papillosum sed globosum. Nomen ~~grivizemum~~ ^{grivizemum} (irregulare alteri vixit
agb) ^{antiferendum} plantae huius non candidus est officis (*C. canariensis*) et potiori jure merito

Instit. rei herb. p. 392, t. 213.

Char. Calyx profunde bilobatus. Corolla ^{tubulosa} alba, speciosa coramari; inferne margine vexillari
nigro, carina accuminata sublongioribus. Stamina monodelpha. Stylus
subulatus, stigma terminale apiculato lobato. Legumen ^{curva oblongum} baccatum isthmio tenui
vergete, ungues inter foveas. — Obs. Fructus ord secundae fere ut
strobilaceae, see 173 n. s. p. 68 E.

Lepidoptera L. sp. pl. ed. II p. 1018. ~~Roth fl. hexag.~~ 1797. exd. 19m. ~~hexag.~~

[illegible]

Non. Monif fl. vord. 1. LXXII f. 1. J. Baach. hift. 2. p. 289. -

Jan. Morio fl. v. r. 1. LXXII p. 1. *S. v. r. 1. 0*
 Hab. in grato. *fl. v. r. 1. 0* *S. v. r. 1. 0*
 morio, in *S. v. r. 1. 0* *S. v. r. 1. 0*
 insula *S. v. r. 1. 0* *S. v. r. 1. 0*
 insula *S. v. r. 1. 0* *S. v. r. 1. 0*

Obs. Ab. h. differt L. Digitalis Hook. cum ipsis a planis comp. fl. longioribus, unguiculatis, et calycis tubo imprim. integro: vixit etiam ex Hook. *Bot. Beech. pl. aeg. et urb. p. 29* tab. V. - Calibus quibusdam asperis.

L. alba L. ^{et} calycibus ^{et} bracteolatis, latis nervis rubris, et viridis
 albis ^{et} calycibus ^{et} bracteolatis, latis nervis rubris, et viridis
 ortu inter folia L. Ternis ^{et} calycibus ^{et} bracteolatis.

Subtrib. II. Anthyllideae Koch syn. fl. german. p. 123.

Alae corollae planae. Stamina ~~separata~~ ^{filamentis separatis latioribus} ~~aut adhaerentibus~~. Filia pinnato, pinnato.

Conspectus Generum

Stanimira monadelphica — Vid. N^o 1.

Stamens Diadelph. var. N. 2.

Alac ^{convly} *liberae*

Alae cum carino connexae

Quoniam

Antyllus

2. Calyx reflexing clasped
Physanthyllis

Calyx campanulatus agestus
Hymenocarpus

Ononis L.

Gen. l. ed. t. 263
Char. Calyx campanulatus, quinquefolius, corolla tetragynata alig lobis, corina acumi-
nata sublongioribus. Stamina monandria. Stylus geminulato-ascendens, signato subop-
tato pubescentibus. Epimenium longiusculum semine inclusum. Cap. triflorum. Ad. 2-3
Sect. I. *Begonia* Griseb. sp. fl. v. n. 17-12

Sect. I. *Dugranus* *Yntio*
Lucenae *ovatum* *vactum*

[illegible]

Icon. Arch. icon. bot. 1. t. VII f. 14. *Simulium circa 1890*

[illegible]

Ononis arvensis ^{non L.} Lightf. fl. Scot. p. 386. Sm. fl. brit. 2. p. 758 excl. var. ? Best. fl. ibt. vii. p. 37
O. caulibus decumbentibus erectis, affinis, albescens, unifloram pilosius, ramulis pinnatis foliatis floribus
v. platis oblongis acutis, stipulis glabris, calyci subaequalibus, seminebus v. minutis; ~~seminebus~~
Icon. Sow. angl. bot. t. 682. ^{fl. Dan. t. 783} Fl. dan. t. 68. Presl hist. stir. p. 60 — Ulyss. Gladst.
Hab. in humilis incul. Lago. Fl. yulii. Fl. aestate. H.
Obs. Differt a praecedente ~~habitu~~ ^{habitu} tantummodo foliis acutis. An cum Smithio ^{h. onis} in unum speciem
reducenda? *O. arvensis* L. 1484. nat. ed. x. t. 2. p. 1159. est *O. hircina* Jacq.

Ononis spinosa L. 14. pl. 2. t. 5. p. 1086. excl. var. a. Ref. 1879. delm. y 42. Hort. fl. austr.
2. 9. 516. Pl. ex. p. 517. Alt. fl. jedr. y 163. excl. syn. L.

[illegible]

Tarn Regi.

Hab. in arvis, laevis humilis ad riv. ad Zava, Schinia, Iran, Gales, Pagan

et in impeliq. Fl. var. gr. asplata.
Obs. Spec. a d. Bernhartii propit nuntio est. pini ^{indubitanter longioribus.}
Hoffl. fl. austr. 2. p. 316.

Ononis repens L. p. l. d. D. 1006. ~~Reich. syn. p. 179. pub. p. 179.~~

[illegible]

hab. in arenis maritimis una Baden, ubi legit consil. G. S. Kunzinger. Fl. rose.

Pl. aspera. $\frac{1}{2}$

Pubes caulem undique vestit, sed unguis^{et} alterve longior est ac in latere opposito

Ononis brachystachya Vif.

[illegible][illegible][illegible]

Ononis nictitans L. p. pl. ed. II p. 1007. Gussone. syn. fl. sic. 2. p. 256

0. ^{head}longer, carities evenly lanuginous - immortary, slender ovality bicurvally glab, flange
by linear-alloy subintery, flexibly ^{speci}spicily, brachy thepulanty ovary ventricory
vanilla-albity calyces ^{oblong}oblongity
height. 2, sect. 2, 4, 17 f. 4.

Sten. Dillen. h. altham. 2. 24. 4. 27. Mon. h. 1. 2. sect. 2. 2. 1. 1.
Hb. in insula Loma d. Attornaria in gloriois temp. luvulid. Castle ferns rose. H. v. 1. 1.

Medicago ^{cordata} Desv. in Lam. enc. bot. d. r. Pad. III. p. 608
~~maritima~~ ^{lanceolata}

[illegible]

Syn. M. maculata W. sp. pl. III. p. 1412. ~~Hyss. fl. austr. II. 386.~~ *Nob. fl. austr.*
504. *Alsch. fl. austr. p. 127.* *Pitt. bot. weyn. n. 596.* (excl. *syn. descr.*) - *Non. Monj.*
fl. ind. t. 2. *Sow. engl. bot. t. 166b.* *flora tetr. inv. t. 88 p. 12.* *Camer. herb.*
med. t. 25.

Hab. in demerit & herbis circa Larn, ⁴/palat, et in rugulis Oryza et Chrys
 fl. flav. fl. vire. ③

H. flav. fl. ven. C
Obs. Nomen generum a cl. Desrousseaux huius plantae insignis et a
^{utrum insensibile sit}
Willdenowio ipso ^{in hisce} injuria mutatum est ^{et alia} quod inde accipit aut

11 *Medicago* ^{my} *carstenkii* Lag. collect. 1. p. 86. Hort fl. austr. fig. 286. (chb. fl. ex. g. 501.)

[illegible]

syn. M. Bocca Balb. fl. Guin. II. p. 84 - Leon. Ray. ii. var. t. t. b. 156. ^{apt.} Nac. et Balb. fl. Guin. II. t. XI.

Hab. in graminosij circa Agass, ubi legit Dr. Neumayer. R. Pers. musci m. t.

12 *Medicago Gerardi* Walst. & Kit. in W. sp. pl. III. p. 1415. Vfl. 1 liny. solum p. 41. Hort. fl. austr.
III. p. 387. Rhod. fl. ex. g. 502. Alab. fl. austr. g. 177.

[illegible]

Syn. M. villosa DC. fl. p. IV. p. 545. *M. polymorpha* & *viridula* L. (syn. d. II. p. 1098) — *Nov. fl. rar. tab. XLIII. Morison* hist. pl. sect. II. tab. 15. f. 18. *J. Bauh. hist. pl. 2. p. 385. f. 3.*

hab. in arvis et apricis circa Lav. Sebenic, Mn. Polato, Ragusa, et in insulis Lefina - H.
proflori. fl. vire. ①. ... quatuor ad sex.

Obs. Flor. flori. Pl. vasa^o
Medicago ulmiflora W. & A. pl. III p. 1411. Morif fl. ind. t. p. 482. Guss. syn. fl. sic. II p. 769
M. ovata-lanceolata incipit dentatis, foliis cuneatis-obovatis lenticularibus, pedunculis uniu-
floris, leguminibus cochlearibus subglobosis villatis, gynij suboblongis oblique nervosis reticulatis, uterinis
infra glomeratis spicatis, spinis divergentibus intertextis bracteis carnulatis
Hegn. M. polymorpha x citrifolia L. (pl. ed. II p. 1099). Non. Morif fl. ind. ed. II. Morifon hist. pl. sect. II.

Hydn. M. polymorpha x *ciliat* L. p. gl. ed. II. p. 1699². - Pen. Mois fl. ind. ob. Ind. Moiron h. p. gl. 1 oct. 17

28 - *Pres. 194 fl. ric. II p. 371. M. bicycla*

M. laevis Maly in fl. ung. austr. p. 350 non Willd. et exl. yn.
D.C. cat. h. univ. p. 125 - non - *Monif fl. ric.*

I. tab. XII B. 2. 3.

Herb. in muris, rivulis Lemn. fl. flav. fl. Apr. Mayo. ©

Obs. Scroph. repas peccis a cl. Monif prolati, forma ^{plant} ^{obstant} ^{legum} ^{exhibetur} ^{ab ejusdem} ^{cl. Quichy} ^{scroph. repas} ^{utary} ^{M. prolati} ^{quoad} ^{plantam} ^{jadrasas} ^{a fl. Alchinger} ^{Maly} ^{in fl. ung. austr. p. 350} ^{quoad} ^{plantam} ^{invol. Lemn. ab egr.} ^{est M. dimidiata} ^{var. brevifolia} ^{quoad} ^{plantam} ^{invol. Lemn. ab egr.} ^{Potterii} ^{littam} ^{peritum} ^{ad ex. p. Medicago} ^{liberly} - *M. procer* D.C.
invol. Lemn. ab egr. p. 350 ^{quoad} ^{plantam} ^{invol. Lemn. ab egr.} ^{Potterii} ^{littam} ^{peritum} ^{ad ex. p. Medicago} ^{liberly} - *M. procer* D.C.
invol. Lemn. ab egr. p. 350 ^{quoad} ^{plantam} ^{invol. Lemn. ab egr.} ^{Potterii} ^{littam} ^{peritum} ^{ad ex. p. Medicago} ^{liberly} - *M. procer* D.C.

Medicago tuberculata W. p. fl. d. II p. 1410. *Monif fl. ric. I p. 443. Pres. fl. ric. II p. 370.*

M. stylis ^{orto} ^{lanceolatis} ^{primis} ^{dentatis} ^{foliis} ^{curvatis} ^{obovis} ^{superioribus} ^{hombis} ^{ovulatis} ^{pedunculis} ^{bisectis} ^{leguminibus} ^{aristis} ^{cochleatis} ^{ovatis} ^{productis} ^{utrinque} ^{gibbosis} ^{ternariis} ^{glabris} ^{gynis} ^{subquaternis} ^{avariis} ^{utrum} ^{acutis} ^{utrinque} ^{laevibus} ^{tuberculatis} ^{tuberculis} ^{avariis} ^{utrum} ^{acutis} ^{utrinque}

Syn. M. pinetorum D.C. fl. p. v. p. 569. - non *Monif fl. ric. tab. XLIV. Rivin.*
tab. iv. t. 86 f. 6. Monif hist. fl. ric. II. t. 15 f. 6.

Herb. in sativ. circa Zava et Ragusa. fl. flav. fl. Apr. Mayo. ©

Obs. Tubercula ^{nunc} ^{utrum} ^{lateris} ^{adnata} ^{ipsum} ^{aequant} ^{nunc} ^{ultra} ^{utrum} ^{quibus} ^{libris} ^{et} ^{auribus} ^{eminent} ^{longi} ^{tubercula} ^{inveniantur} ^{per} ^{globulosa}

Medicago turbinata W. p. fl. d. III p. 1409. *Hort. fl. austr. p. 501. Monif fl. ric. I p. 443.*
M. stylis ^{orto} ^{lanceolatis} ^{primis} ^{dentatis} ^{foliis} ^{obovis} ^{superioribus} ^{hombis} ^{ovulatis} ^{pedunculis} ^{bisectis} ^{leguminibus} ^{aristis} ^{cochleatis} ^{ovatis} ^{productis} ^{utrinque} ^{gibbosis} ^{ternariis} ^{glabris} ^{gynis} ^{subquaternis} ^{avariis} ^{utrum} ^{acutis} ^{utrinque} ^{laevibus} ^{tuberculatis} ^{tuberculis} ^{avariis} ^{utrum} ^{acutis} ^{utrinque}

Syn. M. polymorpha ^d ^{turbinata} ^{W. p. fl. d. II p. 1098. M. oligosperma}
Pres. fl. ric. p. 370 et 194. fl. ric. II p. 375. Monif hist. fl. ric. II. t. 15 f. 5
t. XLV A. Rivin. tab. iv. t. 86 f. 3. Monif hist. fl. ric. II. t. 15 f. 5

Herb. in herbis prope Spalato loco dicto alle Botrelle unde nunc prof. Petter.
M. flav. fl. Apr. Mayo. ©

Sect. II. *Lupulina* ^{rev. in D.C. prod. II p. 172.}

Legumen ^{scirpiforme} ^{monospermum} ^{inermis}

Medicago lupulina ^{W. p. fl. d. II p. 1099. Hort. fl. austr. p. 385. fl. fl. austr. p. 501. Alchinger}
M. stylis ^{orto} ^{acuminatis} ^{basibus} ^{subrotatis} ^{foliis} ^{obovis} ^{utrinque} ^{imaginatissimis} ^{aristis} ^{cochleatis} ^{ovatis} ^{productis} ^{utrinque} ^{gibbosis} ^{ternariis} ^{glabris} ^{gynis} ^{subquaternis} ^{avariis} ^{utrum} ^{acutis} ^{utrinque} ^{laevibus} ^{tuberculatis} ^{tuberculis} ^{avariis} ^{utrum} ^{acutis} ^{utrinque}

Obs. In diffinientis radicigenis speciebus character praevalens inter ipsas
 presentia et dispositio nervorum in altera aut utroque leguminis parte hinc solum
 tenuitas vel crassities. Caeteri characteres potius a pubescentia, foliorum
 forma, stipularum dispo., pedunculorum longitudine, rursus ex ipso
 nervorum numero, et apicis forma ^{magnitudine} aut defectu ^{hinc illuc} foliacei congeri.

Trigonella L.

Gen. pl. ed. II p. 275.

Char. Calyx caezumilatus quinquefidus. Corolla tetragynata decidua alij cavis stylo
 breviora supra angulum connexis. Stamina didypha, filamentis utroque extre-
 mitate aequalibus. Stylus globosus, stigma ^{ovario} ~~obtusum~~ ovario recto. Legumen
 lineare oblongum vix diffrangens polyspermum. — Lf. Didyphum Ord.

Dreandri Lf. p. 28.

Phob. Hexap. 500.

glaciata Steud. cat. h. boreal. 1808. p. 112. Koch syn. fl. germ. ed. II p. 181.

Trigonella ~~gracilis~~ ^{floribus axillaribus subterminalibus, leguminibus}
~~gracilis~~ ^{umbellatis, subterminalibus, pedunculatis}

T. diffusa, foliis cuneatis-obovatis, obovatis, ^{leguminibus} ~~obovatis~~ ^{umbellatis, subterminalibus, pedunculatis}
 nervis reticulato-nerviis foliis, rostro suo vix longioribus, nervis, subterminalibus,
 nervis, granulato-scabris.

Lyn. T. prostrata DC. fl. fr. v. p. 371. Alsch. fl. jard. p. 176. — Pott. bot. weing. u. 974.

T. Danum graecum B. L. p. fl. ed. II p. 1095. — Non. Moris fl. ind. t. I. IV.

T. Dan. hisp. fl. II p. 385 f. 1. Dalech. hisp. fl. I p. 164 f. 2. — Hb. Sic.

linea rostrata. Pistacica.

Hab. in pasuis rivicis et locis agris interis circa Zara, Sebenico, Veglije, Spalato,
 et prope Durno, maxime in rivis Veglije et Lacinia. Fl. pallid. ochroleuca. Fl.
 vere. ©

Obs. Species huius militis est T. Danum graecum L. quae pro istius ^{caulis erectis} ~~differt~~ ^{leguminibus}
 stylo longioribus rostrum stylo longiore excedentibus et seminibus ^{subterminalibus} ~~subterminalibus~~

Trigonella monspeliaca L. p. fl. ed. II p. 1095. Hort. fl. austr. II p. 383. Phob. fl. exl.
 p. 500. Alsch. fl. jard. p. 176. Pott. bot. weing. u. 973.

T. prostrata, foliis cuneatis-obovatis rhombicis floribus axillaribus subterminalibus
 leguminibus ^{umbellatis, subterminalibus, pedunculatis} ~~obovatis~~ ^{umbellatis, subterminalibus, pedunculatis}

Non. Waldb. et Kir. fl. rar. hung. II p. 142. Pers. tab. rar. t. 80. Moris hisp. fl.
 ed. II. t. 17. f. 4.

Hab. in agris multis ad vias circa Zara, Sebenico, Spalato, Durno, et in agulis
 Cherso, Veglije, Lacinia. Fl. lutei exiguu. Fl. vere. ©

Trigonella corniculata L. p. fl. ed. II p. 1094. Vp. hng. del. p. 50. Hort. fl. austr. II p. 382.
 Phob. fl. exl. 500. Alsch. fl. jard. p. 176. Pott. bot. weing. u. 972.

T. erecta, foliis cuneatis-obovatis mucronatis-similibus, floribus axillaribus longe pedunculatis
 leguminibus ^{oblique nervis} ~~oblique nervis~~ ^{oblique nervis} ~~oblique nervis~~

29. ~~M. ney~~ ~~H. H.~~ ~~Feb. 6 Aug 1899.~~ Spikes represent integument instead of scales.
occurent.

Melilotus italica Lam. fl. p. ^{II} p. 594. Hist fl. austr. II. p. 365. f. l. l. fl. ex. p. 498.

Monif fl. sand. I. p. 460.

M. spuriis infundib, lanceolatis-lanceolatis, lumbicis-dentic, superioribus lineari-
 integris, plus minusve denticulatis, obliquis, infundib, oblongis, superioribus oblongo-obovatis,
 vixim, laxifloris, alii corintheaeaeque vixim brevibus, superioribus oblongo-globosis
 obliquis ^{pendulis} ~~pendulis~~, stylis reclinatis-vixim, glabris, stylo laterali.
 Host. Fl. austr. Ital. 365

Lyn. M. rotundifolia Tenor. fl. rep. prod. p. 56. *Trifolium multichne italicum* L. p.
fl. ed. II. p. 1075. *T. italicum* W. p. pl. III. p. 1556. — Ten. Ten. fl. rep. t. 186. f. 3.
Camer. herb. med. t. 29. Minima bip. pl. t. 1. f. 1. f. d. t. 16. f. 4. semi-nudis.
Hab. in fasciis collinis et agris circa Sebenico, Rari, Almonia, Poggio, Suedia,
ad pedem montis Moson, et in rivula Lerina. Fl. flavi, fructus immixti
cruni. Fl. anc. O

Sect. II. *Magioris* Ser. in H. prod. II p. 188

Legumina transversa subaristata vicia

Melilotus elegans Salzm. in DC. prodr. II. p. 188. Moench fl. rar. I. p. 462, Pers.

14n. fl. vic. II. p. 323.

100 ft. 11. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840. 841. 842. 843. 844.

Icon. Monif. fl. card. tab. LVII.

Ab. in herbary loc Plagium ditto in nr. Leine. H. pallid.: H. vire ♂

Sect. III. *Campylorhynchus* Ser. l. c. p. 189.

Legumina ^{dense} ~~acuminata~~ et concentric plicato-sulcata.

Melospiza sulcata Serp. Pl. atl. II. p. 193. *Rebb. Pl. ex. p. 198.* *Merr. Pl. ind. I, 463.*
amphibolus

N. hypolepis *camulata* *laevius* *dentic*, *repens*, *intellig*, *foliis* *argute* *serwis* *obliq*,
inferius *obovatis*, *repens* *unite* - *obliq*, *venis* *obliq*, *his* *canis* *pauca*,
repens, *repens* *aliquo* - *subtus* *repens* *glab*, *1/4* *marginis* *repens*.
Sen. Influen *Melby* *indica* *L. p. p. d. II. p. 107*. *S. mauritanicus* *W.*

Syn. Myrtum Melaleu. F. indica L. fl. Ind. II p. 1077. *S. mauritanicum* W.
fl. Ind. III p. 1554. — Leon. Moris fl. Ind. L. LIX A. J. Barb. Hist. fl. II p. 371/4.
Ch. v. Ind. Linn. V. fl. Ind. V. v. c.

Hab. in agris circa Schmale, Iran, Persia. Fl. flav. fr. vir. $\odot$
Stem erectum et hirsutum. compacta calyx.

Obs. Soter varietas florum leguminosorum caespitum et hic est M. compacta satem. M. helica
 et β compacta Minor fl. radii 404. t. 418. B. — ~~Vox~~ ^{Vox} graeca GOUTIS raga
 latine rypis subinde et pronuntianda est, cum littera γ graecorum γ latine
 nun respondat, sed saepe nexio videtur scriptum gallice, qui cum γ latine
 eodem modo ~~pronuntiant~~ ^{pronuntiant} ac γ , perperam loco ~~longius~~ ^{heio} illud scribere solent. ~~perperam~~
 Pakida, Pakidoghimu, Caetomiy et reliqua —

Trifolium Journ.

Instit. v. l. h. v. j. 404. t. 228.

Des. Caty × *gymnophylla* vel *obolobolus*. Corolla persistens, petalis nunc ovariois inflexis in tubum corollae, nunc 1-2o vixite libris, alij carinam oblongam superantibus. Stamina corollae adnata. Dactylus, filamentis apice subulobatis. Stylus globosus. Legumen induratum, nunc ortum 1-2-fermen, nunc oblongum 3-4-fermen. — *Clas. Dactylus* et *Brundia* 1944. 100.

Conspectus Lectionum

Quercus calycis exannulata Nutt. W. N. 1.

Frax calycis annulo pilisve cincta
Lagopus

1. *Harporhynchus corollae* _____ 2

Flores umbellae capituli apetalae

2. Flors sensibly — — — — 3

Calycospermum
flor. pedicellati — vid. N^o 5

3. Calyx demum inflatus. — — — 4

Calyx immitatus
Paramesur.

4. *Pekla ovina* cornata

Vexillum a petalis liberum

Verillium ^{*Ragifera* *Wibben*} ~~*a petalis*~~ *libanum* — 6
5. *Verillium* ~~*a petalis*~~ *libanum*

Vesicastrum
~~S. Vesillum a pet. of libanin~~
~~Setaria ovacea cornuta~~
~~Setaria ovacea cornuta~~
~~Setaria ovacea cornuta~~
 Olusoria

6. *Veritellum convexum*
Veritellum convexum
L. L.
L. L.

Vexillum complicatum
Lepinaster

Sept. 1. *Lagopus* Koch. syn fl. germ. v. 1. p. 167.

Flory ^{conformis} ~~acutifolia~~ ~~longifolia~~: calycis fruce annulo intus cincto callosi, auriculis, tubo et
dentibus immixtis: petalis ovatis connatis, venis floris. Legumen calycis
tubo inclusum, ~~semine~~ ^{semine} ~~laevi~~.

Spica aut capitulum * *Annua* *heri medium*
Compositum terminale solitarium, pedunculatum. Folia omnia alterna.

Inula angustifolia L. sp. pl. ed. II p. 1083. Hort. fl. angl. III p. 72. Pet. fl. ex. J. 493.
Alch. fl. ind. p. 166. Pet. bot. weng. n. 953.

[illegible]

Leont. Trin. tab. iv. t. 16. Parlat. icon. 698. Monion huff gl. sect. t. t. 13 f. 1. - Nigr. Indiv.
Triglochin, promissus huff et repens huff.
Hab. in pariet. et vici. huff agror. tokiy Deludat. Fl. vesp. Fl. v. c. 3
Fl. v. c. 3. 82. t. 1. h. 1. c. 1. t. 338.

Trifolium intermedium Guss. cat. h. in Boissard. p. 82 et synfl. sic. II. p. 335.

7. *caud. long. pilis long. oblongo-lanceolatis adpress. pilisq. m. pulv. canis. subulatis p. long. cylind.*
ca. oblong. ~~interf. p. long. cylind.~~ ^{oblong.} pilisq. d. h. subulatis p. long. cylind. terminatis tubum ~~secundum~~ ^{recessum}
expandit. subquadratis infus. sublongior

hab. in aëre humido circa Plagijs insulae Levia, ^{unde vixit} ~~ubi~~ legit prof. Solis H. albi: H. ven
it acuta. ②

3. *Thlasium inaequalis* B. *Molineri* Sering in St. p. 190. ^{Ad. H. ex. 199.} *Mong. fl. 1847.*
I. p. 468. *Pers. syn. fl. sic. II. p. 335.*

Syn. I. incusculentum Vig. 1890. Delia 79. Hort fl. austr. II. p. 28. (exhib. col.) Ad.
Nöthen. Pfl. in Nothfl. et in herb. norm. n. 136.
Pfl. jard. g. 167. Pfl. bot. wien. n. 900 L. - Icon. Dandel. t. 169. l. Band. hift
pl. II. p. 276 f. 4. Chab. scagr. p. 161. f. 6. Spinn. herb. iur. t. 17.

4. *Infulum Heltatum* L. sp. pl. ed. it. p. 1085. immo p. 1086 f. 7. g. 1. h. 1. g. 1. d. 1.
Alsch. fl. jard. p. 167. R. bot. weng. n. 968.

Icon. low-engl. bot. t. 1545. Lam. enc. bot. pl. 613. f. 4. a. b. c. d. Jacquin. de fr. et sem. t. 183.

Daml. ion t. 866.
In amig bouque houlaly whing Salinity prapery H vori. Jc. vor. O

2. *Asplenium latifolium*, var. *latifolium*
Asplenium latifolium (L.) Sw. f. *latifolium*. Pet. bot. weng. n.
 962. Moif. fl. ind. t. p. 484.
 f. *latifolium* oblongo subtriangul. pubescenti, stipulacul.

Syn. T. vixidem Lari fl. inf. II p. 189. D. irregularis Fourv. act. Boul. III p. 331. — Pan
Lar. C. i. t. b. l. c. 1. Low. conf. bot. t. 220. Pluken. phyt. t. 113. f. d.

6 *Myriophyllum repens* L. var. *trifid.* p. 46. Pet. bot. wurg. n. 970. Maly. ex.
pl. imp. austr. p. 353.

pl. imp. austr. 353.
 3. caulebus piliis vix vix; subintegerrimis pubescentibus, hirsutis, ~~caulis~~ ^{caulis} canaliculatis linearibus
 vix, ^{pubescentibus} ~~integerrimis~~ ^{pubescentibus} ~~caulis~~ ^{caulis} ~~canaliculatis~~ ^{canaliculatis} ~~linearibus~~ ^{linearibus}

11. *Trifolium* ~~hercynicum~~ Bocconi var. in Act. dell' Acad. Ital. I. p. 191. Feb. fl. ex. p. 491.
Alch. fl. jod. p. 168.
T. pubescens, foliis ^{denticulatis} cuneato-oblongis, inferius cuneatis, strobiliformibus, stipularum caudicibus
linearis-stanceis, spicis oblongo-^{ovatis} cylindricis, ovisque subgeminatis, foliis involutis,
calycis dentibus lanceolatis-subulatis connatis, rectis irregularibus tubo decemnerariis
breviter corollae subaequalibus.
Bon. Boccon. nup. d. g. ens. var. t. 104 f. inf. ext.
Hob. in fasciis Xenox pro Zar, et ad vineas prope lacum Boccagnazzo. Cor. rubella.
fl. acetate. ©
12. *Trifolium tenuifolium* Xenox. fl. nov. prod. p. 44. Del. prod. 14. t. nat. II. p. 206.
T. pubescens, foliis ^{denticulatis} cuneato-oblongis, inferius cuneatis, strobiliformibus, stipularum
caudicibus linearis-stanceis, spicis ovatis ovistylis solitariis sessilibus foliis involutis,
calycis ^{pubescentibus} dentibus linearis-subulatis rectis, irregularibus tubo decemnerariis
aequantibus corollae duplo brevioribus.
Syn. *T. Xenocanum* Boiss. et Griseb. Diagn. pl. orient. II. p. 26. *T. Bocconi* β. longifl.
vum Boy et Charb. in exp. scient. de Mor. part. botan. *T. Bocconi* β. tenuifolium
Griseb. spec. fl. venet. I. p. 23. — Bon. Xenox. fl. nup. t. 177 f. 3.
Hob. in ^{apertis agris} circa Zar, ubi legit A. Andrich. Fl. albi. fl. vir. ©
Obt. Sam. ^{corymbis majoribus} calycis ^{inaequalitate} albis calyce duplo longioribus, et rostris elongatis carinum valde excavatum
dente apice ^{concavitate} a punctura, a quo stromma diffinit el. Boissier l. c. quam confer.
13. *Trifolium striatum* L. sp. pl. ed. II. p. 1085. Hort. fl. austr. II. p. 375. Feb. fl. ex. p. 491. ~~Act.~~
fl. jod. p. 168.
T. pubescens, foliis ^{denticulatis} obsoletis denticulatis cuneatis, strobiliformibus, inferius cuneatis,
strobilis, stipularum caudicibus lanceolatis-stanceis, spicis ovatis oblongis subternariis
sessilibus foliis involutis, calycis ^{pubescentibus} dentibus subulatis rectis subaequalibus
tubo decemnerariis ^{max. ventricosis} brevioribus.
Bon. Waldst. et Kit. fl. rar. hung. I. t. 25. Lav. encl. bot. t. 1843. Vaill. bot. parisi. t. 33. f. 2
Dart. II. nup. 869.
Hob. in fasciis vicinis, et gentis circa Zar, in Ullabith, circa Velika, sign. et in
insulis Lepim et Langola. Corolla parva rubella. fl. vir. ©
14. *Trifolium rubrum* L. sp. pl. ed. II. p. 1084. by. nup. del. p. 50. Hort. fl. austr. II. p. 374.
Feb. fl. ex. p. 492. Alch. fl. jod. p. 168. Pet. bot. univ. v. 967.
T. adpressum pubescens, foliis ^{denticulatis} cuneato-obovatis nervosis, inferius obovatis, stipularum caudicibus
lanceolatis villosis, spicis ovatis ovistylis solitariis sessilibus foliis involutis, calycis ^{pubescentibus}
dentibus lanceolatis ^{case commixtis} irregularibus rectis semine recurvo potius
tibis tubo decemnerariis cylindricis corollae longioribus.
Bon. Low. encl. bot. t. 903. Vaill. bot. parisi. t. 33. f. 1. Samml. icon. t. 870.

- [illegible]

[illegible]

obliqu^e irregularis obov^{at} rep.
Obs. Petio^{li} I. ^{lappaceus} ^{separatim} et 7. ^{lanceolatus} ^{medius}: ab illis

~~Trifolium sphaerophyllum~~ Desf. fl. atl. II p. 201

9 - *Trifolium affinis* Ehrh. bot. VII p. 165. Hort fl. austr. II p. 369. Pöchl. fl. ex. v.
194. Monif fl. scand. I p. 479.

7. patule villosum, ~~stigma~~ ^{stigma} ~~altrax~~ ^{altrax} ~~negativ~~ ^{negativ} ~~opposit~~ ^{opposit} ~~strobil~~ ^{strobil} ~~ovoid~~ ^{ovoid} ~~obovate~~ ^{obovate}
obovate dimidiata obtusius utrinque, ~~stigma~~ ^{stigma} ~~altrax~~ ^{altrax} ~~negativ~~ ^{negativ} ~~opposit~~ ^{opposit} ~~strobil~~ ^{strobil} ~~ovoid~~ ^{ovoid} ~~obovate~~ ^{obovate}
nig. strobili triplicatis capitulum subrotundum totum scilicet ~~capitulum~~ ^{capitulum}
involucris, calicis, ~~stigma~~ ^{stigma} ~~altrax~~ ^{altrax} ~~negativ~~ ^{negativ} ~~opposit~~ ^{opposit} ~~strobil~~ ^{strobil} ~~ovoid~~ ^{ovoid} ~~obovate~~ ^{obovate}
tubo dicussu villosulo long. longioribus corollae subglobois.

Icon Waldst. et Kit. pl. rar. hung. 1. t. 50.

Hub. in aprilis 2 vias circa Cittavulvi, ubi legit d. P. Nisito. H. rubell. H.
vireo

Sect. II. *Parmery* *Eriob. pic. florum*. t. p. 28
~~*Parm. rhytidota*. t. p.~~

Florae conformis: calycis foveae laevi tubo et drubney immutari; petalis
omnibus connatis - Legumen exsertum, siliqua, seminibus, tuberculis

26 *Trifolium lavigatum* Poir. voy. en Barb. II p. 219. Desf. Fl. Atl. II. p. 195. Jacq. syn. ^{ambr.}
H. ar. II. p. 340. *Trifolium lavigatum*

[illegible]

Syn. *D. strictum* Waldb. et Kit. fl. rar. hung. 1. h. 37. Hist. fl. austr. 4 p. 366. Ped.
fl. ex y. 1891. non L. — Non. Desf. l. c. t. 208. Waldb. et Kit. l. c. t. 37.

Michel nov. gen. pl. t. 25. f. 7.

hab in parvum lumen prope Zann, et inter Urshika et lign. H. carnei. H. mag.
 Junis. ☉

Obs. D. strickum L. (opt. ed. II. p. 1079) juxta descriptionem ibidem datam ab hoc
 Desfontainium cum D. parvifloro Ehrh probe convenit, recte observante
 d. Hoch in syn. fl. german. ed. II. p. 141, excludit tantum Michxali synonymum
 nostrum perstruit, quod non minus ~~est~~ ^{cujusque citatio a Linnaeo postea} ~~sapienter~~ ^{opportunitate}
 et omnino compendiosius causa est habenda.

- [illegible]

[illegible]

H. albi. 71. April. 4

[illegible]

Triplarum piliforme L. p. gl. d. t. p. 1088. *Junon* p. n. fl. n. t. p. 347. Smith encl. fl.

2.4.v.3.f.310

2. 2. v. 3. p. 310
D. guillelmi, *caulibus* *diffusi*, *filibus* *uncati* - *obtusis* *reductis* *conjugatis*, *terminalibus*, *obtusis*, *perianthio* *axillaribus*, *capitulis* *umbellatis*, *subquadratis* *lobis* *dentatis*, *antheris* *obtusis*, *calycis* *densis*, *reticulatis* *supra* *bracteas*, *ventris* *angustis* *levis*, *gynemine* *stylis* *bracteis* *aequalibus*.

Syrphoctonus *micans*
Syrph. T. *micans* Vijs. fl. lyb. p. 48. hoch 1 mm. flegel d. II. p. 195. ♀. platform B. mic.
flegel Leing. in Zl. prod. II p. 266. — *hon.* Vijs. l. c. t. XIX f. 3. ♂. tag 1 mm.
Ship. Brit. d. III - t. 14 p. h. Michel. gen. sp. t. 25 f. 8.

hab. in grandibus apris montium Papporum, nec non in montibus inter
Cantonem & Cantonem de Canton. Mexici lateri. 30. May, 1860.

Ob. Ab hoc differt I. minus Im. l. c. quod roburque excedit, Apiculis ovatis nec
basi latioribus ac medio, capitulo subaequilongis. Pollicetum in istis irregulariter
petiolatum, phlo. ^{reperit} intermedio longius petiolato -
Lotus α .

P. Dargenium Vourn.
Sen. pl. d. VI p. 388
Just. ra herb. p. 391 t. 211.

[illegible]

Hygicum

Sect. I

~~Longenium Journ. inst. v. herb. p. 391 p. 211~~

Sept. 1: *Petropedotobus* Scop. fl. carn. II. p. 87.

Flora sublobata
Quercus *Carina* *corolla* *rostrata*. *Legum* *cyathaceum* *quadralobum* aut *pentapetalum*

Lotus tetragonolobus L. sp. pl. ed. II p. 1089. Hist. fl. austr. II. p. 379.

*L. flab. ellipt. inflex. divaricat. apiculis ^{obtus.} serratis, pediculis unifloris foliis
nullis longioribus, lymenibus ^{obtus.} alio planis, ipso angustioribus.*

Syn. Tetrazonella elegans Joth. tent. fl. germ. T. p. 323 (sch. Kexip. 506. Allch. fl. par. p. 176).
— Kon. Nag. fl. austra. IV. t. 361. Chap. h. p. II. p. 244 f. Grevh. & Proct. ann. t. 153.

Ann. - tetr. ivr. t. 29.

Hub. in part ^{humid} ^{carine apice fusco} ^{fl. roseo} ^{ven. 7} ^{l. 1} ^{l. 2} ^{l. 3} ^{l. 4} ^{l. 5} ^{l. 6} ^{l. 7} ^{l. 8} ^{l. 9} ^{l. 10} ^{l. 11} ^{l. 12} ^{l. 13} ^{l. 14} ^{l. 15} ^{l. 16} ^{l. 17} ^{l. 18} ^{l. 19} ^{l. 20} ^{l. 21} ^{l. 22} ^{l. 23} ^{l. 24} ^{l. 25} ^{l. 26} ^{l. 27} ^{l. 28} ^{l. 29} ^{l. 30} ^{l. 31} ^{l. 32} ^{l. 33} ^{l. 34} ^{l. 35} ^{l. 36} ^{l. 37} ^{l. 38} ^{l. 39} ^{l. 40} ^{l. 41} ^{l. 42} ^{l. 43} ^{l. 44} ^{l. 45} ^{l. 46} ^{l. 47} ^{l. 48} ^{l. 49} ^{l. 50} ^{l. 51} ^{l. 52} ^{l. 53} ^{l. 54} ^{l. 55} ^{l. 56} ^{l. 57} ^{l. 58} ^{l. 59} ^{l. 60} ^{l. 61} ^{l. 62} ^{l. 63} ^{l. 64} ^{l. 65} ^{l. 66} ^{l. 67} ^{l. 68} ^{l. 69} ^{l. 70} ^{l. 71} ^{l. 72} ^{l. 73} ^{l. 74} ^{l. 75} ^{l. 76} ^{l. 77} ^{l. 78} ^{l. 79} ^{l. 80} ^{l. 81} ^{l. 82} ^{l. 83} ^{l. 84} ^{l. 85} ^{l. 86} ^{l. 87} ^{l. 88} ^{l. 89} ^{l. 90} ^{l. 91} ^{l. 92} ^{l. 93} ^{l. 94} ^{l. 95} ^{l. 96} ^{l. 97} ^{l. 98} ^{l. 99} ^{l. 100} ^{l. 101} ^{l. 102} ^{l. 103} ^{l. 104} ^{l. 105} ^{l. 106} ^{l. 107} ^{l. 108} ^{l. 109} ^{l. 110} ^{l. 111} ^{l. 112} ^{l. 113} ^{l. 114} ^{l. 115} ^{l. 116} ^{l. 117} ^{l. 118} ^{l. 119} ^{l. 120} ^{l. 121} ^{l. 122} ^{l. 123} ^{l. 124} ^{l. 125} ^{l. 126} ^{l. 127} ^{l. 128} ^{l. 129} ^{l. 130} ^{l. 131} ^{l. 132} ^{l. 133} ^{l. 134} ^{l. 135} ^{l. 136} ^{l. 137} ^{l. 138} ^{l. 139} ^{l. 140} ^{l. 141} ^{l. 142} ^{l. 143} ^{l. 144} ^{l. 145} ^{l. 146} ^{l. 147} ^{l. 148} ^{l. 149} ^{l. 150} ^{l. 151} ^{l. 152} ^{l. 153} ^{l. 154} ^{l. 155} ^{l. 156} ^{l. 157} ^{l. 158} ^{l. 159} ^{l. 160} ^{l. 161} ^{l. 162} ^{l. 163} ^{l. 164} ^{l. 165} ^{l. 166} ^{l. 167} ^{l. 168} ^{l. 169} ^{l. 170} ^{l. 171} ^{l. 172} ^{l. 173} ^{l. 174} ^{l. 175} ^{l. 176} ^{l. 177} ^{l. 178} ^{l. 179} ^{l. 180} ^{l. 181} ^{l. 182} ^{l. 183} ^{l. 184} ^{l. 185} ^{l. 186} ^{l. 187} ^{l. 188} ^{l. 189} ^{l. 190} ^{l. 191} ^{l. 192} ^{l. 193} ^{l. 194} ^{l. 195} ^{l. 196} ^{l. 197} ^{l. 198} ^{l. 199} ^{l. 200} ^{l. 201} ^{l. 202} ^{l. 203} ^{l. 204} ^{l. 205} ^{l. 206} ^{l. 207} ^{l. 208} ^{l. 209} ^{l. 210} ^{l. 211} ^{l. 212} ^{l. 213} ^{l. 214} ^{l. 215} ^{l. 216} ^{l. 217} ^{l. 218} ^{l. 219} ^{l. 220} ^{l. 221} ^{l. 222} ^{l. 223} ^{l. 224} ^{l. 225} ^{l. 226} ^{l. 227} ^{l. 228} ^{l. 229} ^{l. 230} ^{l. 231} ^{l. 232} ^{l. 233} ^{l. 234} ^{l. 235} ^{l. 236} ^{l. 237} ^{l. 238} ^{l. 239} ^{l. 240} ^{l. 241} ^{l. 242} ^{l. 243} ^{l. 244} ^{l. 245} ^{l. 246} ^{l. 247} ^{l. 248} ^{l. 249} ^{l. 250} ^{l. 251} ^{l. 252} ^{l. 253} ^{l. 254} ^{l. 255} ^{l. 256} ^{l. 257} ^{l. 258} ^{l. 259} ^{l. 260} ^{l. 261} ^{l. 262} ^{l. 263} ^{l. 264} ^{l. 265} ^{l. 266} ^{l. 267} ^{l. 268} ^{l. 269} ^{l. 270} ^{l. 271} ^{l. 272} ^{l. 273} ^{l. 274} ^{l. 275} ^{l. 276} ^{l. 277} ^{l. 278} ^{l. 279} ^{l. 280} ^{l. 281} ^{l. 282} ^{l. 283} ^{l. 284} ^{l. 285} ^{l. 286} ^{l. 287} ^{l. 288} ^{l. 289} ^{l. 290} ^{l. 291} ^{l. 292} ^{l. 293} ^{l. 294} ^{l. 295} ^{l. 296} ^{l. 297} ^{l. 298} ^{l. 299} ^{l. 300} ^{l. 301} ^{l. 302} ^{l. 303} ^{l. 304} ^{l. 305} ^{l. 306} ^{l. 307} ^{l. 308} ^{l. 309} ^{l. 310} ^{l. 311} ^{l. 312} ^{l. 313} ^{l. 314} ^{l. 315} ^{l. 316} ^{l. 317} ^{l. 318} ^{l. 319} ^{l. 320} ^{l. 321} ^{l. 322} ^{l. 323} ^{l. 324} ^{l. 325} ^{l. 326} ^{l. 327} ^{l. 328} ^{l. 329} ^{l. 330} ^{l. 331} ^{l. 332} ^{l. 333} ^{l. 334} ^{l. 335} ^{l. 336} ^{l. 337} ^{l. 338} ^{l. 339} ^{l. 340} ^{l. 341} ^{l. 342} ^{l. 343} ^{l. 344} ^{l. 345} ^{l. 346} ^{l.}

Seet. III Lotea Medik. phil. bot. 1. p. 204

Ob. Flores ^{subumbellati.} ^{See the Z...}
Cassia 10/10/10. Legumen compressum longum

2. *Lotus ornithogalifolius* L. p. pl. ed. II p. 1091. Vj. 1472. del. p. 40. Hort. fl. austr. II p. 380.

Nitb. fl. ex. p. 505. Alsh. fl. jard. p. 175. Pet. bot. univ. n. 574

foliis rhombic-obovatis, stipulis rhombic-ovatis subcoriatisque, pediculis brevibus
 quibusque foliis floribus calycis subaequantibus, leguminibus linearibus, torulis
 nervis.

6

Lotus parviflorus Desf. fl. atl. II. p. 206. Moris fl. cord. I. p. 513. Jun. syn. fl.

ii. II. p. 357.

L. pilis patulis villosus, foliis cuneato-obovatis oblongis, stipulis ovatis pedunculatis
bi-sexfloris, dentibus calycinis linearibus tubo longioribus, leguminibus oblongis rectis
calyce vix longioribus

Syn. *Lorymenum parviflorum* Ser. in St. petr. II. p. 208 — Non. Desf. l. c. t. 211. DC.

ii. fl. yell. var. t. 30.

Hab in apris montibus Mojubi Colladus dicti prope ^{Lomina} ~~desplum~~ ^{desplum} ~~Vythia~~, und
 communicavit cl. Boneracini. Fl. flavi exsertione coeruleo-iridescentis.

Obs. ^{vera} ~~forma~~ ^{forma} ~~maxima~~ *L. edulis*, * qui differt pedunculatis subreflexis, et cordatis flavis
calyce longioribus, vexillis carinatis ^{repentibus}
 et leguminibus ^{curvis, a longe calyce} ~~rectis~~ ^{supra} ~~longioribus~~

Non. Carn. ic. p. 163. Pin. tetr. m. t. 75.

Herb. ad viciis circa Zam, in saxosis agris circa Vini, & Helot, in praevenis
circa Regis, et in insulae Oper. fl. lutei. fl. vere. 0.

Sect. IV. Eulotus Ser. in D. y. m. II. p. 210.

Herb. conjugati. Carina robusta. Legumen longum.

2. Lohus corniculatus L. p. p. ed. II. p. 1092. Vif. herb. d. m. 40. Hst. fl. austr. II. p. 381. Jedd.
fl. ex. p. 506. Alch. fl. j. d. p. 175. Ped. bot. w. n. 370.

L. foliolis semibovatis terminalibus, ovatis, stipulis oblique subordatis - ovatis, ~~perennibus~~
~~perennibus~~ semi-erectis, dentibus calycis et lobi truncatis subulatis, ~~et~~
~~et~~ leguminibus rectis.

Non. Low. engl. bot. t. 2090. Ped. fl. dan. t. 991. Hst. p. 373 f. 2. Zambel.
1st. plant. ven. t. 280.

Var. p. ulatius: foliolis stipulis calycibusque ulatis -

Var. p. hirsutus: ~~basibus~~ ^{calycibus} ~~ulatis~~ ^{ulatis} - Syn. L. corniculatus p. ~~conspicua~~ ^{conspicua} Desf. p. 111.
354. Vif. herb. d. m. 40.

Var. p. tenuifolius: foliolis stipulis, alisque corollae angustioribus - Syn. L. tenuifolius
Desf. p. 506. L. corniculatus p. ~~tenuifolius~~ L. t. c. -

Herb. in juncis ad viciis in tota Bohemia frequent. fl. lutei, corollae undulatae
ulatis: fl. vere in autumno 4.

4. Lohus cythreus L. p. p. ed. II. p. 1092. Jedd. fl. ex. p. 505. Alch. fl. j. d. p. 175. Ped. bot.
w. n. 371. Koch syn. p. 12. II. p. 196.

L. foliolis adpressis campensibus, foliolis oblongis - ovatis, stipulis ovatis, perennibus brevibus
perforatis, dentibus calycis oblongis lanceolatis, leguminibus subincurvis.

Syn. L. peregrinus Hst. fl. austr. II. p. 380 ex herb. Potenschl. t. non L. - Non

Allium. fl. j. d. t. 11 f. 1. Barcl. ic. stirp. t. 1031 - Buxb. cent. v. append. f. 24.

Herb. ad viciis et saxe montibus circa Vini et Regis, et in insulae Oper. et
Carole. fl. fl. fl. vere. 4.

Obs. Haec agnoscitur est L. cretensis L. qui statim agnoscitur est non ^{argentea et} ~~argentea et~~ ^{perenne} ~~perenne~~.

Lohus angustifolius L. p. p. ed. II. p. 1099. Hst. fl. austr. II. p. 380. Moris fl. rand.

II. p. 502. Koch syn. p. 12. II. p. 198.

L. foliolis parvis villosis, foliolis cuneatis - ovatis, stipulis ovatis, perennibus brevibus
flos, dentibus calycis ^{longioribus} ~~longioribus~~ - subulatis, ^{leguminibus} ~~leguminibus~~ ^{longioribus} ~~longioribus~~.

in L. hirsutus Desf. fl. at. h. p. 190 - L. gracilis Walst. et Kit. fl. v. d. m. 40.

III. f. 254. L. diffusus Sm. fl. bot. p. 494. - Non. Walst. et Kit. t. c. t.

229. Loisel. fl. gall. II. t. 16. Low. engl. bot. t. 925 (communis pubes) Moris. fl. p. 1.

Herb. II. t. 18 f. 1.

6 in saxosis agris insulae Lipa ubi invenit et Potenschl. in capis herbis vici.
fl. lutei. fl. aestate. 0.

II
Sect. III *Woeberia* Moench. *meth. p. 143*

Plorif. subsolitarii. Carina rotata. Legumen uncinatum trigonum longum
Lotus edulis L. sp. pl. ed. II. p. 1090. *Rehb. fl. ex. p. 506. Pet. bot. weing. n. 572*

*L. pilis paucis pubescentibus, foliis pinnatis oblongis, stipulis subcordatis ovatis, pinnis unis-
pinnatis dimidiatis calycibus oblongis lanceolatis, leguminibus carinis duabus
connatis semibilocularibus.*

*Don. laron. icon. pl. II. t. 157. Pers. Alp. pl. exot. t. 268. Moench. hyst. p. sect. II. t. 18
p. 5. et in unguis saporis, vixit. Carina*

*Hab. in agris ad S. Stephanum prope Salatem, et circa Paganum fl. albi, carinae
apice nigro. fl. vireo.*
Obs. hanc carinam calyx subaequantem exornantem vixit.

Sect. V. *Dorycnium* Dur. *instit. rei herb. p. 391 t. 211.*

Plorif. umbellati. Carina obliqua. Legumen rectum trigonum calyx subaequantem
Lotus Dorycnium L. sp. pl. ed. II. p. 1093. *Moench. fl. sil. II. p. 303. capitulis apiculatis*
L. suffruticosum adpressis pubescentibus, foliis stipulisque angustis, cuneatis spatulatis, leguminibus
hypocrateribus glabris calyce subaequale longioribus

Syn. Dorycnium suffruticosum Will. hyst. de gl. de Saug. h. III. p. 16. D. pubescens
Pers. syn. pl. II. p. 354. Vif. Hyst. de Saug. p. 33. D. monspeliense Will. p. 1. III. p.

1096. Hyst. fl. austr. II. p. 382. D. pentaphyllum Rehb. fl. ex. p. 507 exot. syn.
Will. - Icon. Herb. bot. Hyst. p. 100 f. 1. Moench. hyst. p. sect. II. t. 18 f. 1. Lot. in. D. p. 81 f. 1.

*Hab. in praetibus, saxosis circa Paganum et in rupibus Carinae fl. albi, carinae apice
obscurioribus. fl. vireo.*

*Obs. Synonyma et nomina a Linnaeo huius plantae adscripta hanc speciem respiciunt,
cum referantur referri possunt.*

Lotus Crankii Vif. capitulis pinnatis
*L. suffruticosum foliis longis villis pubescentibus, stipulis oblongis cuneatis, leguminibus
curvatis glabris calyce subaequale longioribus*

Syn. Lotus Dorycnium Crank. Hyst. austr. v. p. 402. non L. Dorycnium herbaceum
Will. hyst. de gl. de Saug. h. III. p. 417. Hyst. fl. austr. II. p. 382. Al. Hyst. fl. austr. p. 65
Pet. bot. weing. n. 346. D. sabaudum Rehb. fl. ex. p. 507. - Icon. Will. l. c.

+ 214. Moench. hyst. p. sect. II. t. 18 f. 14.
Vas. argentea: pilis erectis longis uncinatis. vix in specibus multibus circa Dorycnium
Hab. ad marginem pratorum circa Carinam, et in monte Dickers. fl. albi, carinae apice nigro.
fl. aestate.

*Obs. - speciem nimis affinem, et non potest differre: rectius fructum cum L. ibidem, et
L. latifolium in unam conjugendam.*

Sect. VI *Borjeana* Rehb. fl. ex. p. 507

Plorif. subumbellati. Carina obliqua. Legumen oblongum interius rectius brevius divisi-

Lotus hispidus L. p. pl. D. II. p. 1091. Vif. thys. del. p. 40. Host fl. austr. II. p. 381.

Alsch. fl. par. p. 175. Pet. bot. voy. n. 573.

L. tomentosa - *Gilliesii*, foliis stipulaceis cuneatis - oblongis obtusis, leguminibus oblongis turpiter glabris, calyce subaequalibus.

Syn. Bonjeana hispida Phil. t. c. *Dorycnium hispidum* Ser. in DC. prod. II. p. 203.

— Non. Aust. bot. voy. t. 336. *Syn. L.* n. 176.

Subtrib. IV. *Galegeae* DC. prod. II. p. 243.

Alsch. compta glabra. Stamina didypha ~~filamentis~~ ^{filamentis} aequalibus. Legum. unilocularia retusis glabris. Filia uni-multipara impari pinnata.

Bonjeana ~~compta~~

Bonjeana *Ny.*

Host. luyd. bot. p. 372.

Char. Calyx irregulariter quinquefidus, dente infimo longiori. Corolla tetrapetala, albis, glabris, recurvis coracae superae longiorum connatis, denum. ovallibus longis. Stamina a petalis libris, filamentis ovallibus apicibus superae cartilagineae adnatis. Stylus filiformis, stigmati terminali capitulo. Legumen coracae inclusum, uniloculare intrinsecum — *Alsch. compta* DC. *Decand. Syst. ex.*

Bonjeana *compta* Phil. t. c. *Dorycnium* *hispidum* Ser. in DC. prod. II. p. 203. *Phil. t. c.* p. 339. *Alsch. fl. par. p. 175. Host. luyd. bot. p. 372. Vif. thys. del. p. 40.*

canerens, foliis pinnatis trifoliatibus, infimum foliis orbiculatis orbiculatis, superiorum lanceolatis, petalis subulatis, pedunculis axillaribus foliis longioribus, capsula ~~filamentis~~ ^{filamentis} ~~subulatis~~ ^{subulatis} ~~incurvis~~ ^{incurvis} ~~leguminibus~~ ^{leguminibus} ~~oblongis~~ ^{oblongis} ~~obtusis~~ ^{obtusis} ~~glabris~~ ^{glabris} ~~marginibus~~ ^{marginibus} ~~stipulis~~ ^{stipulis}.

Syn. B. plumbosa Phil. fl. par. p. 184. *B. Bonjeana* Vif. thys. del. p. 40. Host fl. austr. II. p. 381. *Alsch. fl. par. p. 175. Host. luyd. bot. p. 372. Vif. thys. del. p. 40.* — *Syn. Junc.* *Sam. en. bot. p. 614. Bot. t. hist. p. 504 f. R. 2. opt.* *Host. vine. II. t. 184.* — *Ny. L.*

Hab. in collibus saxosis agris thym. dolomiticis libralibus. Fl. violaceo-caeruleiscentibus. Fl. caespit. 2.

Alsch. compta ~~adnatis~~ ^{adnatis} ~~filamentis~~ ^{filamentis} ~~subulatis~~ ^{subulatis} ~~incurvis~~ ^{incurvis} ~~leguminibus~~ ^{leguminibus} ~~oblongis~~ ^{oblongis} ~~obtusis~~ ^{obtusis} ~~glabris~~ ^{glabris} ~~marginibus~~ ^{marginibus} ~~stipulis~~ ^{stipulis}. *Alsch. compta* ~~adnatis~~ ^{adnatis} ~~filamentis~~ ^{filamentis} ~~subulatis~~ ^{subulatis} ~~incurvis~~ ^{incurvis} ~~leguminibus~~ ^{leguminibus} ~~oblongis~~ ^{oblongis} ~~obtusis~~ ^{obtusis} ~~glabris~~ ^{glabris} ~~marginibus~~ ^{marginibus} ~~stipulis~~ ^{stipulis}. *Alsch. compta* ~~adnatis~~ ^{adnatis} ~~filamentis~~ ^{filamentis} ~~subulatis~~ ^{subulatis} ~~incurvis~~ ^{incurvis} ~~leguminibus~~ ^{leguminibus} ~~oblongis~~ ^{oblongis} ~~obtusis~~ ^{obtusis} ~~glabris~~ ^{glabris} ~~marginibus~~ ^{marginibus} ~~stipulis~~ ^{stipulis}.

Glycyrrhiza *Dum.*

Host. vine. p. 339. t. 210.

Char. Calyx quinquefidus, dentibus longioribus superioribus ad medium connatis. Corolla pentapetala, albis, glabris, recurvis coracae recte acutae. Stamina didypha. Stylus filiformis, stigmati terminali capitulo. Legumen oblongum adnatis coracae inclusum 1-4 loculare. — *Alsch. compta* DC. *Decand. Syst. ex.*

Antrozalus L.

Gen. pl. ed. VI p. 385 ex. p. plur.

Sen. gl. ed. VI p. 385. excl. p. plur.
Chae. Calyx quinquepartitus, dentibus superioribus unidentatis. Corolla tubulosa, carina
obtusius nuda. — Cesp. radice per an. decem ann. l'yst. n. —
Rehb. H. ex. p. 509.

Sent. I. Hypoglossidei ~~St. prod. syst. nat. II. p. 281~~

Radix persicif. Stipulis inter / comitibus a pectus libere. Flors procreantibus,
calycibus immutatis.

Aphrodesia Onchophora L. p. pl. ed. II p. 1070. Hist. Nat. p. II p. 358. Pl. b. fl. ex. 510

A. adnata pilulosa, stipulis repentinis comatis oppositis, foliis linear-lanceolatis
 sub 12-jugis, spinis ^{inermis} ~~longis~~ ^{longis}, pedunculis foliis longioribus, vix
 linear-oblongis truncatis alioq. dyale. longior, leguminibus orb-oblongis trijugis
~~adnatis~~ ^{hirsutis} ^{L. var}
 a tenuissimis Ait. h. h. var. 73 non des. A. linearis ^{foliis} ^{longis} ^{des.}

~~Asperula~~ *Asperula* a temperary Alt. h. h. m. p. 13 non def. A. Linnaeus, p. 13
Syn A. leontine Alsch. H. J. p. 178 ex loc. non Wulf. — Non Gay fl. austr. 2.
1. re. Rost. bot. may. t. 2665! Chus. h. p. II. p. CCXXXVIII.

Hab. in denticulatis hirsutibus montium vallis bith. Flor. aestate. 7.
H. denticulatis hirsutibus montium vallis bith. Flor. aestate. 7.

Hab. in densest rocky mountainous mountainous vegetation.
Hb. A singular branching leafy plant with distinctively

Asperula purpurea Lam. enc. bot. I p. 214. fl. bl. ex. p. 509.

Ac. umbelliferis, thymis omnibus essentis oppositifolia, foliis ovatis ovatoque, infirmis
acute emarginatis - lobatis, subdecompositis, spicis capitatis, perennantibus foliis longi-
oribus, nervis oblongo - ovatis acue lobato alij sequitiorum, leptomorphis subdant

Syn. G. leonardus Pilsb. *ex loc. non Wolff.* - *Leon H. astrag. t. 12.*

Dec. 21. 1872. - Soci. Pure Sci.
Hab. in saxos. agris. munit. Bickon sub jago Weglow fl. apr. 2.

Ob. Spirit a graduato et ab A. contin. Vulp. Cynomyx sp. et filis auct.
brutis.

Sect. II Dimidiiflora. *St. prod. syst. nat. II. p. 284.*

*Radix peruviana. Stipulae a petiolo et inter se libere. Flores purpurei,
semitae laxae spirales, calycibus immixtis.*

Astragalus Müllerii Steud. & Hochst. in fl. od. bot. zeit 1827 p. 72

A. cinereus, cinereus, stipulis latis, foliis oblongis breviter ~~cuneatis~~ ^{obtusatis}
5-7 jugis, spicis oblongis, laxis, pedunculis fl. longioribus, bracteis ^{paucis} pedicellis
~~bracteis~~ longioribus, calycibus ovatis, adnatis, vexillis coracis integris albis
leguminibus ^{stipulis} ~~leguminibus~~ velis minis calyce dyle longioribus
Legumina leguminibus ^{stipulis} ~~leguminibus~~ velis minis calyce dyle longioribus
L. 236. 1820 T.B. 2006. 1. 18. April 16

Syn. A. argenteus Benth. in *Vij. fl. rar.* in bot. Zeit 1829. T. B. eximib. p. 18. Alch. fl. ¹⁸⁰⁸
Reichb. fl. exc. p. 510. ¹⁸⁰⁸ *quad. fl. delon*
jacq. p. 178. *Bot. bot. weyn.* n. 192. (4) *virgatus* *Reichb. fl. exc.* p. 510. ¹⁸⁰⁸ *quad. fl. delon*
A. argenteus Benth.
non Pall. et *excl. syn. gem.* — *Reichb. fl. delon.* t. XLVI.

Hub. in saxosis apicis circa Zarz, ^{Salonia} Sabinus, Scardina, Imu, in monte Moser pign

Spilato, Memedova, prae Krainsko, Vjsoznica et Beljak, prae Lijn, nec non in
invalij Onje, Chorf, Orso, Oglic et Levina fl. virens violaceo, alij
alij fl. vire. ~~et~~ fl.

[illegible]

Obs. Similitudine A. virgato Pall. qui differt ex parte rubre axium speciei et calycis
 et communiter calyce parvulo longioribus. Quod in nigra diffinitur hothm
 vestiente albo petala longa: Quibus reliquis diffinitur ~~1860~~ ¹⁸⁶⁰ ~~1860~~ ¹⁸⁶⁰
 diffinitionis ut ab hothm videri videtur a virgato et specimen
 hothm videri est. Long et breviter notum p. p. p. p. videri, dem illa
 et hothm ex Arcturum proculat carilium, plerum latitudinem,
 et specis multitudine longioribus aliam plantam p. p. p. p.

Sect. III. Vesicarii D.C. l. c. p. 288

Paris perennif. - Stipulae a petiolo et inter se libere. Flor. purpureae,
 aut albid., calycibus veisicis. exl. 1911. Sact. et Kunt.
 Bot. II. meth. II. p. 358. Schb. Mexic. 510.

Astragalus deflexus L. 17. fl. ed. II. p. 1071. Hort. fl. austr. II. p. 358. f. 68. fl. ex. p. 510.

A diffusis jili centro affinis adpresso nervis, foliis oblongis 3-4 l. p. 7
capitulis oblongis, pedunculis folii brevibus pedicellis longioribus, albugineis
oblongis perib. alba patula in apice inflatis longioribus.
semilocularibus calyce ~~inflato~~ longioribus.

— Leon Waldb. et Rie

Syn. as albidus Welst. et Kt. fl. rar. hung. 1. p. 39. — Leon. Welst. et Kt.
l.c. t. 40. Cart. bot. mag. t. 32 18. Allion. fl. ped. t. 80. f. Drill. fl. du Jura p. 3. t. 2. f. 1.
l.c. Osmat. ut legit. prop. Clementi. fl. violacii.

P. t. 40. Cart. bot. maj. f. 32 v. n. 1. p. 1.
Hob. in saxis apricis montij Ghuyat ubi legit prof. Clementi. Fl. violacei.
Fl. maj. Lemn. t. 1. post specimen authenticum
Obs. Astragalus pastellianus Pollin. hort et pro verm. fl. nov. p. 19. non differt
ab haec non floribus luteis, et leguminibus calyce Ruelli brevitate. Lactes
planta nostra creditur grandior et foliis recte majoribus ut in iconis
~~Athenae~~ et Kibichiana vel omni parte minor ut in illis Athenae et
Villarsii, in quibus tamen calyx male purpureo cyathiformis et nervis brevif.

Sept. IV. Lesamei H. l. c. p. 287.

laris annua. Stipulae a petiolo et inter se liberae. Flos perispermatus et
affixi dense capitati aut spirali, calycibus immixtis.

Artemisia serotina L. 4th ed. p. 166. Bot. fl. accepr. t. y. 358. Herb. Mex. p. 316.
 descr. fl. par. p. 168. Bot. cur. n. 137.

A diff^{us} villis, foliis oblongis, subrecinjis, capitulis ^{sub}sertatis axillaribus, lymenibus
stellatis capitulis ^{terrib} lanceolatis ~~et~~ mucinis, hapteris, dorso canaliculatis.

Non. Jans. hyst. des pl. d'Aix. n. 12. Pluchea. alm. t. 79 p. 3. fr. 10. terr. irr. t. 104 (alluv.)
Columm. ceptiv. 1. p. 304 p. 3.

Herb. in aprig. anis. circa Zan, Khivir, Iracu, Spalato, Ragusa, etiam in montib.
Velle Rha, Shijot. Flor. caeruleis, et albis. fruct. flavis. fl. vere. 0

Sect. V. *Bucrates* M. l. c. p. 290.
vel *intusinae* *hemiconus*

Sect. V. Baccatae Pl. l. c. p. 290.
 vil. infusae semiconnatae
 Jedis annua. Spulac a petalo et calice a lobis. Hony ~~perigonium~~ calycis
~~laxi immutabilis~~ Lychnis arvensis.

Asplenium harveyi L. p. h. d. II p. 167. Root fl. auct. II p. 357. Rhiz. fl. em. 5th. Alud
fl. juv. p. 178. Rh. bot. near n 134. Rh. sp. fl. germ. d II p. 206.
o long. - uncut. - em. - uncut. - 1 bit. - uncut. - about 1. spec.

A adnatum pubescens, foliis subrotundis, supra glabris, infra vix ciliatis, nervis oblongis, spec
subcylindricis, latius pedunculis, folio brevioribus, lepidibus tenuibus, hirsutis
subtus ~~in~~ ^{ad} ~~causam~~ ^{causam} ~~corp~~ ^{corp} ~~canaliculatis~~ ^{canaliculatis} -

J. Lam. ^{leafy} ~~fruit~~ ^{accompanying} ~~fruit~~ 22 f. 4.
Don ^{fruit} ~~fruit~~ ^{accompanying} ~~fruit~~ 105. *Clus.* ^{fruit} ~~fruit~~ ^{accompanying} ~~fruit~~ 11 p 234 f. 2. *Labet.* ^{fruit} ~~fruit~~ ^{accompanying} ~~fruit~~ 11 p 234 f. 2.

Hab. in apris ^{siccissimis}, herbis cum rariis spinis, Fraxin., palato, Vexone, Jagers, et in
infusis Charphi, Bylex, ^{Marye} et Linnæ. Fl. ann. ©

Sept. VI. Galeiformes St. f. c. p. 294.

Parix perennif. Stipulae a petiolo et inter se libere. Flor. ochroleucis,
salsitruis immixtis. Legumina in calyce hirsuta.

Asplenium glycyphyllos L. sp. pl. ed. 2. p. 1067. Hort. fl. austr. 2. p. 257. Pebb. fl. ex. p.
511. ~~Black fl. ed.~~ ^{Red. br. waving.} n. 133. ^{invol. ovate-oblong. ped. pedunc.}

A. glabripes, a slender elegantly 5-7-jugit, spiril ovate-oblong, rarely pedunculate, folis brevibus, liguamentis linearibus subnatis aruatis, glabris erectis, duceis communibus, dorso centraliter.

Acron - low. crypt. bot. t. 203. *Prun.* ter. inv. t. 103. *Sedum.* fruct. p. 548. -

Hab. in sylvestribus et pascuis circa Verulam, in Hunn, ad Ogoje, prope Regum,
ad Regum et Veterano Junno. Fl. aestiv. 7.

Sect. VII. *Caprini* ~~Leob. ⁱⁿ f. ⁱⁿ f.~~ *Spic. fl. rum. 1. p. 59*

Sed. VII. *Inverni*
Spix ferrug. *Apyulae* *pallida*, *adnatae*. *Alcyon* *inimulati*

Astragalus illyricus Bernh. sel. sem. h. Exsicc. 1836.

A. ^{pubescens, a. n. sp.} ~~pubescens~~, *Apuleia lanceolata* - lanceolata, *pubes.* 12-20-jajif. *light*, *very* *dark* - *oblong*.

~~In nostra sentia categoria~~ ~~gady~~ ~~layung~~ ~~et~~ ~~ayuphony~~ ~~de~~ ~~in~~ ~~planta~~ ~~no~~
A. mantropi / Et hic A. Inguanthe. ^{non} Denique tropicis ayupis acutis ut in n. 11
de altitudinis arboribus quibus vides in quibus a Mantropi acceptis, et
ayupis Lamarkii et Caudellii optime representat. In nostra planta pte ubique
et confusis semine. canescendo, ut in planta a Lamarkii dicitur, et in
specimen Douroepithio ab ipso et Caudellio citato.

Ind. III. Hedy sarcae

си. си

Prunus dioica - *Lepidum transectum* in *lenticularis*, not in *artificialis* more
spermy seedling. Embryo curved, cylindrical in germination, plicately epigeal.

Subm B. I. Cornilleau D.C. p. 2. II. p. 308.

Leptocarpus teretica aut *crispus*. Hox umbellat., aut solitaria.

Scopians L.

Gen. pl. ed. II. p. 278.

Char. Calyx breviter campanulatus, quinquefidus, dentibus apicibus
ultra medium saepe. Corolla tubulosa, caeva lobi 5-partiti,
apice recurvato. Stamina 5-loba, filamento altero
apice 2-loba. Ligamen amentum setosum brevius
amentum 2-lobum saepe. Caps. 2-loba. Ov. 2-loba. 1-loba.

Temnodactylus subulley L. f. p. 2. 11 p. 1054. ~~1841~~ ~~1842~~ ~~1843~~ ~~1844~~ ~~1845~~ ~~1846~~ ~~1847~~ ~~1848~~ ~~1849~~ ~~1850~~ ~~1851~~ ~~1852~~ ~~1853~~ ~~1854~~ ~~1855~~ ~~1856~~ ~~1857~~ ~~1858~~ ~~1859~~ ~~1860~~ ~~1861~~ ~~1862~~ ~~1863~~ ~~1864~~ ~~1865~~ ~~1866~~ ~~1867~~ ~~1868~~ ~~1869~~ ~~1870~~ ~~1871~~ ~~1872~~ ~~1873~~ ~~1874~~ ~~1875~~ ~~1876~~ ~~1877~~ ~~1878~~ ~~1879~~ ~~1880~~ ~~1881~~ ~~1882~~ ~~1883~~ ~~1884~~ ~~1885~~ ~~1886~~ ~~1887~~ ~~1888~~ ~~1889~~ ~~1890~~ ~~1891~~ ~~1892~~ ~~1893~~ ~~1894~~ ~~1895~~ ~~1896~~ ~~1897~~ ~~1898~~ ~~1899~~ ~~1900~~ ~~1901~~ ~~1902~~ ~~1903~~ ~~1904~~ ~~1905~~ ~~1906~~ ~~1907~~ ~~1908~~ ~~1909~~ ~~1910~~ ~~1911~~ ~~1912~~ ~~1913~~ ~~1914~~ ~~1915~~ ~~1916~~ ~~1917~~ ~~1918~~ ~~1919~~ ~~1920~~ ~~1921~~ ~~1922~~ ~~1923~~ ~~1924~~ ~~1925~~ ~~1926~~ ~~1927~~ ~~1928~~ ~~1929~~ ~~1930~~ ~~1931~~ ~~1932~~ ~~1933~~ ~~1934~~ ~~1935~~ ~~1936~~ ~~1937~~ ~~1938~~ ~~1939~~ ~~1940~~ ~~1941~~ ~~1942~~ ~~1943~~ ~~1944~~ ~~1945~~ ~~1946~~ ~~1947~~ ~~1948~~ ~~1949~~ ~~1950~~ ~~1951~~ ~~1952~~ ~~1953~~ ~~1954~~ ~~1955~~ ~~1956~~ ~~1957~~ ~~1958~~ ~~1959~~ ~~1960~~ ~~1961~~ ~~1962~~ ~~1963~~ ~~1964~~ ~~1965~~ ~~1966~~ ~~1967~~ ~~1968~~ ~~1969~~ ~~1970~~ ~~1971~~ ~~1972~~ ~~1973~~ ~~1974~~ ~~1975~~ ~~1976~~ ~~1977~~ ~~1978~~ ~~1979~~ ~~1980~~ ~~1981~~ ~~1982~~ ~~1983~~ ~~1984~~ ~~1985~~ ~~1986~~ ~~1987~~ ~~1988~~ ~~1989~~ ~~1990~~ ~~1991~~ ~~1992~~ ~~1993~~ ~~1994~~ ~~1995~~ ~~1996~~ ~~1997~~ ~~1998~~ ~~1999~~ ~~2000~~ ~~2001~~ ~~2002~~ ~~2003~~ ~~2004~~ ~~2005~~ ~~2006~~ ~~2007~~ ~~2008~~ ~~2009~~ ~~2010~~ ~~2011~~ ~~2012~~ ~~2013~~ ~~2014~~ ~~2015~~ ~~2016~~ ~~2017~~ ~~2018~~ ~~2019~~ ~~2020~~ ~~2021~~ ~~2022~~ ~~2023~~ ~~2024~~ ~~2025~~ ~~2026~~ ~~2027~~ ~~2028~~ ~~2029~~ ~~2030~~ ~~2031~~ ~~2032~~ ~~2033~~ ~~2034~~ ~~2035~~ ~~2036~~ ~~2037~~ ~~2038~~ ~~2039~~ ~~2040~~ ~~2041~~ ~~2042~~ ~~2043~~ ~~2044~~ ~~2045~~ ~~2046~~ ~~2047~~ ~~2048~~ ~~2049~~ ~~2050~~ ~~2051~~ ~~2052~~ ~~2053~~ ~~2054~~ ~~2055~~ ~~2056~~ ~~2057~~ ~~2058~~ ~~2059~~ ~~2060~~ ~~2061~~ ~~2062~~ ~~2063~~ ~~2064~~ ~~2065~~ ~~2066~~ ~~2067~~ ~~2068~~ ~~2069~~ ~~2070~~ ~~2071~~ ~~2072~~ ~~2073~~ ~~2074~~ ~~2075~~ ~~2076~~ ~~2077~~ ~~2078~~ ~~2079~~ ~~2080~~ ~~2081~~ ~~2082~~ ~~2083~~ ~~2084~~ ~~2085~~ ~~2086~~ ~~2087~~ ~~2088~~ ~~2089~~ ~~2090~~ ~~2091~~ ~~2092~~ ~~2093~~ ~~2094~~ ~~2095~~ ~~2096~~ ~~2097~~ ~~2098~~ ~~2099~~ ~~2100~~ ~~2101~~ ~~2102~~ ~~2103~~ ~~2104~~ ~~2105~~ ~~2106~~ ~~2107~~ ~~2108~~ ~~2109~~ ~~2110~~ ~~2111~~ ~~2112~~ ~~2113~~ ~~2114~~ ~~2115~~ ~~2116~~ ~~2117~~ ~~2118~~ ~~2119~~ ~~2120~~ ~~2121~~ ~~2122~~ ~~2123~~ ~~2124~~ ~~2125~~ ~~2126~~ ~~2127~~ ~~2128~~ ~~2129~~ ~~2130~~ ~~2131~~ ~~2132~~ ~~2133~~ ~~2134~~ ~~2135~~ ~~2136~~ ~~2137~~ ~~2138~~ ~~2139~~ ~~2140~~ ~~2141~~ ~~2142~~ ~~2143~~ ~~2144~~ ~~2145~~ ~~2146~~ ~~2147~~ ~~2148~~ ~~2149~~ ~~2150~~ ~~2151~~ ~~2152~~ ~~2153~~ ~~2154~~ ~~2155~~ ~~2156~~ ~~2157~~ ~~2158~~ ~~2159~~ ~~2160~~ ~~2161~~ ~~2162~~ ~~2163~~ ~~2164~~ ~~2165~~ ~~2166~~ ~~2167~~ ~~2168~~ ~~2169~~ ~~2170~~ ~~2171~~ ~~2172~~ ~~2173~~ ~~2174~~ ~~2175~~ ~~2176~~ ~~2177~~ ~~2178~~ ~~2179~~ ~~2180~~ ~~2181~~ ~~2182~~ ~~2183~~ ~~2184~~ ~~2185~~ ~~2186~~ ~~2187~~ ~~2188~~ ~~2189~~ ~~2190~~ ~~2191~~ ~~2192~~ ~~2193~~ ~~2194~~ ~~2195~~ ~~2196~~ ~~2197~~ ~~2198~~ ~~2199~~ ~~2200~~ ~~2201~~ ~~2202~~ ~~2203~~ ~~2204~~ ~~2205~~ ~~2206~~ ~~2207~~ ~~2208~~ ~~2209~~ ~~2210~~ ~~2211~~ ~~2212~~ ~~2213~~ ~~2214~~ ~~2215~~ ~~2216~~ ~~2217~~ ~~2218~~ ~~2219~~ ~~2220~~ ~~2221~~ ~~2222~~ ~~2223~~ ~~2224~~ ~~2225~~ ~~2226~~ ~~2227~~ ~~2228~~ ~~2229~~ ~~2230~~ ~~2231~~ ~~2232~~ ~~2233~~ ~~2234~~ ~~2235~~ ~~2236~~ ~~2237~~ ~~2238~~ ~~2239~~ ~~2240~~ ~~2241~~ ~~2242~~ ~~2243~~ ~~2244~~ ~~2245~~ ~~2246~~ <

II. p. 249. *Phob. fl. ex. g.* 541. *Alch. fl. j. g.* 179. *Pitt. bot. warr. n.* 841. —
l. ... 2-5 flori. lagonis

St. John's wort - lanceolate acutif, tomentose, peduncles 2-5 fls, cymose
~~St. John's wort~~ ^{St. John's} *acutif, extremely aculeatis, indivisibly inermis.*

~~Spec. Nisi tot. inv. t. g~~

Non. Arisaema Herb. et fl. anpr. et p. 349. non L. — Non. Arisaema.

Syn. Sc. minutum 1861.
 var. f. 98. *Mont. kept. 11 Oct. II. f. 11. f. 2.* *leaving. Succ. 19 a. fl. vi. II. y. 50.*
 *leaving. Succ. 19 a. fl. vi. II. y. 50.*

Jav. B. andersonii: *lymanii* *diversa* *clavata* *ovata*:
Hab. ad riv. et marginem cynosu in *apice* *citra* *pan* *Abur* *pan* *gala*.
Jav. B. andersonii *diversa* *clavata* *ovata*:
Hab. ad riv. et marginem cynosu in *apice* *citra* *pan* *Abur* *pan* *gala*.

Prof. et in domo Vespini, Orici, Lomani. H. lutea. H. magis. H. ...
... glabris ut luteis, nunc solis coctis domo

Obt. Don't audibly laugh by hunches, frowns, or any
means for any, intently gazing except, or usually.

Cornilla L.

Gen. pl. ad. II. p. 279 excl. 1 per.

Synpl. ad. lt. p. 279 ex. sp.
Chas. calyx breviter campanulatus purpureus, dentibus sagittatis albis median
natis. Corolla lobis ovatis dentat. calycem longioribus, cavitatem accensu-
matis - rostrata. Stamina didyma, filamentis albicantibus apice dilatatis. Sep-
men contractum - articulatione ~~natura~~, articulis minusque — Clap. Didymis
Art. Decandria l'yst. n. x. —
Drosera infl. p. 650 t. 418.

Sech. F. Emery Journ. inst. p. 650. t. 418.

Upper perianth calyx large, long. *Lycium latifolium*, and
occasionally.

Convolvulus *serotinus* L. sp. pl. 2. II. p. 1146. Vif. / Arg. ed. lat. p. 32. Hort. fl. and fr. II. p. 345. fide.
 1/2 562. *Alch. fl. lat.* p. 179. *fl. lat. serotinus* L. 290.

H. exc. p. 542. Alsch. H. jadv. p. 179. Pet. b. w. u. 290.

C. fruticosa erecta ramis angulatis. Infloresc. ~~erect~~ ^{erect} cymes longae, pauciflorae, oviger.

Bonaveria Scop.

Introd. ad hist. nat. p. ~~310~~ ³¹⁰

[illegible]

14th Oct. -
Donaria leucandra ~~Lept.~~ Abb. fl. ex. p. 541

[illegible]

Scen. Sibth. et Sm. fl. graec. VIII t. 212. Lam. enc. bot. gr. Reg. Gaertn. de fr. ca.
 sem. II. t. 153. Char. hisp. gr. II. p. cccxxvi. — Murr. Bot. ussuri.

hab. a tief, ich agist, in großer tiefen gegen taxoffene tiefen Dolanti. H.
am in asphalten. ②

Herbaceous ^{in aspect} ^{offic.} Cornelian ^{prun.} stipulif. leafy, fls. glabrous, fls. 4-5-jugif
obovate acuminate viteris, umbellif. long pedunculat. 5-10 fls. fls.
brif. luteif. ligunda hyssopari long. viteris glab. indurcent. yve a
Cornelian ^{prun.} viteris.

Ornithogalys L.

Gen. pl. ed. II. p. 278.

Gen. pl. ed. 4. p. 288.
 Chor. Calyx tubulosa campanulata, 5-lobata, lobis, rubris, rugulosis, bisp. cordatis.
 Corolla tetragloba, lobis rotundatis, lobis. Stamina diadelph. filamentis, calycis
 apice dilatatis. Gynem longius brevius vixitum compressum contractu-
 sculatum, subulig. reniditibus monospermis. — Chap. Diadelph. Ord. Dica-
 dia. 1946. 1. x. — 1947. — Feb. fl. x. p. 540.

[illegible]

Fritab. spic. fl. rusc. 1. ⁶³ ~~pistilata~~ utrinque ligulata

[illegible]

London

brunellia
Don. Sibth. et Sm. fl. gr. VIII t. 714. Dalech. hist. pl. i. p. 493.
Hab. in arvis hirsutis insula Lacoon prope Bugeia. Fl. parvi lutei. Fr. Agg. Nervis. ©

Hippocrepis L.

Pin. pl. 2. II. p. 28

Char. Calyx coramulatus, quinquefidus, dentibus saepe brevibus 2. et medio
 curvatis. Corolla tetragyna coramulata. ovata. Stamina
 didyma, filamentis oblongis apice dilatatis. Legumen longum
 conflatum articulatione, articulis internis superioribus immixtis
 exasperatis ~~articulis~~ demum nudantibus monospermis. Clap.
 Didymis ord. decandris. Lich. 200.

Hippocrepis corolla L. sp. pl. 2. II. p. 1080. Vif. 1. Kny. delm. p. 37. Herb. fl. europ.
 II. p. 1. Feb. fl. exc. p. 580. Alsch. fl. jud. p. 178. Pet. bot. weig. n. 419.

H. ~~radix~~ caulibus basi suffrutescentibus, parvis umbellatis foliis longioribus
 leguminibus subseratis, dorso flexuosis nix tuberculatis scabris

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Var. ~~glauca~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}
 H. ~~radix~~ ^{glauca} ~~fl. exc. p. 580.~~ ^{fl. exc. p. 580.} ~~Alsch. fl. jud. p. 178.~~ ^{Alsch. fl. jud. p. 178.} ~~Pet. bot. weig. n. 419.~~ ^{Pet. bot. weig. n. 419.}

Syr. V. *Biranea* (Papr. proc. sociol. p. 37) non Verdy. ^{Verdy} *Biranea* Maly en pl. imp. arkt. p. 362. *Bir.* in Sp. 1948. v. 3. p. 267. *Evren argentinum* Juss. cat. sem. h. in Acadif. 1826 p. 4. — Non. *Morif* fl. sand. t. LXX. *carvaca*

Ab. in cultis, horti, incolis Latic. Corolla ^{cambraca} ~~fulva~~ ^{alba}, ovillo caeruleo-striata, carinae
apice atropurpureo. fl. scab. ③

Pl. Hinc rugosa non lanceolata acuta sed p^{er} h^{oc} obliqua et abrupte mucronata, et ^{te recte} ~~non~~ ^{non} ~~mutata~~ sed oblique truncata: h^{oc} quibus divergat ab iis. Florae sordidae
vires in plantis tunc fere tunc olivaceae.

Vicia carnabre L. p. pl. ed. II. p. 1035. Herb. fl. vesp. II. p. 350. Pub. fl. car. 528.
V. flatis oblongis ^{orb-}~~lanceatis~~ ^{lanceatis} ~~oblongis~~ ^{multijugis}, stipulis integerrimis,
^{apiculatis} ~~linearibus~~ ^{brevibus} ~~tortosis~~ ^{floribus} folio brevioribus, pedicellis calycis
~~tubis~~ ^{calycis} ~~subglobosis~~ ^{dubius} ~~calycis~~ ^{apice} ~~retineis~~ ^{tubo} ~~breviter~~ ^{leguminibus}
oblongo-~~rhomboidibus~~ ^{globois}.

Obozo - Shombij glabrij.

Syn. V. *Pseud. Jay*. ft. *austro-It.* g. 16 nov St. V. *nichitang Comp.* 1879. *austro p. 388.*

Eriom. cassidatum Nov. Jay. l.c. t. 229. *Cantk l.c. lib t. f. 2.*

Hab. in glühender sonnenhitze, und communis pfl. Altschuler. Fl. weiss.

29 *Vicia Gerardi* DC. fl. fr. IV. p. 591. ex. syn. Hoch syn. fl. germ. 2. V. p. 213. Borkst. ^{2.} ^{Doyle}
 monog. syn. 4. p. 1. V. p. 1. in Nuov. ann. d. l. nat. Bot. 1847. p. 481. ^{et fl. Ital. VII. p. 481.}
 sub 15-jugis ^{unijugis.} ^{Gerardi (Cav.)}

V. floris oblongis lanceolatis obtusis mucronatis multifloris ^{sub 15-jugis} ^{nerviis} ^{marginibus} ^{concoloribus}
calycis sessilibus longimucronatis sepius multifloris ^{marginibus} ^{multis} ^{vix} ^{obtusius} ^{nerviis} ^{sepius}
calycis tubo brevioribus, distinctis calycinibus exsertentibus brevissimis
^{stati} ^{horum} ^{calycis} ^{tubo} ^{longior.}

V. Candolleana Pers. fl. usq. v. p. III.

Lycium *V. galleprovincialis* Pers. em. bot. mag. C. V. p. 87. *V. Conna* var. *Savardi* Sav.
fl. bulv. IV. f. 506 Grub. sp. fl. vinal. I. p. 82. — Non. *V. galleprovinc.* p. 498.
f. 19.

Hub. in f. 19.
in sylvaticis montium Vellebith und meist prof. Altschungen, et ex Dolomiti
non less speculij ubi ~~habet~~ acceptum upon a cl. prof. Bernharti. fl. caerulea.
fl. caerulea. 24. abuss. p. 81. et fl. mag. v. p. 116. de vrb. fl. imp. VII p. 485.

22. *Vicia* *villosa* *Robt. Hart. fl. german. 2. p. 187. var. p. glaberrima Koch syn. fl. german. ed. II*
p. 216. Berol. locut

[illegible][illegible]

Obt. Vicia polyphylla Waldb. et Kit. fl. v. hung. II. p. t. 254 nec Desf. est eadem ac
V. villosa Roth, et utique vix tantum vixit pube coarctata, longiorum.

23 Vicia Cracca L. p. pl. d. II. p. 1035. Hort. fl. augs. II. p. 331. Fels. fl. ex. p. 528.

V. foliis oblongis lanceolatis mucronatis, stipulis angustis subdeceunjugis, stipulis
remixtis integerrimis, crumibus linearibus, pedunculis densifloris foliis longioribus, pedicellis calycis
tubo brevioribus, dentibus calycis apertis brevibus irregularibus, vexillis ad
medium coarctatis, leguminibus linearibus, stipite calycis subaequalibus.

Sem. long. ang. bot. t. 1168. Sem. et pube. fl. purpureo-roseo. fl. p. k. fl. et axillares.

Hab. in humis montium Major et Brethor. fl. purpureo-roseo. fl. augs. II.

Obt. In vix fl. quidam longior at in speciemibus plantae germanicae, id sine
quodam cum analysi iconis angl. bot. non citat.

24 Vicia tenuifolia Roth tent. fl. germ. I. p. 309 et II. 2. p. 183. Hort. fl. augs. II. p. 332.

Fels. fl. ex. p. 528.

V. foliis lanceolatis linearibus mucronatis, stipulis angustis subdeceunjugis, stipulis
remixtis integerrimis, crumibus linearibus, pedunculis densifloris foliis longioribus, pedicellis
calycis tubo subaequalibus, dentibus calycis apertis brevibus irregularibus, vexillis ad
medium coarctatis, leguminibus linearibus, stipite calycis subaequalibus.

Var. p. laxiflora: floribus laxe racemosis, foliis angustis linearibus - Syn. V. tenuif.
lin. var. laxiflora Griseb. spec. fl. rumel. I. p. 82. - Sem. long. fl. var. cent. V. app.
t. 35.

Var. y. luxurians: floribus imbricatis majusculis, foliis late linearibus purpureis
abrupte mucronatis.

Hab. in locis apricis circa Lissim, et Citharistia in ins. Creta, et circa Nereid.
ad vix apricis in insula Bragg: var. y. pube flava - fl. blavini, alij
pallidioribus. fl. augs. II.

25 Vicia ochroleuca Den. fl. nap. p. 42. Pers. syn. fl. sic. II. p. 290. Fels. bot. univ. p. 1099.

V. foliis oblongis linearibus mucronatis, stipulis angustis subdeceunjugis, stipulis
remixtis integerrimis, crumibus linearibus, pedunculis longis racemosis, pedicellis
calycis tubo brevioribus, dentibus calycis apertis brevibus irregularibus, vexillis ad
medium coarctatis, leguminibus linearibus, stipite calycis subaequalibus.

Sem. Den. fl. nap. t. 174.
Hab. in sylvestribus montibus circa Lissim, et Citharistia in ins. Creta, et circa Nereid.
ad vix apricis in insula Bragg: var. y. pube flava - fl. blavini, alij
pallidioribus. fl. augs. II.

26 Vicia atropurpurea Desf. fl. atl. II. p. 183. Germ. syn. fl. sic. II. p. 289. Pers. fl. ital. VII. p. 493.

V. foliis oblongis linearibus mucronatis, stipulis angustis subdeceunjugis, stipulis
remixtis integerrimis, crumibus linearibus, pedunculis longis racemosis, pedicellis
calycis tubo brevioribus, dentibus calycis apertis brevibus irregularibus, vexillis ad
medium coarctatis, leguminibus linearibus, stipite calycis subaequalibus.

16 *Vicia grandif* Loisel. fl. gall. ed. I. p. 460. Morif. fl. var. I. p. 568

V. flabris linearis-lanceolatis acutis 2-5-jugis, infimis ovatis, cirsidis simplicibus, stipulis semijugis integris cuneatis superius bipinnatis linearibus, pedunculis filiformibus 1-5-floris foliis multis longioribus dentibus calycis lanceolatis cuneatis inaequalibus leguminibus 2-3-jugis bipinnatis, hyeme tub. calycis brevioribus.

Syn. *Ervum grande* DC. cat. h. morif. p. 109. fl. flex. p. 526. E. truncatum

Ref. syn. pl. II. p. 509. E. longispicum Tenor. fl. neap. prod. p. 59. E. tetrapetrum

Alsch. fl. jedr. p. 194 ex herb. nec L. — Non Loisel. fl. gall. hb. XII.

Hab. in saxos. herbis et cirsidis circa lacum Breznovce prope Carn. in monte Tell. supra latior et in rupibus Clus. Open et Loma. Fl. perennans. semet fl. acutis. C.

Obs. Smellina V. tetrapetrum L. Ervum tetrapetrum L. quod differt foliis repandis lobatis ac in illa minus acutis oblique, cirsidis simplicibus, cirsidis, corolla minus exserta.

17 *Vicia hirsuta* Koch syn. fl. germ. ed. I. p. 191

E. flabris linearibus oblique repandis cirsidis vixis, apicibus ovatis, stipulis semijugis lanceolatis, pedunculis filiformibus 2-8-floris multis angustis foliis subrepandis, dentibus calycis lanceolatis inaequalibus leguminibus oblongis-shombrii. Hyeme tub. calycis brevioribus.

Syn. *Ervum hirsutum* L. p. pl. ed. II. p. 1039. Herb. fl. austr. II. p. 337. fl. flex. p. 527.

— Non. Low. engl. bot. t. 990. Des. fl. dan. t. 639. Sprin. tab. inv. t. 50 f. 2. Lechl. II. p. 147. II. p. 76 f. 1.

Var. β *leptocarpa*: leguminibus lobatis, stipulis simplicibus integris linearibus. Morif. fl. p. 515. E. *Derrovii* Ten. fl. neap. prod. app. V. p. 22. E. *pubescens* B. *leptocarpa* Ten. sylt. pl. p. 364. E. *sardum* Spr. syst. reg. cur. post. p. 546. E. *Derrovii* Ait. herb.

Hab. in saxos. herbis, et inter agiles circa Derrovij, Zagorje, Lebnice, Verpog, Istri, Verpog et in deserta Lomae. Fl. ex albidis caulescentibus. fl. ligr. Ten. C.

Obs. Quoad *Derrovium* generum *Vicia*, *Ervum* et *Lathyrus* vide infra post *Lathyrum*.

Lathyrus Wimm.

Fl. fil. II.

Char. Calyx perispermis et perispermatis. Stylus apice unguicatus vel dilatatus. Legumen linearis, oblongum, vel shombrium. — Clasp. Dicotyledonae Ord. Dicotyledonae Syst. II.

Subgen. I. *Ervum* L. gen. pl. ed. II. p. 271.

Corolla calycis subequalis. Legumen shombrium. Folis apice cirsidis.

1 *Lathyrus* Lenc. trif.

L. fl. repandis subrepandis cirsidis filiformibus, stipulis lanceolatis integris, pedunculis filiformibus 1-5-floris foliis subrepandis, leguminibus ovatis-shombrii.

Syn. *Ervum Lens* L. *sp. pl. ed. II* p. 1039. *Ny. shrig. delin. p. 34. fl. ex p. 526*
Alsch. fl. jard. p. 175. Cier Lens W. sp. pl. III p. 1114. - Den. pin. t. h.
iss. t. 35. Matth. in Diof. p. 451. Nees. hyst. shrig. p. 859. - Hlyr.
Lichia - Lögiviza.

Hb. cultis in apic. ob remissa edulia. fl. ex albidis purpureis fl. a. p. t. 0

2. *Lathyrus nigricans* Vis.

L. foliis bi-quadrifidis, cunctis mucroniformi, stipulis semi-statis dentatis, pedunculis aristatis uni-bifloris foliis longioribus, leguminibus quadrato-rhombicis glabris.

Syn. *Ervum nigricans* Sibth. fl. taen. - cauc. II p. 164. *Ny. shrig. delin. p. 34. fl. ex p. 526. Alsch. fl. jard. p. 174. E. lentoides Den. fl. mag. prod. suppl. II p. 68. E. delvaticum Presl p. 27. & Benth. p. 30? - Non Monf. fl. rar. t. XXXI p. 2. Den. fl. rar. t. 167. p. 2.*

Hb. in arvis apicis circa Schinua, Tria, sp. t. et in bursis Open et Livia. fl. purpureo-carmineo, alii albidis. fl. v. 0

Obs. Quid sit *Ervum storionae* L. difficile dictu, nam ex iconibus in amoen. acad. IV. p. 326 huius plantae ascriptis illa *sp. n.* exhibet V. *Lathyrum* L., et non originalis Monfroi unde Linnaeus nomen speciei hanc p. 2. t. 167. p. 2. nunc probabiliter refert. Descriptio vero ejusdem plantae in Amoen. acad. t. c. cum E. nigricante Sibth. bene quadrat.

3. *Lathyrus Leukula* Vis.

L. foliis bi-quadrifidis, cunctis ~~mucroniformi~~ ^{filiformi} vel mucroniformi, ~~leguminibus~~ ^{leguminibus} ~~rhombicis~~ ^{filiformi} ~~stipulis~~ ^{stipulis} ~~semi-statis~~ ^{semi-statis} ~~dentatis~~ ^{dentatis} ~~pedunculis~~ ^{pedunculis} ~~aristatis~~ ^{aristatis} ~~uni-bifloris~~ ^{uni-bifloris} ~~foliis~~ ^{foliis} ~~longioribus~~ ^{longioribus} ~~multis~~ ^{multis} ~~aristatis~~ ^{aristatis} ~~leguminibus~~ ^{leguminibus} ~~obovatis~~ ^{obovatis} ~~rhombicis~~ ^{rhombicis} ~~puberulis~~ ^{puberulis}.

Syn. *Ervum Leukula* Schreb. in Thurm. fl. germ. I fasc. 32. Monf. fl. rar. I p. 573 Koch syn. fl. germ. ed. II p. 220. E. uniflorum Desv. fl. neap. prod. suppl. II p. 68. Schaff. et Maur. fl. rom. prod. p. 248. *Ny. fl. ex p. 526. Pelt. bot. vray. n. 371. Cier ervoides Boryl. fasc. fl. parjul. p. 27. Hb. fl. austr. II. p. 338 - Icon Schaff. et Maur. t. c. tab. v f. 2. Desv. fl. neap. b. 172 f. 2.*

Hb. in saxosis sylvestribus ad Bergolze prope Lubnia, et circa Jagy, nec non in insulis Posman, Cherp et Livia. fl. vixille purpureo-aeeruleo, alii albidis. fl. v. 0

Obs. Speciei hae minus differt a praecedente pedunculis unifloris, qui interdum biflori; aut multis, cum non raro aristati occurrunt; vel foliis superioribus filiformi differt, cum saepius eodem gaudeant; vel stipulis modo bipartitis modo lanceolatis, quam forma leguminis, quae in L. *Leukula* sunt bipartite alternate idque obovata, dum in L. nigricante sunt atropaeae extremitate apiculatae late ita ut signum quadrilaterum fere regularem exhibeant, quod minus bene expressum in rone flor. romane ducit curvatus ex illa fl. neap. p. 2.

— *Penny* ~~hiffe~~ ^{Lathyrus} ~~affinis~~ ^{affinis} et *Cicer* *Dura*, *plaventy* *altany*
stammum *infra* *apicem* *invenit*, *legumina* *tergide* *ex* *spirans* *ovuli* *aut* *sharbo*, *et* *semine*
huj *invenit* *gibbif* *affinchem* *thijura* *pecij* *C. arichnum* *L. paze* *in* *Delan*
tie *elitus* *ob* *semine* *edulis* *sub* *nominis* *ilgira* *Grah.* *Grah.* *Hani.* *Hanick.*

Lathyrus II. *Lathyrus* *L. gen. phed. p. 267.*
Corolla *calycem* *superante* *Legumen* *lineare* *oblongum*. *Hic* *circloph.*

Sect. I. *Aphaca* *Dura* *intus* *herb. p. 6.*

Folia *nulla*, *petalo* *foliiformi* *aut* *circuliformi*.

1. *Lathyrus* *aphaca* *L. gen. phed. p. 109. W. ph. Del. p. 39. Bot. fl. austr. p. 324.*

Hab. fl. ex. p. 533. Alsch. fl. pad. p. 172. Pet. bot. univ. n. 530.

L. phed. p. 109. W. ph. Del. p. 39. Bot. fl. austr. p. 324.
L. phed. p. 109. W. ph. Del. p. 39. Bot. fl. austr. p. 324.

Non. Low. east. bot. t. 116. Lib. i. con. it. p. 70 f. 1. Dod. pempt. p. 545.

Hab. ad *reg. et* *in* *salsis* *thij* *Delan.* *sup. p. 71. fl. austr. p. 324. fl. austr. p. 324.*

Sect. II. *Chimene* *Del. in* *ist. nat. p. 378.*

Petiolis *in* *finibus* *aphyllis*, *superioribus* *foliolatis*, *inferioribus* *filiformi*.

3. *Lathyrus* *Chimene* *Del. fl. fr. IV. p. 578. Pet. bot. univ. n. 533. Alsch. fl. pad. p. 172.*

L. caulibus *bracteatis*, *legumina* *tergide* *ex* *spirans* *ovuli* *aut* *sharbo*, *et* *semine*
huj *invenit* *gibbif* *affinchem* *thijura* *pecij* *C. arichnum* *L. paze* *in* *Delan*
tie *elitus* *ob* *semine* *edulis* *sub* *nominis* *ilgira* *Grah.* *Grah.* *Hani.* *Hanick.*

Syn. Pison. Chimene *L. gen. phed. p. 1027. Bot. fl. austr. p. 321. Chimene* *pet. bot. univ. n. 535.*
fl. austr. p. 321. Bot. fl. austr. p. 321. Chimene *pet. bot. univ. n. 535.*

Hab. in *agris* *et* *in* *salsis* *thij* *Delan.* *sup. p. 71. fl. austr. p. 324. fl. austr. p. 324.*

6. *Lathyrus* *temuifolius* *Desf. fl. atl. II. p. 160. Monf. fl. nord. t. 1. p. 582.*

L. caulibus *bracteatis*, *legumina* *tergide* *ex* *spirans* *ovuli* *aut* *sharbo*, *et* *semine*
huj *invenit* *gibbif* *affinchem* *thijura* *pecij* *C. arichnum* *L. paze* *in* *Delan*
tie *elitus* *ob* *semine* *edulis* *sub* *nominis* *ilgira* *Grah.* *Grah.* *Hani.* *Hanick.*

Syn. L. auriculatus *Desf. fl. atl. II. p. 160. Monf. fl. nord. t. 1. p. 582.*
fl. atl. II. p. 160. Monf. fl. nord. t. 1. p. 582.

Hab. in *agris* *et* *in* *salsis* *thij* *Delan.* *sup. p. 71. fl. austr. p. 324. fl. austr. p. 324.*

Sect. III. *Eulathyrus* *Desf. in* *Del. pad. p. 367.*

Petiolis *in* *finibus* *aphyllis*, *superioribus* *foliolatis*, *inferioribus* *filiformi*.

7. *Lathyrus* *sahary* *L. gen. phed. p. 1030. W. ph. Del. p. 39. Bot. fl. austr. p. 324.*

14 *Lathyrus sphaerius* Herz obs. pl. III p. 39. Hort fl. austr. II p. 326. Abb. fl. ex.
p. 534.

[illegible]

Syr. L. angustata Roth. et Sm. fl. gr. prodr. II p. 66. et fl. græc. VII p. 84.
L. truncatipes Desf. fl. atl. VII p. 453. — *fl.* —
Marit. fl. cap. d. i. p. 591. — *Desf. fl. atl. VII p. 453.* — *fl.* —
Denf. fl. atl. VII p. 453. — *fl.* —
fl. atl. VII p. 453. — *fl.* —

Ab. in herb. et parum ad Brasilię ppe. Schum. et sine Popul.
H. conici. H. Maj. Linn. ③

15 *Lathyrus incognitus* L. p. fl. ed II p. 1030. Hort fl austr II p. 325. fl. h. f. m.

[illegible]

~~Leon. Jacq. h. v. m. n. tab. 86~~

Var. β . stars: entirely ovate, anisoprenich, lanceolate, linear-obov. - syn.
L. star bif. in exsert. zur bot. zeit. 1829 I.B. p. 19. Alab. fl. juss.
p. 172. Pet. bot. wey. n. 536. Koch syn. fl. germ. 2. II. p. 221.
var. γ . L. star bif. in exsert. zur bot. zeit. 1829 I.B. p. 19. Alab. fl. juss.
p. 172. Pet. bot. wey. n. 536. Koch syn. fl. germ. 2. II. p. 221.

Obs. Species arborescens in plures magnitudines plethorice elongata apice conchiformi
aut adpresso, quod d. statim strium per plures annos cultu constantem
deff. in hoc circuli omnes ~~complectitur~~ ^{complectitur} sunt omnes & plethorice ^{angustior} lance-
tem arborumque referunt: in alijs nullum speciem. Hanc
variant corolla longius vel vix exserta; hyemina pubescentia
vel glabra.

[illegible]

42
22. Lobbying monitoring Beruk in third row. bot. ed. II. p. 233.

L. nuda tuberosa, caulibus ~~anguste~~ ^{obtusiusculis} triangularibus, plus bi trijugis, pilosis
~~ellipticis - sessilibus lanceolatisve sulcatis laevibus, pilosis alatis, stylo linearibus.~~
*Syn. Bot. technomy L. f. pl. 2 p. 1028. Hort fl. austr. II. p. 323. Webb. Fl. ex.
 t. 537. O. prostratus Hoff l. c. p. 322. — Bon. Sav. engl. bot. t. 1153. ed.
 II. p. 537. Thall. herb. sup. ic. hb. I. sp. in lat. in 1-57.
 Pl. den. 1781. Thal. herb. sup. ic. hb. I. sp. in lat. in 1-57.
 Hab. in glaciis montium Arctice, et ad Gorgone in Thracia. Pl.
 purpureo-virens. Tb. van Z.*

Lathyrus variegatus Vif
L. indic tuberosa crasse fibrosa ^{mucronata}, caulis angulatis, folijs bi-trinervijs,
pilis late ovatis acuminatis, ~~basis oblique sub-rectis~~, petalis angustis
~~oblongis~~, ^{membris dimidiatis}, ~~stylis linearibus~~, ^{strobili calycum imbricatis} tubum aperturam
stylis linearibus.
Syn. *Brosia variegata* Desv. fl. neap. prod. suppl. t. p. 52 et fl. nap. II.
p. 146. Burdet. fr. ital. VII p. 924. O venetius Mill dict. n. s. pl. p.
exc. p. 536. O multiflorus Sieb. in bot Zeit. 1821 p. 97. O. vernus
Aleg. p. 104. p. 141. Pet. h. v. n. 573 gen. L.
Sibth. et Sm. fl. grec. c. VII p. 81. — Leon. fl. nap. t. 68. Sibth.
et Sm. l. c. t. 691. Column. ~~fl.~~ phytob. p. 60. Clus. hist. pl.
VI p. 232.
Juss. Vrii. Koisko, Castelnuovo, Japyx.

VI f. 232.
Hb in glaucis umbrosis, circa Zann, Irni, Kirovsko, Castellana, Joziz,
in montibus Olleth, Suleja, et in iugis arb. fl. rosei, viridis
perpetuo striata.
In a. 1. v. v. (18 v. v. L.) proceriora v. v.

869. *Ab. hoc spec. differt L. verum (p. 6 verum L.) praesentia minus*
drupa multiflora, semibus calycinis omnibus tubo brevioribus, et legumi-
nibus etiam junioribus luteis, quae in nostris siliis glandulosis - scabris.

24 *Lathyrus niger*. ~~17~~
L. radia fruticiformi, caulibus angulatis ramosis, foliis subsessilibus, pinnatis
 orbiculis oblongis mucronatis, pinnulis angustis alatis, mucronis densifloris, densi-
 calycis omnibus tubo brevioribus, ^{style} style linearis -
 Syn. *Ostrya niger* L. p. p. l. ed. II. p. 1028. *Lathyrus radia* Walp.
 588. Alab. fl. p. 122. Pitt bot. univ. v. 871. - ~~W.~~ Non Low. cap.
 Bot. t. 2788. Bed. fl. den. t. 1170. Pin. t. 1. v. t. 58
 Hab. in umbrosis ad Bellefonte prope Zane et in monte Odissonis prope Sign.
 H. purpureo-virens - Plante vivand nigrescit. Fl. immo. 4.

Tab. IV. Caesalpiniae N. Br. gen. rem. p. 19.
Stamina omnia libera. Legumen coriaceum. Embryo rectus.

Ceratonia L.

Gen. pl. ed. II. p. 368.

Char. Flors polygami aut divisi apetalae. Perianthium calycinum quatuor-
partitum. Stamina 5 sub styli hypogyni marginem inferiorem
stigma subrotundatum sessile. Legumen coriaceum compressum
saturius compressius induratum rectis spinis pilosis in lobis
pluribus inaequalibus divisum. — Cl. Polygamia Ord. Polygalia,
vel Diocia Ord. Pentandria Syst. sex. —

Ceratonia siliqua L. sp. pl. ed. II. p. 1518. Vrf. schiz. Deln. p. 30. Tab. fl.
aust. II. p. 662. Phil. fl. ex. pp. 544. Alch. fl. jud. p. 57. Ped. bot. univ. n.
237.

C. inermis, foliis ovatis oblongis plenis. —

Icon. Lam. enc. bot. pl. 859. Carr. ic. t. 113. Blackw. herb. t. 209. Camer. agit.
pl. p. 139. — Myr. Rozas.

Hab. in collibus, locisque agris saxosis insularum, Sicula, Lesina, Lipari. Fl. rubra. Fr.
austriaca. L.

Ufus. Fructus dulcis edulis, ex aqua decoctus in morbis praecum et postea
que emolliens natione adhibetur.

Pinj

Munda a Ligno d. 10 Dec. 1850.

Chara hispida L. sp. pl. ed. II. p. 1624. Nchb fl. ex. p. 149. Gaunter.

Dim. p. 17.

Ch. caule cernuente, ramisque densis reviridis setis-hispidis, bracteis quaternis longioribus linearibus mucosae obovatas superimixtis.

Iron. Pluken. ^{phys.} ~~an.~~ t. ~~143.~~ f. 6. bed fl. dan. t. 154. Low engl. bot. t.

463. Gaucher. l. c. l. II. f. XIV. Coll. et Germ. fl. parif. atl. pl. xxxviii. B.

Hab. in aquis stagnantibus circa Naranta. Fruct. aestate. C

Obs. Ab hac Drullen vixit Ch. hispidum β ? delmaticum
fl. delm. I. ² ~~max~~, cum in ~~specie~~ ^{planta} adae polygonos intermedium vixit
longior brevior et off. nuncius pediculus hanc sufficienter
distans ut hinc ^{distans} ~~specie~~ ^{specie} ~~specie~~ ^{specie} possit -

Flora Delmatica
addenda et emendanda.

Vol. I.

p. 32. Characeae sic dipneumae.

Chara Vaill.

Mem. de l'Acad. des sc. 1819.

Char. Sem. ac generif.

Subgen. I. Eithara.

Caules tubuli cernui spiculati, laxis, extrinsecus test. Nuclei spirales
subrotundi, apiculati, laxis, extrinsecus test. Nuclei spirales
subrotundi, apiculati, laxis, extrinsecus test. Nuclei spirales

Chara fedia A. Braun ^{not. I. 1834} ^{anal. d. sc. p. 334} Kütz. phyc. germ. p. 258. Gant.

ver. dist. pertract. Char. Austr. Vind. 1847. p. 18.

Ch. caule cernuente, diffo gracili, ramis subrotundis, bracteis linearibus
nucleis ovatis, apiculatis, laxis, extrinsecus test. Nuclei spirales

Syn. Ch. vulgaris L. p. pl. 1624. ^{not. I. 1834} ^{anal. d. sc. p. 334} Kütz. phyc. germ. p. 149. — Icon. low. angl. bot.
t. 336. Gant. l. c. t. II. p. 12-13. Coll. et Germ. atl. phyc. pl. xxxviii.

Hab. in aquis insularibus ^{Vigla ovata et laxis} ^{not. I. 1834} ^{anal. d. sc. p. 334} Kütz. phyc. germ. p. 149. Gant. l. c. p. 20.

Chara fragilis Desv. in Lorr. not. p. 126. Kütz. phyc. germ. p. 149. Gant. l. c. p. 20.
Ch. caule cernuente, diffo gracili, ramis subrotundis, bracteis linearibus, nucleis ovatis, apiculatis, laxis, extrinsecus test. Nuclei spirales

Syn. Char. vulgaris L. p. pl. 1624. ^{not. I. 1834} ^{anal. d. sc. p. 334} Kütz. phyc. germ. p. 149. Gant. l. c. p. 20.

Ch. Henrici Bruch. in Flor. 1826 II. p. 489. Ag. syst. alg. p. 129. Ch. longibracteata
Palenb. Krypt. fl. p. 200. Kütz. phyc. germ. p. 843. — Icon. Henrici. l. c. t. xxxiv-xxxv.

Gant. l. c. t. II. p. xv. a. b. Vaill. mem. de l'Acad. des sc. 1819. tab. 3. f. 1. Cop. et Germ. atl. phyc. pl. xxxviii.

Chara repens Bauer of Moench. A. Braun ^{not. I. 1834} ^{anal. d. sc. p. 334} Kütz. phyc. germ. p. 149. Gant. l. c. p. 20.

Ch. caule cernuente rigida subrotunda, ramis cernuente, bracteis linearibus, nucleis ovatis, apiculatis, laxis, extrinsecus test. Nuclei spirales

Syn. Char. repens L. p. pl. 1624. ^{not. I. 1834} ^{anal. d. sc. p. 334} Kütz. phyc. germ. p. 149. Gant. l. c. p. 20.

Hab. in aquis ad ripas fl. Henrici p. 149. Gant. l. c. p. 20.

Obs. Offic. Ch. ~~Boeckii~~ ^{May.} Gmel. sed qua jungetur cum Ch. ~~boeckii~~ a illo ~~phlo.~~
in fl. ex p. 149, sed differt caule tubuloso, pinnatis, coriatis, nec uiridis, et ~~gemmae~~
univisa obita. In nostra nuculae ~~exsertis~~ ^{exsertis} pedunculatae et vixtae, nec ellipticae
et hinc quod plerumque in Ch. ~~boeckii~~ ^{neganda}

Subgen II. Nitella Ag. 1841. p. 22111
Caulis exting. ^{medi.} Nuculae ~~exsertatae~~ ^{exsertatae} apice ~~cornu~~ ^{cornu} decidua ~~nuda~~
Cham flexilis L. p. 1624. fl. ex p. 148.

Ch. caule gracili ~~rumpo~~ ^{rumpo} pellucid. ~~rumpo~~ ^{rumpo} ~~unif~~ ^{unif} ~~apice~~ ^{apice} ~~bi-tripunctat.~~ ^{bi-tripunctat.} ~~nuculis~~ ^{nuculis} ~~sublongis~~ ^{sublongis}
diff. ~~antheris~~ ^{antheris} ~~solitariis~~ ^{solitariis}.

Syn. Nitella flexilis Ag. l. c. p. 124. - Non. ~~Quadr.~~ ^{Quadr.} ~~ic. bot. VIII f. 1071.~~ ^{ic. bot. VIII f. 1071.} ~~San. angl. bot.~~ ^{San. angl. bot.}
~~Hab. in vasis luteo flumensibus prope~~ ^{Hab. in vasis luteo flumensibus prope} ~~deris p. Funt. Jul. Sept.~~ ^{deris p. Funt. Jul. Sept.} ~~excl. syn. p. 1070.~~ ^{excl. syn. p. 1070.}

~~Cham nitida~~ ^{Cham nitida} ~~in in low eogl~~ ^{in in low eogl} ~~62. p. 148.~~ ^{62. p. 148.} ~~phlo. ex p. 148.~~ ^{phlo. ex p. 148.}
~~Ch. caule gracili rumpo pellucid. rumpo elongat. rumpo~~ ^{Ch. caule gracili rumpo pellucid. rumpo elongat. rumpo} ~~fructif. fructif. fructif.~~ ^{fructif. fructif. fructif.} ~~excl. syn. p. 1070.~~ ^{excl. syn. p. 1070.}

Syn. Conferva nitida Müll. in ad. fl. den. bot. v. 6. Cham flexilis var. nitida
Ch. caule gracili ~~rumpo~~ ^{rumpo} pellucid. ~~rumpo~~ ^{rumpo} ~~fructif.~~ ^{fructif.} ~~fructif.~~ ^{fructif.} ~~excl. syn. p. 1070.~~ ^{excl. syn. p. 1070.}

var. p. nitida: rumpo fructif. ~~excl. syn. p. 1070.~~ ^{excl. syn. p. 1070.} Syn. Ch. flexilis var. nitida phlo.
non bot. VIII p. 37. Ch. nitida ~~in in low eogl~~ ^{in in low eogl} ~~bot. p. 1703.~~ ^{bot. p. 1703.} ~~phlo. ex p.~~ ^{phlo. ex p.}
148. - Non ~~low eogl~~ ^{low eogl} ~~bot. p. 1703.~~ ^{bot. p. 1703.} ~~phlo. ic. bot. VIII t. 1072.~~ ^{phlo. ic. bot. VIII t. 1072.}

Hab. in vasis luteo flumensibus prope ~~deris~~ ^{deris}: var. ad saxa litorum maris circa
Lithuania ~~in vasis luteo flumensibus prope~~ ^{in vasis luteo flumensibus prope} ~~deris~~ ^{deris} ~~bot. p. 1703.~~ ^{bot. p. 1703.} ~~phlo. ic. bot. VIII t. 1072.~~ ^{phlo. ic. bot. VIII t. 1072.}

Obs. Conferva nitida Müll. in ad. fl. den. v. p. 6. t. 261. rumpo fructif.
excl. syn. p. 1070. ~~phlo. ic. bot. VIII t. 1072.~~ ^{phlo. ic. bot. VIII t. 1072.} ~~phlo. ex p.~~ ^{phlo. ex p.}

Lychnothamnium Reprecht
Subgen III. ~~Cham nitida~~ ^{Cham nitida} ~~in in low eogl~~ ^{in in low eogl} ~~62. p. 148.~~ ^{62. p. 148.} ~~phlo. ex p. 148.~~ ^{phlo. ex p. 148.}

Caulis exting. ^{medi.} Nuculae ~~exsertatae~~ ^{exsertatae} apice ~~cornu~~ ^{cornu} decidua ~~nuda~~
Cham nitida L. p. 1624. fl. ex p. 148.

Cham Nitella L.
C. caule ~~rumpo~~ ^{rumpo} ~~exsertata~~ ^{exsertata} ~~apice~~ ^{apice} ~~cornu~~ ^{cornu} decidua ~~nuda~~
caerulea ~~exsertata~~ ^{exsertata} ~~apice~~ ^{apice} ~~cornu~~ ^{cornu} decidua ~~nuda~~

Syn. Cham Nitella L. ~~exsertata~~ ^{exsertata} ~~apice~~ ^{apice} ~~cornu~~ ^{cornu} decidua ~~nuda~~
Hab. in foris insulae Lema, ubi litorum a ~~Matth. Boten~~ ^{Matth. Boten} ~~exsertata~~ ^{exsertata} ~~apice~~ ^{apice} ~~cornu~~ ^{cornu} decidua ~~nuda~~

Al. Nitella. ~~exsertata~~ ^{exsertata} ~~apice~~ ^{apice} ~~cornu~~ ^{cornu} decidua ~~nuda~~
Deris. ~~exsertata~~ ^{exsertata} ~~apice~~ ^{apice} ~~cornu~~ ^{cornu} decidua ~~nuda~~
a. Nitella. ~~exsertata~~ ^{exsertata} ~~apice~~ ^{apice} ~~cornu~~ ^{cornu} decidua ~~nuda~~

Oed. fl. Dan. tab. 564. Rehb. ic. ^{bot. XI} ~~fl. græc.~~ f. 1436

Hab. in arenosis maritimis insularum Ossero et Cherso. Fl. a. Junio ad Sept. 24.

P. 57. = post Agrostidem verticillatam Vill. addatur

Agrostis pungens Schreb. gramm. II p. 46. Rehb. fl. exc. p. 140.²

A. paniculae thyrsioidae compactae acutae folio supremo vaginatae ramis sparsis, apice densifloris, flosculis pedicellatis glabris, glumellis muticis, foliis distichis convolutis rigidis, ligula barbata.

Icon. Cavan. ic. et descr. II t. III. Rehb. ic. ^{bot. XI} ~~fl. græc.~~ f. 1437.

Hab. ad litus scopulorum circa insulam Lesina, unde misit Al. Stalis. Fl. aestate. 24.

54 = Ante Digitalium Sep. addendus

Chaeturus Link.

in Schrad. Journ. 1799. 4.

Char. Flores laev. paniculato-spicati pedicellati. Stamina unguis bivalvis compressa
valvis acuminatis, altera longe aristata. Stamella bivalvis mutica, valvas in-
feriore trifida, superiore bifida. Paleae sub-truncatae. Rigmata plumosa.
Class Triandria, Ord. Digynia Syst. nat.

Chaeturus fasciculatus LK. loc. c. II p. 313.

Ch. Culm. erecto, foliorum ligula elongata, paniculae ramulis erectis subspicatis.

Syn. Polypogon subspicatus W. enc. bot. hex. 1 p. 88. P. fasciculatum Prof. syn.
pl. 1. p. 80.

Hab. circa Lara unde misit ex. Doct. Dom. Papafava. Panicula fusco-virens. Fl.
verc. O.

56 = Post Calamagrostidem Pygmaea inseratur

Calamagrostis montana Hoff. gram. austr. IV. p. 27. et fl. austr. I. p. 104.

Rehb. fl. exc. p. 26.

C. panicula ^{stricta} ~~stricta~~, sub anthesi patula, glumis lanceolatis acuminatis, glumellae pilos superantibus, arista ex imo dorso, geminata, valvae glumellae exterioris ~~fl.~~ pilos superante.

Syn. = Arundo montana Pand. fl. helv. 1 p. 200. Ar. varia Schrad. fl. germ. 1. p. 216

A. acutiflora Schrad. l. c. p. 217. Calamagrostis acutiflora DC. fl. fr. V. p. 253

Rehb. in botan. p. 44, et fl. exc. p. 26. C. sylvatica Hoff. gram. austr. IV. p. 28, et
fl. austr. I. p. 105. — Flora Rehb. in bot. XI f. 1442. 1443. Hoff. gram. austr. IV.

t. 46, et 48.

Hab. in montosis circa Tagus, ubi legit Fr. Nummayer. Panicula purpurascens,
vel straminea. Fl. aestate. 2.

58. Agrostide stolonifera addatur

A. stolonifera sp. maritima; paniculae spiciformi compacta, foliis rigidos glaucis — Syn. A. maritima Lam. enc. bot. I. p. 60. Rehb. fl. exc. p. 140.² — Flora.

A. maritima Lam. enc. bot. I. p. — Rehb. fl. exc.

P. 58. Post *Melinum multiflorum* Cav. add.

Melinum paradoxum L. sp. pl. ed II. p. 90. Hort. fl. austr. I p. 95. Rel.
fl. ex. p. 28. —

M. panicula elongata nectante, ramis gracilibus flexuosis, glumis ovatis
longis acutis, glumellae pilosae valvae exteriores aristatae.

Syn. *Agrostis paradoxal* L. sp. pl. ed I p. 62. — Nov. Hort. ii. gramm. austr. III

Bochb. ic. bot. XI. f. 1460. Pluker. phyt. t. XXXII. f. 2.

Hab. in sylvaticis umbratis agri Bagusini, ubi legit Neumayer *Panicula*
viridens. Fl. ver. 4.

P. 58. Post *Stipa pennatam* L. addenda

Stipa tortilis Desf.

Stipa tortilis Desf. atl. I p. 99. Malg. ev. pl. austr. p. 21. Gussone Syn. fl. sic. I p. 12

Boetol. fl. ital. I p. 688.

St. aristis longis basi contortis glumellaeque exteriores villosae, supremae glabris
foliis canaliculatis, supra pubescentibus

Syn. *Stipa palustris* Boetol. et Sm. fl. gr. prodr. I. p. 65. — Nov. Boetol. et Sm. fl. gr. I

I. t. 86. Desf. fl. atl. I t. 31. f. 1. Bouche mus. tab. 99. —

Hab. in apricis sacosis aridis insulae Lesina, ubi invenit Mattia Boetol. fl. ver.

P. 59. *Leucostachys Stipae Aristellae* L. adduntur insulae Lesina et Syracusa
ille Panicis conciformis add. mus. Lesina ad vias agrestes

P. 62. *Phalaris canariensis* L. oritur quaque in insula Lesina.

P. 62. *Phalaris argutia* L. oritur hactenusmode circa Syracusa et Tragusa. add. ione
Cavan. ic. et Dur. pl. I. t. 64. Barrel. ic. pl. 700. f. 1. post nov. sequentes

Phalaris nodosa L. syst. veg. ed XIII p. 88. Boetol. et Sm. fl. gr. prodr. I. p. 37

Boetol. fl. ital. I p. 389.

P. radice perenni, culmi nodis inferioribus globosis incrassatis, tercia cylindrace
attenuata, spiculis obovatis, glumae valvae navicularibus integris crosse
glumellis omnibus villosis, accessoria unica lineari-subulata immixta.

Syn. *Ph. tuberosa* L. mont. pl. atl. p. 557. — Nov. Boetol. et Sm. fl. gr. I tab. 5

et M. quid. fructu coll. Nov. de Vinea

p. 34. — *Equisetum elongatum* W. ist neuerer Name, jedoch ist es gebräuchlich.

E. ramosa Schreb. cat. pl. horti. 1807, quod igitur proferendum.

p. 36 - post genus etc

p. 37. pinnatifid oblongif corr. pinnatifid oblongif

p. 38. ante Aspidium *Centium insect*

Aspidium aculeatum W. & A. v. p. 258

[illegible]

Syn. *Rhynchospora aculeatum* L. in vol. 2. p. 1552 ex part. Hist. pl. austr. II. p. 675.

~~Gen. ^{Jan} Hoof left of. vol. XIV 1-3 f. 15. Rother. also. 1-1832~~

Aspidreum haplolum Den. mem. in Dr. eua unov. Jalc. Nagl. 1832. p. 25.

~~— Non. Den. l. c. t. IV. f. 7. Moif. Sept. 18. l. c. t. IV. f. 13. Platten~~

~~alm. 180 f.~~

also 1800 f. 1.
Hab. in umbrosis, glenibus montium Vellebitz et Bickovo. Prut. apud 7.
H. 2. 1/2. * l. b. attenuabili et im

Obs. Differt a sequente ^{planifolia} ~~planifolia~~ ^{reticulat} ~~reticulat~~ (nec tamen bifi attenuat) et in
 illis) ^{reticulat} ~~reticulat~~ ^{et bifi} ~~et bifi~~ ^{superum} ~~superum~~ ^{acumulat} ~~acumulat~~, nem in A. lat.
 bifi et hinc ^{est} ~~est~~ ^{caeris} ~~caeris ^{sed} ~~sed ^{acumela} ~~acumela~~ ^{caerit} ~~caerit~~.~~~~

Pinnularia recta var. *recta* + *capitata*, *repens* Moench.
 p. 42. *Chilanthum* ~~*pinnatifidum*~~ *oson* L. et Ch. *pinnatifidum* V.
~~*Chilanthum* *pinnatifidum* V.~~ *orientalis* *gigas*
 in saxibus et nemoribus agris ibidem Lecina.

un sautoir et un anneau de fer. Les uns

f. 43. *Opilobolus rugosus* L. var. *hab. et in monte Orizaba*
Opilobolus parviflorus L. var. *hab. et in monte Orizaba*
Opilobolus parviflorus L. var. *hab. et in monte Orizaba*

Don. Basal. icon. v. 252 p. 2. and visit Al. Salis; S. Clemente page number Leina varum H. ag. 4.

hab. in saxif. scopuli S. Clemente pag. novam Lehm.
Bot. Tern. Lam. in bot. pl. 66a f. 3. differt a nostra fere stylis brevioribus
lobis, elliptico-lanceolatis, et spicae apice filiformi longiore, fere in usum
coarctata.

p. 36. post genus *Cetrach* W. in *am*
Comitatus *San*

Gravimetry Sur.

Syn. flie. p. 3.

Char. Son. indurata det. Kuhn. *Sonchus oleraceus* - *Char. Cryptogamia*. Bot. Kuhn. 1848. p. 426. *Sonchus oleraceus* L.

Erasmia leptophylla Sw. l. c. p. 23 et 218. Berl. ~~amer.~~ ital. p. 426. ~~Sw.~~ 1919 p. 111.

II. p. 254

II. p. 284. *rubra*
G. pumila *hyemalis* *glaberrima* *granulata* *cuneiformis* - *obovata* *flabellata* *lobata*, -
lanceolata *lanceolata* *L. p. p. II. p. 1553* - *Per. Magnol. bot. mon. p. 5.*

Syn. Polypodium leptophyllum L. f. pl. D. II. p. 1553 - Icon. bot.

Barcl. ic. pl. 421.

Parcel: 12. pt. 431.
Hb. ad mens. et mens. apri. Jozsefi. uel. Capt. J. Neemayer. Fruct. & vire in aestate. (4)

p. 50. Pollinia diffractis. sp. Habitat etiam in demetis collium circa
Jagusa

p. 52. Heteropogon glaber Pers. habitat etiam in rupibus circa ins.
Lerina.

ibid. Saccharum striatum sp. habitat circa Jagusam ad Ombla (nec
Ombla) et ad Grenchetta. Post Saccharum add.

Imperata Cyrill.

Pl. var. vna veg. II. p. 26.

Char. flor. hermaphroditi numeroso-jicati, spiculis geminatis neutris, glan-
bralis unispermis ^{ext. tri} sericeo-lanata glomerata bralis membranaceis. Paleae ob-
longae. Stigmata plerumque ~~plena~~ ^{ext. tri} exserta. Clap. Brachii ord.
Dyggvi. 194. var. —

Imperata cylindrica Beauv. agrost. p. 7. Rich. syn. fl. grm. II. p. 395
Syn. Lagurus cylindricus L. p. 96. et II. p. 120. Saccharum cylindricum Lam.
enc. bot. I. p. 588. Pers. fl. ital. I. p. 232.

Sp. L. ital. Hort. fl. austr. I. p. 66. — Imperata arundinacea
in grm. IV t. 40. Lam. enc. bot. fl. 40 f. 2. Dant. icon. II. — Mly.
Zabod

Hab. in graminibus insulae Sansego, in agris circa Jella in insula Lerina
et in insula San Pietro prope Jagusa. Flor. aestiva. 7.

Hb. sub litt. a. iconis Lamarckii flor. exhibitus Brachii quod salum.

p. 53. lin. 33. lacus Moringe corr. lacus Moringe.

p. 54. antea diffinimus sup. adarsidus (Chaetanthus)
p. 56. post Calamagrostidem Epigejss inuen

Calamagrostis montana Hort. grm. austr. IV. p. 27. et fl. austr. I.
p. 104. Phil. fl. ex. p. 26.

C. panicula stricta, sub antheri palea, glan-
glanellae pilos. superstitibus, anthera ex imo dorso, geminata, vltima glanellae
exterioris pilos. longior

Syn. Arundo montana Gaud. fl. helv. I. p. 200. A. vana Schrad. fl. grm. I. p. 216.
A. acutiflora Schrad. l. c. p. 217. Calamagrostis acutiflora L. fl. fr. V. p. 255.
p. 14. et fl. ex. p. 26. — Non. Phil. 12. bot. III. 1442. 1443.
Hort. grm. austr. IV. t. 46 et 48.

Hab. in montibus circa Jagusa sub lignis fr. Neumayer. Panicula purpurea,
et straminea. fl. aestiva 7.

p. 56. Agrostis, Monifera add.

A. Monifera p. maximum: panicula spiculis compacta, filis nigris glaucis — Syn.
A. straminea fl. L. fl. fr. II. p. 92. A. maximum Lam. enc. bot. I. p. 104. Phil. fl. ex.

And in giving receipt early enough. Letter, not insert gr. Math. Boston
H. W. C.

p. 59. Leaf natively *Nipae Anisotae* L. acuminata insularis Lessing ex

Giugiana: illo Panicis crueiformis adha in- L. lina ad vias agrorum

p. 62 *Phalaris* ^{canariensis} L. venter grisea in noua Lesina.

p-62 *Phalaris apiculata* L. var. *candicans*, var. *Zera* (L.) Trin. & Hook.
Japan Cava. ~~et~~ *et* *var.* *t. t. d. d.* David. *et* *var.* *t. t. d. d.* *et* *var.* *t. t. d. d.*
Phalaris nodosa L. var. *reg.* c. XIII. p. 89. *et* *var.* *t. t. d. d.* fl. gr. *et* *var.* *t. t. d. d.*

Berol. fl. ital. I. p. 339.

P. nuda peruviana
~~P. nuda~~ ludaei nudi imprimis glabro-mucronati, spica cylindrica - attenuata

speculis oblongis, glaucis, coloris nigris, calicibus integris, corollae, glaucis, speculis
oblongis, albis, unguibus ^{linearibus} ~~oblongis~~, lobis nigris

Syn. Ph. tuberosa L. mant. pl. alt. p. 557. — Icon. 18th. et Am. fl. gr. 5. tab 56.

Burcl. ic. pl. 10. Mont. cat. frimp. agr. bonon. tab. icon. n.º 43.

Abd. in herb. incul. *Lexia* und nicht *prof. Phil.* H. vers. 4.

* *Phalaris minor* Detz. det. the J. S. Maly in pl. aufg. 26. Berl. H.
Ital. I. p. 342

ital. l. p. 342

P. variis annua, culmi vixi aequalibus, spica cylindrica, spiculis oblongis,
glumae vixi navicularibus, ovip, glumellis ovatis, villis, necnon
semita lineari-subulata minutis.

Amida Utsun - 1000 - 1000

Syn. Ph. aegyptia W. sp. pl. 1. p. 326. Hort fl. aegypt. 1. p. 86. Phob. fl. aegy. 32

- Non. Host ic. gram. austr. II, p. 39. Philb. ic. bot. ^{xif} 1493. Th. F. Nees jun. pl.

à l'herbier et ajoutée avec Schwinn, Salomon, Bruno pour l'ajout, et en 1891

offered to Leomin. H. m. : 2. inclusion caused by a permeable hydrogen

[illegible]

Spencer

Phalaris paradoxa L. ff. pl. 2. II. p. 1665. Hort. fl. angl. I. p. 87. Feb. 4.

exc. p. 32. Mely ca. pl. austr. p. 26.

P. nodosa annua, spica oblonga ^{abortiva} bry. attenuata, spiculis oblongo-^{abortivis} obtusis, lemnas

relat. nasicularibus ~~capitulis~~ ^{capitulis} ~~dentatis~~ ^{dentatis}, floralibus ~~interioribus~~ ^{interioribus} ~~capitulis~~ ^{capitulis}, ~~quibus~~

fully accepting brief ^{other} residing.

Icon. Reichb. ic. bot. XI p. 1491. Hott gram. austr. II t. 40. 16th. et sur. fl. gr.

f. 26. 58.

Ab. in salis insularum Once, Levine et dale Jozeg. fl. mag^a, Peruv.

Spei mit-Hammer: 0

Bl

Baldinger ~~Carl May at Lehr~~
G. M. S.

~~Flow water~~ n 99

~~Char. Flory paniculati, sub anthesi apert. Spiculae uniflorae.~~

Obs. Characterum generum Phalaridis sic emenda

Char. Hory composita nitens perist. Spiculae floratae unius hermaphro-
ditae fertile, uno altorum stentis specimenformi. Gleba hermaph-
rodica conspicua glebulae longior valis alata canaliculata. Paleolae oblongae
stigmati glebulae. Laryngis glebulae testis. -

Post Phalaridis ~~infera~~ adventu

Baldingeri Gaertn. May. et Schr.

Flor. der Wetter. n. 99.

Hory penicillati sub anthesi effusi. Spiculae floratae unius hermaphroditae
fertile, ~~duobus~~ ^{pluribus} sterilibus specimenformibus. Gleba hermaphrodica
conspicua longior valis alata canaliculata. Paleolae oblongae. Stigmati
glebulae. - Laryngis glebulae testis. - Clasp. Inordini Ord. Symplicis lyp. vix.

Baldingeri arundinacea Robt. fl. ex. p. 33. et ic. bot. XI p. 20

Syn. Phalaris arundinacea L. sp. pl. ed. II. p. 32. Hort fl. conf. II. p. 85. Alech. fl. p. 14.
p. 245. Arundo chlora W. sp. pl. I. p. 457. Calamagrostis chlora Roth. fl.
oxon. p. 37. Baldingeri chlora Gaertn. May. et Schr. fl. Wetter. n. 99. Syngraphis
arundinacea Trin. fund. agror. p. 125.

Don. Th. F. Nees gen. pl. pr. XI. n. 6. D. fl. du. t. 259. Low. engl. bot. t. 402
et 2160 f. 2. Hort ic. grm. conf. II. t. 33. Leef. herb. hb. 7. f. 3.

Hab. in pasc. montan. prope Zar, und nicht prope Alchinger. fl. May.
Tunis. Panicle alba, vel ex alba et purpurea var. 2.

Obs. Radix fibrosa. Culm. erecti, foliis arundinaceis lanceolatis - linearibus, glau-
cis alba obso. vel pinn. Panicle acuta aut et prope anthesin bifida. Flos
hermaphrodica, in floribus parvulis patula, et glebae valis conspic-
ua et canaliculata id est omnimodis distincta a Phalaride prope distincta.

p. 65. Post Phleum tenui Schrad. adsp.

Phleum arvense L. sp. pl. ed. II. p. 88. Hort fl. conf. I. p. 84. Phb. fl. ex.
p. 31.

Ph. spica cylindrica bsp. glauca attenuata, floribus oblongo-lanceolatis, glau-
cis acuminato-mucronatis nervis valis, radice annua

Syn. Phalaris arvensis W. sp. pl. p. 328. Phalaris arvensis Gaertn. et Sch. 1798. 107.
II p. 409. - Don. Phb. ic. bot. XI. f. 1482. D. Th. F. Nees gen. pl. pr. XIII n.
2 f. 17-21 Low. engl. bot. t. 322.

Hab. in arvis insulae Lissa, und nicht d. P. Witske. Spica straminea fl.

Obs. ^{supra} ~~infera~~ ^o agerum paleolarum oblongarum distincta -

Phleum nodosum All. in fl. ges. II. p. 233. Hort fl. austr. I. p. 83.
Ph. ~~*spica cylindrica*~~, *sterilis oblonga, pennis*

Ph. ~~Spica cyclotoma~~, florulis oblongis, glumis

Phleum aegyptium Will. *hisp.* 108 pl. de Saugl. II. p. 61. Hist fl. aegypt. 2 p. 84.

Pl. spica cylindrica bifi. glomeris truncatis, florum breviter ~~longis~~
 corollis-tubulosis, glomeris ^{uncatis} ~~coniformibus~~ truncatis apice inflatis corollis
 numerosis. herbis.

Syn. Phleum paniculatum Thed. fl. angl. p. 26. *Phalaris agnes* Retz obs. 64.

IV. p. 14. *Chitachloa ager* Beauv. agron. p. 37. — Ten. Reichb. ic. bot.

xl. f. 1487. Th. 2. Neq. ju. fl. 13. n. 3 f. 22. Say. ic. fl. var. I. t. 14.

Hort ic-gram-aufw-II, b. 37

1101. 15
Hab. in ~~homo~~^{alluv.} ~~gemma~~ arena gelata insulae Lerina, und nupsz jng Itali. Spec
unvoll. *PC* *incaute acutata*. ⑤

p. 63 Post Pleura Becktonie DC. apud ut i. 68 regimen prole a Pl.
utq. non differt, hoc postremo iten Pl. prole variet maxim
habund, ad.

Alchemilla agraria L. *prostrata* 788. Herb. fl. caesp. *typ.* 81. *pub.* fl. ex.

Phl. spica *sp.* trif. *truncata*, fl. *oblongo-lanceatis* *undique* *canis* *ulaz*,
crispi, *glauca*, *acuminato-angustis* *oblongo* *truncatis*, *angustis* *glauca*, *ulaz* *junioribus* *celis*,
 17 *strictis*, *ulaz* *affluent*, *indis* *phlo* *parum*

Non. Dec. ii. bot. xi f. 1485. Red. fl. du. t. 263. Host ic. grm. austr. III. t. 10.

Schweinb. p. 12 t. 3.

Hb. in ramifying clusters by sympyces, ^{with} ~~some~~ few lvs. generally white leg'd & narrow
 yew. H. acut. 4 ^{inches}

668. ab. loc. d. part. ^{young} flowers ^{infundibular} tubularly, ^{reflex} reflex, ^{anthers} anthers

f. 66 ob. 66. *Ida hii celibis, ingenua antiquae nobis, antist. potibz.*
Grassia abiecerunt et C. auleat minter, pueri in religiois circa Hagis.
66. puer natum Aime apudiam corp lacum Boccagnis.

p. 69. Locum natalem Armi agrosae vixi lacum Dougnoffo.
 p. 70. Horum Armae athermutha Best praeclatiss, vixi A. hifp
 10th capit. bth. III p. 19 a p. 19 illa vixi A. hifp
 ab A. p. 19 glauca lacum schief angustiss, p. 19.

p. 68 Post Aream mediam Jou.

Aira caespitosa L. sp. pl. ed. II. p. 96. Host fl. austr. I. p. 114. Richb. fl. exc. p. 50.

A. panicula laxa verrucata, flumis apice crispis emarginatisve floralis ^{subaequantibus} ~~basibus~~, glomerellae
basi pilosae valva exterior truncata crenulata dorsa basi ~~basi~~ aristata, interiori bifida,
culmo nudipubulo, foliis lineantibus ~~paniculis~~ ^{paniculis} superis herbaceis.

Syn. Deschampsia caespitosa Beauv. - ess. D'une nouv. Agr. p. 12. — Icon. Richb. ic. bot. 21. ^{pl. 2} ~~pl. 1~~
Sow. engl. bot. t. 1453. Host ic. gram. austr. II. tab. 42. (caesp. fl. herb. n. tab. IV. p. 8.

Hab. in pratis circa Ivrea, ubi legit A. Candrich. Spiculae stramineae vel e viridi
et purpureo variae. Fl. aequalis. 4

Obs. Specimina nostra e loco majori formam contractam exhibent, et varietati montanae ^{ad. f. h.}
icon 1683 exhibita, vel ^{rectius} ~~pl. 1~~ var. **B.** a d. Minio descriptae in Spec. gram. tom. III. ^{potius}
quam vno speciei typo respondent.

p. 82. Port Poae Engrortum

Poa pilosa L. p. fl. ed. II. p. 100. Hort fl. austr. I. p. 150.

P. panicula aequali, ramis inferioribus tenuissimis ^{intra basin} ~~sub~~ pilis in tracha
patulis; spiculis linearibus 5-12-floris, florculis adnatis obsoletis univariis,
basi natis, culmis decumbentibus, spinis on lobatis, radice annua.

Syn. Engrortis pilosa Beauv. ess. d'une nouv. Agrost. p. 21. Richb. flex. p. 48. Poa verticillata
Cav. ic. et descr. pl. I. p. 63. Engrortis verticillata Beauv. l. c. Richb. flex.
p. 48 — Non Richb. ic. bot. XI f. Cav. l. c. t. 93. Hort ic. grm. austr.
II. t. 68.

Hab. in cultis et ad vias stationis Sagerinae ubi legit Neesmayr. Panicula purpurea.
H. aestate. ©

Ver. p. arundinacea : ^{major} *speculis laxifloris*, ^{ex vint et pinguis vici} ~~in vint~~ — *Syn. M. arundinacea* Schrad
 l. c. p. 336. *M. libanotis* Hook fl. austr. l. p. 118. *Pubb. fl. ex. p. 44. M. albissima*
 Link h. bot. duer. l. p. 196. — *Son. Pubb. ic. bot. xi p. 1607.*
Hab. in sylvestribus agris *Pagopuni* : var. *intra vici* ^{speculis in quibus alba volucres} *fl. septem 4.*
Bb. Nulla certe est varietas *differt a specie* ^{sed} ~~per~~ *ex loco umbros pinguis*
proceris fit, laxiflora et pallidiora —

p. 85. ad *Cynarum* *crinitum* loco involucri *glaucae* *mentis* loco involucri
crinitis *mentis*: post *Equisetum* addenda

Lamarckia Moench

Char. ~~flor~~ ^{meth. gl. p. 201.} ~~racemosa~~ ^{racemosa} ~~paniculata~~ ^{paniculata} ~~spiculae~~ ^{spiculae} ~~pubescentes~~ ^{pubescentes} ~~exsertae~~ ^{exsertae}
 strobili mixtae. hermaphroditi inferiori, et strobili sterili superioris antheris, cum, hinc inde
 glomeris mixtae. glomeri bivalves lanceolati - subulati. stamella intra
 inferiori apice antheris bisp. superiorum breviora sunt. paleae ovatae
 oblique recurvatae. ligam. cordatae glomeris. Caryops. glomeris
 laxe circumdatis. — Chap. Ind. ed. Regni 1794. ex —

Lamarckia aurea Macneil l.c. Th. F. Near gen. pl. proc. XIV p. n. 9.
Syn. *Cynurus aureus* L. J. fl. 2 II. p. 10. Sibth. et Sm. fl. gr. jordan. i. p. 89.
Chrysomys cynurinus Pers. syn. fl. l. p. 80. Host fl. austr. i. p. 20.
Astr. aureus Beauv. ex Desf. nov. agrost. p. 14. pub. fl. ex. p. 36.
— Don. Th. F. Near l.c. t. 9. f. ~~pub.~~ ic. bot. xi p. 1515. Sibth. et Sm.
fl. gr. i. t. 89. Host ic. gram. austr. III. t. 4. Bonann. graph. t. 6. p. 3.
Sedula Oken ubi novam plantam invenit Doct.

Hab. in open, stony insular grass near ...
 No. *Panicum latifolium* ^{fl. Moq.} ^①
 Obs. *Panicum latifolium* ^{sp. nov.} ^② ^③ ^④ ^⑤ ^⑥ ^⑦ ^⑧ ^⑨ ^⑩ ^⑪ ^⑫ ^⑬ ^⑭ ^⑮ ^⑯ ^⑰ ^⑱ ^⑲ ^⑳ ^㉑ ^㉒ ^㉓ ^㉔ ^㉕ ^㉖ ^㉗ ^㉘ ^㉙ ^㉚ ^㉛ ^㉜ ^㉝ ^㉞ ^㉟ ^㊱ ^㊲ ^㊳ ^㊴ ^㊵ ^㊶ ^㊷ ^㊸ ^㊹ ^㊺ ^㊻ ^㊼ ^㊽ ^㊾ ^㊿ ¹ ² ³ ⁴ ⁵ ⁶ ⁷ ⁸ ⁹ ¹⁰ ¹¹ ¹² ¹³ ¹⁴ ¹⁵ ¹⁶ ¹⁷ ¹⁸ ¹⁹ ²⁰ ²¹ ²² ²³ ²⁴ ²⁵ ²⁶ ²⁷ ²⁸ ²⁹ ³⁰ ³¹ ³² ³³ ³⁴ ³⁵ ³⁶ ³⁷ ³⁸ ³⁹ ⁴⁰ ⁴¹ ⁴² ⁴³ ⁴⁴ ⁴⁵ ⁴⁶ ⁴⁷ ⁴⁸ ⁴⁹ ⁵⁰ ⁵¹ ⁵² ⁵³ ⁵⁴ ⁵⁵ ⁵⁶ ⁵⁷ ⁵⁸ ⁵⁹ ⁶⁰ ⁶¹ ⁶² ⁶³ ⁶⁴ ⁶⁵ ⁶⁶ ⁶⁷ ⁶⁸ ⁶⁹ ⁷⁰ ⁷¹ ⁷² ⁷³ ⁷⁴ ⁷⁵ ⁷⁶ ⁷⁷ ⁷⁸ ⁷⁹ ⁸⁰ ⁸¹ ⁸² ⁸³ ⁸⁴ ⁸⁵ ⁸⁶ ⁸⁷ ⁸⁸ ⁸⁹ ⁹⁰ ⁹¹ ⁹² ⁹³ ⁹⁴ ⁹⁵ ⁹⁶ ⁹⁷ ⁹⁸ ⁹⁹ ¹⁰⁰ ¹⁰¹ ¹⁰² ¹⁰³ ¹⁰⁴ ¹⁰⁵ ¹⁰⁶ ¹⁰⁷ ¹⁰⁸ ¹⁰⁹ ¹¹⁰ ¹¹¹ ¹¹² ¹¹³ ¹¹⁴ ¹¹⁵ ¹¹⁶ ¹¹⁷ ¹¹⁸ ¹¹⁹ ¹²⁰ ¹²¹ ¹²² ¹²³ ¹²⁴ ¹²⁵ ¹²⁶ ¹²⁷ ¹²⁸ ¹²⁹ ¹³⁰ ¹³¹ ¹³² ¹³³ ¹³⁴ ¹³⁵ ¹³⁶ ¹³⁷ ¹³⁸ ¹³⁹ ¹⁴⁰ ¹⁴¹ ¹⁴² ¹⁴³ ¹⁴⁴ ¹⁴⁵ ¹⁴⁶ ¹⁴⁷ ¹⁴⁸ ¹⁴⁹ ¹⁵⁰ ¹⁵¹ ¹⁵² ¹⁵³ ¹⁵⁴ ¹⁵⁵ ¹⁵⁶ ¹⁵⁷ ¹⁵⁸ ¹⁵⁹ ¹⁶⁰ ¹⁶¹ ¹⁶² ¹⁶³ ¹⁶⁴ ¹⁶⁵ ¹⁶⁶ ¹⁶⁷ ¹⁶⁸ ¹⁶⁹ ¹⁷⁰ ¹⁷¹ ¹⁷² ¹⁷³ ¹⁷⁴ ¹⁷⁵ ¹⁷⁶ ¹⁷⁷ ¹⁷⁸ ¹⁷⁹ ¹⁸⁰ ¹⁸¹ ¹⁸² ¹⁸³ ¹⁸⁴ ¹⁸⁵ ¹⁸⁶ ¹⁸⁷ ¹⁸⁸ ¹⁸⁹ ¹⁹⁰ ¹⁹¹ ¹⁹² ¹⁹³ ¹⁹⁴ ¹⁹⁵ ¹⁹⁶ ¹⁹⁷ ¹⁹⁸ ¹⁹⁹ ²⁰⁰ ²⁰¹ ²⁰² ²⁰³ ²⁰⁴ ²⁰⁵ ²⁰⁶ ²⁰⁷ ²⁰⁸ ²⁰⁹ ²¹⁰ ²¹¹ ²¹² ²¹³ ²¹⁴ ²¹⁵ ²¹⁶ ²¹⁷ ²¹⁸ ²¹⁹ ²²⁰ ²²¹ ²²² ²²³ ²²⁴ ²²⁵ ²²⁶ ²²⁷ ²²⁸ ²²⁹ ²³⁰ ²³¹ ²³² ²³³ ²³⁴ ²³⁵ ²³⁶ ²³⁷ ²³⁸ ²³⁹ ²⁴⁰ ²⁴¹ ²⁴² ²⁴³ ²⁴⁴ ²⁴⁵ ²⁴⁶ ²⁴⁷ ²⁴⁸ ²⁴⁹ ²⁵⁰ ²⁵¹ ²⁵² ²⁵³ ²⁵⁴ ²⁵⁵ ²⁵⁶ ²⁵⁷ ²⁵⁸ ²⁵⁹ ²⁶⁰ ²⁶¹ ²⁶² ²⁶³ ²⁶⁴ ²⁶⁵ ²⁶⁶ ²⁶⁷ ²⁶⁸ ²⁶⁹ ²⁷⁰ ²⁷¹ ²⁷² ²⁷³ ²⁷⁴ ²⁷⁵ ²⁷⁶ ²⁷⁷ ²⁷⁸ ²⁷⁹ ²⁸⁰ ²⁸¹ ²⁸² ²⁸³ ²⁸⁴ ²⁸⁵ ²⁸⁶ ²⁸⁷ ²⁸⁸ ²⁸⁹ ²⁹⁰ ²⁹¹ ²⁹² ²⁹³ ²⁹⁴ ²⁹⁵ ²⁹⁶ ²⁹⁷ ²⁹⁸ ²⁹⁹ ³⁰⁰ ³⁰¹ ³⁰² ³⁰³ ³⁰⁴ ³⁰⁵ ³⁰⁶ ³⁰⁷ ³⁰⁸ ³⁰⁹ ³¹⁰ ³¹¹ ³¹² ³¹³ ³¹⁴ ³¹⁵ ³¹⁶ ³¹⁷ ³¹⁸ ³¹⁹ ³²⁰ ³²¹ ³²² ³²³ ³²⁴ ³²⁵ ³²⁶ ³²⁷ ³²⁸ ³²⁹ ³³⁰ ³³¹ ³³² ³³³ ³³⁴ ³³⁵ ³³⁶ ³³⁷ ³³⁸ ³³⁹ ³⁴⁰ ³⁴¹ ³⁴² ³⁴³ ³⁴⁴ ³⁴⁵ ³⁴⁶ ³⁴⁷ ³⁴⁸ ³⁴⁹ ³⁵⁰ ³⁵¹ ³⁵² ³⁵³ ³⁵⁴ ³⁵⁵ ³⁵⁶ ³⁵⁷ ³⁵⁸ ³⁵⁹ ³⁶⁰ ³⁶¹ ³⁶² ³⁶³ ³⁶⁴ ³⁶⁵ ³⁶⁶ ³⁶⁷ ³⁶⁸ ³⁶⁹ ³⁷⁰ ³⁷¹ ³⁷² ³⁷³ ³⁷⁴ ³⁷⁵ ³⁷⁶ ³⁷⁷ ³⁷⁸ ³⁷⁹ ³⁸⁰ ³⁸¹ ³⁸² ³⁸³ ³⁸⁴ ³⁸⁵ ³⁸⁶ ³⁸⁷ ³⁸⁸ ³⁸⁹ ³⁹⁰ ³⁹¹ ³⁹² ³⁹³ ³⁹⁴ ³⁹⁵ ³⁹⁶ ³⁹⁷ ³⁹⁸ ³⁹⁹ ⁴⁰⁰ ⁴⁰¹ ⁴⁰² ⁴⁰³ ⁴⁰⁴ ⁴⁰⁵ ⁴⁰⁶ ⁴⁰⁷ ⁴⁰⁸ ⁴⁰⁹ ⁴¹⁰ ⁴¹¹

[illegible]

p. 87. *Locust vabian* ^(Vesalius)
p. 88. *Loca nobilis* *Potbellia nemoralis* 116 pongs in arvensis warthweg
insularum Sanguis, omnia effusa, Moxer pons Zibella, Crogans, Lesum, id
circa Rappin. Chusachum *Potbellia emenda*: *Flora Lantia uni-bisidig*,
vini-bisidig.

1-89- ^{am} ~~subbetting~~ addatur W. 10. pl. I. p. 464. Brit. fl. ind. I. p. 268.

Pottbellia cylindrica W. & A. p. 2 f. 404
P. tenuis pubesca veta, flum. universi, filii rari carbi
Lophyia cylindrica Trin. fund. agr. p. 123. Pub. p. 16. *Pottbella pubesca* Savi. sine camp.
 syn. *Lophyia cylindrica* p. 35. *Moussiea pubesca* Beauv. ep. d'une nouv. agrost. p. 117. — Icon. Th. P. Nery

gen. gl. fr. x n. 16. p. 16. ic. 64. xl. f. 1335. Barl. ic. pl. t. 5. Buccon. imp.
x. p. 1. t. 59.

hab. in arvisij montibus circa Lom; tot' b'get, et in vulg. Lavin. N. m. 0
p. 89. post Pilerum Vin.

Nardus L.

Gen. pl. ed. II. p. 22

Char. Long laxa spicata unilaterali in rachem filiformi articulis parvis
imbric. spicula ^{ovari} glabra. Glum. ^{marginali} ~~remixta~~ breviter ex dent. rachis
prodeh. Glumula bractej, vix extensiore subula - acuminata, rugosa
stigma includente. Stamina tria. Stylus niger filiformi pubescent.
Caryopsi glumula coarctata. - Alog. Ind. ar. Dign. Syd. r.

Hardy strict L. 14 fl. 2. II. 7. 77. Herb fl. a. pr. 1. 1. 68. Feb. fl. ex. 1. 1. 78.

N. fließt zwischen Felsbrocken, große Stämme, grob, rauh, flammig,
rauh.

Don. Th. 7. Nees gen. pl. prae. x. f. ¹⁵ pub. ic. bot. xi. f. Hort ic. grana
aug. II. t. 4. Laes fl. herb. hb. 1. f. 2.

Heb. in brief voorkomende agri. pagetini; ook bij Dr. Neumann. Specie vrom.
H. aspera. 4.

p. 90 ^{part} ~~part~~ *Agrylisma orlean* L. ~~data~~ excluding *Aluk* *fig. p. 90*.
Agrylisma humilis Vrij. *fig. dalm. t. b. Maty en. gl. austr. p. 2.*

~~Aug.~~ ²⁰ ~~21~~ 1911

A. spica oblonga, cylindrica, spiculis brevibus, glomeris infimis brevissimis,
superioribus tri-angulatis, angulis glomeris illis glomerulis longioribus

Icon. Vif. fl. dlen. t I. p. 2.

Hub. ad marginem agrorum insulae Lercina, unde vixit prof. Stiles. Spicae et
herba omni glauca. Fl. acaute.

[illegible]

Obt. affing Aug. ¹ manuscript Publ. fl. ex v. 17 et ic. bot. XI f. 135h ^{quasi} Aug. neglecta Reg.

p. 92. post interim vulgo Will. h'p. (me h'p.) der st-dm Daugh.

Trichium durum Desf. fl. atl. 1. p. 114. Herb. fl. ex. p. 21.

2. *Spica compressa*, pinnatif. quadriflor., flum. valis. umbellif. oblong. lat. pedunc. tripl.
 bipinnatif. late nervosif. sub-alat. canisif. rachis tenax. margin. pilos.

Syn. *Ditium hordeiforme* Hart 12. gram. *syn. IV. p. 3.* — Non Hart l. c.
non Bieb. ? H. A. l. 1. 5

B. nylt. - *S. gnampt.* - *Lyn.* *T. vittipenn* Herb. co-gran. W. p. 4. - *Poa.* *Hyp.* *P. c. t. 6.*
Herb. cultum in insula Linnæ. Fl. apert. o.

~~Indien - Afrika - Ostindien~~ - vj

7. ~~*Spiraea cuneata*, sparsely pubescent, flowers white; sometimes ill-defined oblongif.~~

Innum. glomerum L. p. pl. d. II. p. 127. flab. fl. ex. p. 21.

9. *specie compacta, sparsely 3-4-florif, glumae velis subtransverse oblongo-lanceolatis membranaceis lobulatis multo longioribus brevifloris* ^{antheris} *bracteatis can- natis, rachitibus magis folijs.*

Non Hoit ic. gram. aeph. III - f. 31. - Myr. Pax turka.

Herb. cultum in desert. Linn. H. asph. - ①

p 98. *Hordeum bulbosum* L. minus prope circa Trai.
~~unicellul. & adnatum~~ ~~lucis remote~~

Carex leporina L. sp. pl. ed. II. p. 1381. ~~Pl. p. 1381~~ Koch 114. fl. Germ. 2. II. p. 869.

[illegible]

Syn. C. aralis Good. *trans. of Lin. soc. II. p. 148.* *Hort fl. austr. II. p. 580.* *Dignea*
leprosa Phil. *fl. ex. p. 58.* — *Non Phil. in fl. germ. VIII p. 554 opt. Saw.*
engl. bot. t. 306. Bed. fl. den. t. 1115. Leest fl. hibern. t. XIV p. 6.

Her. in parvis agri pagani ubi Cajit Kienmayer. spirales pinguet, vel
pingues - vitales. 71. aest. 24.

Carex avenaria L. sp. pl. ed. II. p. 1381. Hoff fl. austr. II. p. 588. Rich syn. fl. germ. ed. II. p. 865.

C. spica decem^{oblonga} spiralis ~~oblonga~~ ^{multiflora} confertif alba^{reperit} oblongis, ^{superioribus} ~~inferioribus~~ ^{inferioribus} foemineis, fructibus concoloribus ovatis 7-9-nerviis marginatis utroque latere bifido, cuneatis bracteis orbiculatis, fructibus aequantibus

Lyn. Vigna avenacea Rabb. fl. ex. p. 59. — Non Rabb. ic. fl. germ. VIII p. 551. Hb. ic. germ. auct. p. t. t. 49. Sav. angl. bot. t. 928. Ced. fl. Ind. t. 1266. —

Hab. in herbis et ad vias siccioribus pagorum, ubi legit Dr. Neumayer. H. aestiv.

Spica peruviana 4.
 Usp. *Chispa* ex aqua decolorum inter primas dependentium locum habet et prope mortis
 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830

p-101 post *Canem* *muniatum* L. intertext

Carex remota L. sp. gl. ed. II p. 1383. Host fl. austr. II. p. 580. ^{alpinis} in prae maribus
specie composita, spiculis ~~adrogynis~~ ^{ovatis} tenuibus rematis bractea calicem superante
suffultis, squamis approximatis, periclyps ovato-lanceolatis erectis glume ~~cauicis~~ ^{cauicibus}
in rostrum ~~peristoma~~ ^{scabrum} bidentatum attenuatis glume ovato-capitata longioribus
Syn. *Vignea remota* Rehb. fl. ex. p. 58. — Non Rehb. ic. fl. germ. VIII p. 556 et 557.

Low-engl. bot. t. 832. Host ic. germ. austr. I. t. 52. Less fl. herb. t. XV p. 1.
Hab. in locis umbratis aditioris *Agajunae* ubi legit Fr. Neumayer. Spiculae
stramineae. Fl. Majo, Junio. 4.

5 - p. 102 - *Loc. nobile Canus digitatae* L. additur mons J. Offero.

102 - Caricby viny aduna post C. Linkii

~~ax P. larva pharynx very~~
~~max with that~~

Corax nictitans ^{1. p. 53. fl. apr. - p. 594. 2nd. fl. ex.}

p. 65.
 pro asperata sublinearata solitaria sessile, femineis subgeminis strobis
 difformibus, infra pedunculata, fructibus subglobosis, viridibus, ~~strobis~~
 brevissimis iniquiliter breviter, bracteolis late ovatis strobis marginem
 h. t. bracteis foliaceis basi seminigeris late ovatis

Syn. C. verrea Schkbr. hist. nat. Car. p. 115. Host fl. austr. et progn. num. 614.
- Ten Rhod. ic. p. germ. VIII f. 641. Host ic. germ. austr. t. 181 et IV
t. 86. Schkbr. l. c. tab. 1. f. 46. et t. opp. f. 156.

Hb. in collibus apricis apricis circa Schenico. Plant. Apr. Woj. Specimen
 minutely ferrugineis, hirsutibus, marginibus albidis. 4.
 Obs. specus hinc fructibus, granulis et tamen globulis nec conglomeratis praeditis
 et symmetris, tritibus donata. inconstanter diametris ex aequo
 fructus depressi pro vixis a laniibus separandi probat, et in
 ter utroque ambigit.

Carex distans L. sp. pl. ed. II p. 138. ~~Hort. fl. rom. II p. 199. Febb. fl. ex.~~
p. 66. Alch. fl. jedr. p. 223. Pet. bot. univ. v. 203.
fl. dist. adnata, formis, subternis, obliq. breviter

[illegible]

Buteola orla Engelm.
Don. Flab. ii. fig. germ. VIII p. 62. Torr. angl. bot. t. (234). Hort. ic. germ.
~~after 1 + 22~~ Schk. hist. des Car. no 7 et Pl. n. 68.

~~1841~~ 1842. Schk. n. sp.
 Hab. in part. com. part. Kuin, Derrif, Lebenco, Spalato, Salva
 et Poyuz. ^{et in ignis loco.} Sp. macula ferruginea aut b.
 dia, femineae ex badio viridigen variae. 2.

Carex glauca Scop. fl. carn. II. p. 223. pub. fl. ex p. 66. Pet. bot. ungu.
n. 266.

n. 266.
C. spicae vaginatis antheris, foeniculis duobus pendulis, bracteis plura-
bus submarginatis brevioribus, utrisque submarginatis pedunculatis
terribus, fructibus oblongis pennatis - calyce oblongo integro, ex-
trinsecis, bracteis ovatis antheris subaequalibus.

C. flacca Schreb. spirit. fl. tips. aff. n. 89. Host ii. germ. austr.
 f. 66 et fl. austr. II. p. 66 Schkr. Hoffm. var. p. 182. n.
 fl. ii. fl. germ. VIII. p. 648. Host ii. germ. I. t. 90. Low. engl. bot.
 1806 pub. (recurva Good.) Schkr. l. c. tab. O. P. 57 et 22 n. 113.

Hib. ad fopay et invaly et in partu udy totus Holmatis H. Agn. Mojo. pua

Obs. Differs a *C. caespitosa* L. (caespitosa Dej. (Holtz. ic. Fl. germ. VIII. t. CXXXIX)) principally in
leaves being more long, very prominent large perianth lobes generally are convex suborbately
oblongate - Here perhaps *affinis* with *C. verrucosa* Vill. *C. pinetis* Webb. Fl. gar. p. 222 near
speciosa

Carex hirta L. 17th ed. 1789. Hort. fl. austr. II. p. 511. Fl. ex. p.

C. pilosa, spicis masculis approximatis geminis, feminis utrinque diffunditis breviter pilosis longipennis saepeante multo brevioribus, ^{bracteis} ~~bracteis~~ bracteis pedunculatis oblongis, fructibus ovato-oblongis stratis, ^{bracteis} ~~bracteis~~ bracteis longipennis, ^{bracteis} ~~bracteis~~ bracteis longipennis, ^{bracteis} ~~bracteis~~ bracteis longipennis.

Syn. *C. hordeiformis* Pers. bot. univ. II. 207. var. alior. et ex. 194. — Non. fl. ex. p. 628. Sav. angl. bot. t. 585. Hort. ic. grm. austr. I. t. 76. Schkr. hyst. det. var. tab. III. n. 108. Less. fl. herb. t. XVI. p. 3.

Var. β . *hordeiformis*, Pers. syn. gl. II. p. 548: foliis virginibus globosis — Non. Ped. fl. det. t. 1711.

Hab. circa lacum Bozagnappe. Loci ubi prope Zart promissa cum exultante. Spicae masculae dilute ferrugineae, femineae flavo-virides, dimissa percae. Fl. Apr. Mayo. &.

Obs. Non Lessii pectus bracteis fructibus longioribus.

Carex hordeiformis Vill. hyst. det. gl. det. Deyh. II. p. 221. Koch. syn. fl. grm. II. t. p. 583.

C. glabra, spicis masculis approximatis geminis, feminis utrinque diffunditis breviter pilosis longipennis saepeante multo brevioribus, ^{bracteis} ~~bracteis~~ bracteis pedunculatis oblongis, fructibus ovato-oblongis stratis, ^{bracteis} ~~bracteis~~ bracteis longipennis, ^{bracteis} ~~bracteis~~ bracteis longipennis, ^{bracteis} ~~bracteis~~ bracteis longipennis.

Syn. *Carex rupestris* Wahlenb. in act. botan. 1805. p. 151. Hort. fl. austr. II. p. 610. Fl. ex. p. 62. *C. hordeiformis* Wahlenb. l. c. p. 152. Hort. ic. grm. austr. I. t. 76. — Non. fl. ex. p. 628. Schkr. l. c. t. II. n. 55. Hort. ic. grm. I. t. 76.

Hab. in uliginosis circa Norenta ubi legit Dr. Neumayer. Spicae flavae, t. p. 4.

Obs. Interdum pedunculi inferioris spicas geminas sustinent.

* *Carex hirta* L. 17th ed. 1789. Hort. fl. austr. II. p. 511. Fl. ex. p. 628. Sav. angl. bot. t. 585. Hort. ic. grm. austr. I. t. 76. Schkr. hyst. det. var. tab. III. n. 108. Less. fl. herb. t. XVI. p. 3. Var. β . *hordeiformis*, Pers. syn. gl. II. p. 548: foliis virginibus globosis — Non. Ped. fl. det. t. 1711. Hab. circa lacum Bozagnappe. Loci ubi prope Zart promissa cum exultante. Spicae masculae dilute ferrugineae, femineae flavo-virides, dimissa percae. Fl. Apr. Mayo. &.

Fr 103 ~~Polk~~ (vernom Vill. adendas ment

p. 103 ~~1014~~ ¹⁰¹⁴ C. venusta Vill. ad Ardenas mont
~~1014~~ Carex tomentosa L. Mantz. p. 123. Hort fl. austr. II. p. 600. Philb. Henc. p. 63.

[illegible]

Don. Ridd. ii. fl. germ. VIII f. 638. - Nov. engl. bot. t. 2046. - Schkbr. herb. R. Car. ^{tbl. 7} 2. 28
Leers fl. herb. t. XV f. 7.

Hal. in part only ad Petropolis prope Lernip. H. Majo. Vener. Specie
significaty. 7.

Carex riparia Curt fl. Lond. fasc. IV. Hort fl. austr. #. p. 607. Philb fl. ex. p. 70.

Aluh. fl. jadr. p. 223

Alb. fl. par. p. 245

C. spica masculis & feminis sessilibus, feminis subteris oblongo-cylindricis, imbricatis, pedunculatis, bracteatis bracteis foliaceis late longioribus basi vaginae multum brevibus, bracteis cuneatis, utrinque multum brevibus, in rostrum breviter productis, bracteis lanceolatis cuspidatis brevibus.

Syn. *C. confusa* Ehrh. *beig.* IV. p. 43. Herb. ii. gramin. aeph. I. p. 1. - Non
 fecht. ii. fl. germ. VIII. p. 647. Herb. ii. gramin. aeph. I. p. 93. Law. engl. bot.
 + 519. *conf.* fl. herb. I. VII. p. 2. I.

Herb. ad fossas, nigras, lacum abque fluvium totius Solanaceae. 16 April, 1891.
 Spicae maritima foragineae nitidae: semineae crassae nigrescentibus
 abo-volacae 4.

Carex angustata Gooden. in trans. of the linn Soc. II p. 207. West fl. asept. st.
p. 600. West fl. asept. st.

(C) spica ^{nervosa} majusculis 1-3, fovearum subtiliter cylindricis pedunculatis, bractea folium
longiora unguata, fructibus siliosis inflatis glabris corpe nervosissimis,
in rostrum lineare truncatum contractis, bractea oblonga longioribus

syn. *C. venaria* β L. 10. pl. ed. II. p. 1359. *C. venaria* both text & fl. germ. II. 21.
466 Less. fl. herb. p. — non pub. & fl. germ. VIII. p. 659. off.
Sax. engl. bot. t. 180. Hort. ic. gram. austr. t. 89. Less. l. c. tab. XVI. f. II.

Herb. Leipzigerbotan. vasa Herbaria ubi legit Fr. Neesmaeyer. Specie str.
minuae. 76. Moji, Linn: 4

108 post *Heliocharis palustris* *fl. Br. prod. fl. nov. Hb. I. p. 80. intertextus*
Heliocharis ovata *fl. Br. prod. fl. nov. Hb. I. p. 80. Poem. et. Ich. 19th. reg. II. p. 152*
Zab. fl. ex. p. 77

[illegible]

p III. *Cymoides* aede
Eriophorum L.
Gen. pl. d. II. p. 21.

Char. Hermafrodita, caulis breviter foliaceus, involucribus, bracteol. multifloris
involucris subaequalibus, inferius intermedium floribus ~~subaequalibus~~
et subaequalibus bracteol. duobus subaequalibus. Stamina tria, stylus
filiformis, triplicis, stylis subaequalibus, lanceolata trigona subaequalibus
postfloribus in lumen elongatis obducta. — *Herf. Maritima* C. D. Monog. p. 101

Eriophorum angustifolium Roth. tent. fl. germ. t. p. 24. Herb. fl. auct. t. p. 83.
fl. ob. fl. ex p. 29.

E. culmo subterre, foliis lanatis, canaliculatis apice acuminato-triplicis,
spicis aggregatis, filamentis brevibus.

Syn. E. polytrichon ^{fl. ex p. 29} Herb. fl. germ. auct. t. p. 29 — Non fl. ob. fl. germ. VIII
Th. 7. New gen. p. 18. n. 16. f. 1-7
fl. ob. fl. germ. t. p. 24. l. t. 54. l. d. fl. d. t. 1442. Herb. fl. germ. p. 18. n. 16. f. 1-7
f. 1. a

Herb. in p. 18. n. 16. f. 1-7. Narante ubi legit Dr. Neumayer. Spiculae ferrugineae,
luna candida. fl. vire 24.

p. 113 post *Junium lampocarpum* Roth.
Junium obtusiflorum Ehrh. bot. VI. p. 88. Reb. fl. ex. p. 96.

J. culmis filicis, foliis nodulosis teretibus, cum decompositione divergenti, perianthiis
phyllis aequalibus, omnibus apice rotundato-obtusis capsulae ovato-trigonae
acutae subaequalibus.

Syn. J. sylvestris Herb. ic. germ. auct. III. p. 57 et fl. auct. t. p. 442. — Non
fl. ob. fl. germ. X. f. 901. qd. l. w. encl. bot. t. 2144. Herb. fl. auct. III. t. 86.
ic. gram.

Herb. in p. 18. n. 16. f. 1-7. Narante ubi legit Math. Bollen. Flor. viridis,
capsulae c. flavido ferrugineae. fl. May, Junio. 24.

p. 113 post *Junium insulense* W. W.

Junium desponsum L. sp. pl. ed. II. p. 466. Herb. fl. auct. t. p. 449. Reb. fl. ex. p. 96.

J. culmis filicis, foliis filiformibus subaequalibus, floribus omnibus
longe alaribus distinctis, perianthiis phyllis ^{caput d. d. longioris} eximie acuminatis, interius
acutis brevibus, capsulae ~~obovatae~~

Syn. fl. ob. fl. germ. X. f. 822-26. l. w. encl. bot. t. 802. Herb. ic. germ. III. t. 90.
Loefl. fl. herb. t. XIII f. 8.

Herb. in agris humidis, insulae ^{insulae Siciliae} *Leina* a *Pizzia*, et circa *Narante*. fl. haec
nec fl. vire 24 at forma ejusdem contracta fieri videtur

Obs. *Junium insulense* W. W. hunc unum proximum *Junium insulense* fl. ob. fl. germ. X. f. 822-26. l. w. encl. bot. t. 802. Herb. ic. germ. III. t. 90.
at phyllis perianthiis ^{et filamentis} ~~subaequalibus~~ aequae distinctis.
J. insulense ^{insulae Siciliae} *Leina* a *Pizzia*, et circa *Narante* ubi legit Math. Bollen. Flor. viridis,
germ. 18 f. 877-881.

Lunney Denageja L. fl. 1798 p. 208. Hort fl. austr. I p. 449. Rhod. fl.
exc. p. 95. exl. 194. Naes.

I. culmis floris robustis cymosis flis pinnatis, cymis raris
diversis floris raris cymosis, perianthii floris ordo-lancea
lris capsula subrotunda obis sublongioribus.

Leon Rhod. ic. fl. germ. X. p. 923. Od. fl. den. t. 1160. Hort ic. germ. austr. III
t. 91. Vaill. bot. parisi. tab. XX f. 1.

Hab. in campis humidis circa Seta in insula Lania, unde missa Al. Stali. fl.

p. 118 straminei. fl. m. ①
p. 120. D. Will. Herbst in sua Concom. Spargi (Bot. reg. Dec. 1845 p. 83)
qui est Rhod. ic. fl. germ. IX. p. 796. 797.
nisi cum Dabio Concom. Pallarii desinunt (C. Virgineum Herb.) ad
C. radiatum suum. C. Virgineum ex ipso differt a C. Pallarii Des.
et a C. Thomasii Des. paucis later, quod sane parum!

p. 123 Gladstus triphyllus Roth. orbus etiam in regalis insula Curcola
circumscriptus, et cum hoc idem f. Doucheum Schlecht.

p. 129. Panormum ~~officinale~~ ventosum L. orbus prope capite in ora
extrema insula Lania, et a phanerogolis floris in insularibus
ulgo adhibetur pro Sella ventosa L.

p. 131. Intillaria pyrenaica L. ff. fl. 2. 4 p. 436. Schlecht. bot. p. 149.
~~ex tab. 308~~ Gaert. in Lest. bot. mag. t. 664. Rhod. ic. bot. X. p. 7. et
tab. f. 976. est alia species perianthii laciniis apice coriaceis inermi-
bus reflexis inaequalibus, id hanc notam ~~non~~ ^{me} ~~semper~~ ^{omnibus} in specimenibus
pyrenaeis a d. Pettenbacheris pro sua f. pyrenaica habitis
fructu quassiffi doles, cum ovum e contra Potallion non
tam vixitum tantum exhiberint, nempe notam var. β.

p. 128 post Navium vivum Lort. addit.
~~p. 128~~ Navium polyanthos Lort. rech. sur les Navium. p. 36. et fl. gall. I. p.
236. Rhod. fl. exc. p. 89.

N. flis late linearibus flans obis inaequalibus, raris angustis 8-12 floris,
pedunculis angularibus, perianthii laciniis oris ^{conspicuis} flis ^{conspicuis} corona
cyathiformi peltulata subrepanda duobus longioribus.

Leon Rhod. ic. fl. germ. 18 f. 814 opt. Lobel. in. 149. Barm. ic. fl. 915. Clus.
hist. fl. 49. 155 f. 1. 2.

Hab. in saxis apris napoleonum circa Lania. fl. jam Novembri et Decembri.
Rhod. 7.

Obs. Differt a praedicta cui vixit fructibus prope forma ordo-elliptica, nec
orbo-lanceolata laciniarum perianthii, caryopis brevitate.

p. 127. post Navium pectinatum L. insensitum

Pipileum stansburyi aequilongum, stylis ovario subglobosis brevius longior, stigma
 nuxta. — Stamina brev. filamento connatis et nunc alterius filamen-
 tibus species brev. filamento connatis. *Podocarpus* cum *Pont.*

p. 141. *Allis marginatus* Little. add. icon. 7/66. 10 ft. germ. XI. r. 1887.
p. 142. *Ornithogalum* add.
Ornithogalum pyrenaicum L. sp. pl. ed. 2 p. 440 ex herb! Berol. fl. ital. 18
p. 106.

0. folij canaliculatis unicoloribus glabris, nervis longissimis, pedunculis
fructibus ^{superius} striatis, ~~longioribus~~ ^{apice} bracte ovato-^{apice} reticulatis brevioribus, plerumque
ovato-lanceolatis apice angustatis.

Orthocentrus affinis angustif.

Syn. *O. sulphureum* Steen. et Sch. 1866. reg. VII p. 518. *O. tachyoides* Hart. pl. afr. t. p. 432. non act. *anthurium* *sulphureum* Waldb. et Kit. pl. var. long. t. p. 98. — non. *Orth. ic. fl. germ.* XI p. 1027. Waldb. et Kit. l. c. tab. 95.

Sim in *East. bot. mag.* f. 2623
Dodon. pempt. p. 209. f. 1. et in *insule Otero* prope *Lollen* grand
Herb. in ~~1667~~ *prope* *Namata* ~~unde~~ *unde* *Mr. Botani.* *Flora* *e* *irrit.*
flor. pempt. p. 209. f. 1.

Hyacinthus ornatus var. β . phyllanthus = brevis in flis linearibus
 Lycopodium vacuatum superantia productis.

Coryphaea nemoralis supersede potius.
Habui inquam hanc abominatam ex aquis circa Althoracina a prop. Loria
quo ipse ibi specus riber sponte.

p. 152. *Asphodelina cretica* Vrf. oviter et in sinula Lesina

p. 160 *Asparagus officinalis* L. subjugatus

Asparagus tenuifolius Lam. enc. bot. 2. de Pad. t. p. 291. Richb. fl.
exc. p. 118.

exc. p. 118.
Q. caule herbaceo. lami erecto, foliis parvulatis ~~et~~ sagittatis nervulosis ter-
tibus lacubus, filamentis antheris subrotunda multis longioribus.

Syn. *A. officinalis* p. L. sp. pl. 2. II. p. 448. *A. sylvatica* Walst. et Kit. pl. rar. burg. III p. 223. ~~ut~~ *Hoff* fl. austr. I. p. 439. — *Rein.* Walst. et Kit. l. c. t. 201. *Mab.* hist. in *Dion.* p. 506.

Kit. l. c. t. 201. *Mar. 1881.*
Hab. in umbrosij vltij Bessa in insula Vaglin, ubi legit Doct. Venetner.
H. a om in aestivo. H. pallid. 4.

p. 123. part *Orchidium conopsea* L. addent. etc.

Orchis odoratissima L. f. p. ed. II. p. 1335. Host fl. austr. II. p.

③ *Tabella integrum trilobis, lobis ovatis oblongis, perianthii tenuis, ovae extrorsae per
tuppinis, calice recurvo filiformi annulo subaequantibus bracteis binerviis
minime subaequantibus, bracteis cylindricis elongatis densis, foliis linearibus tenuibus,*

Syn. Euphaderia ~~sp. nov.~~ *adstringens* Rich. in mem. du Mus. IV. p. 38. Pl. ex.
p. 121. — Don. *Reich. bot. VI. f. 814.* Hall. *ic. pl. helv. t. XXVIII. f. 1. var. fl. sup.*

p. 166. lin. 9. *Platanthera bifida* *P. clavata* est *Pl. chlorantha* Curt. pub.
- lin. 24. loc. *absentibus* var. *absentibus*.
p. 108 post *Orchidum provinciale* Barb.

p. 167 post *Orchitum provinciale* Balb

Orchis mascula L. sp. pl. ed. II. p. 1333. Nöb. fl. exc. p. 123. ext. ic. Lag.

[illegible]

tab. 39. f. 12.
 Var. β . spiciosa: lobellis inciso-dentatis, perianthii laciniis ^{apice} acuminato-subulatis.
 — Syn. *O. spiciosa* Host fl. austr. II. p. 327. (tab. fl. ex. p. 140¹⁸) *O.*
maguel Jacq. ic. riv. I. p. 17. *O. palustris* Alsch. fl. jard. p. 210
 ex herb! — Non Jacq. l. c. tab. 180

ex herb! — non suff.
Hab. in partib. aquisq. palustribus circa Lava, et in sylvestrib. circa Caput.
Flous purpurei. Fl. Majo, Junio. &.

Flora purpurea. p. 1. 1. 1.
 Obs. Calcar tunc in specie tunc in varietate ledit ascendens vel descendens,
 folia oblonga, vel lanceolata, lobellum flexu minusve magis inaequa-
 le, medio ~~dentato~~ subrepuliter denticulato, vel biplo cum du-
 bus interjectis, vel denticul. repulente.
Flora purpurea. p. 1. 1. 1.

p. 168. Ad Archim palustrem Læg. dale ^{citatum} ~~specimen~~ ^{Flora jadrans,}
 2. B. muscivora B. speciosa pertinet.

p-173
genus *B. mosnanae* *B. speciosa*
part *Orchidum coroffham* L.
part *Orchidum* intera *speciosa* genus

p. 173 post orchid
p. 174 post orchid inverte species group
Tinea Bion.

Tinea Bivon.

Tinea Bion.
Ex fam. *Orchid.* nov. gen. in Giorn. d. sc. e lett. Ital. ~~1833~~ 1833. p. 149.

[illegible]

Tinea maculata vrf.

Linea maculata Vg.
Syn. *Latyrum maculatum* Desf. fl. atl. II. p. 219. *Orchis atlantica* W. p. pl. IV. p. 42.
Berol. d. var. ital. Dec. II. p. 42. et Amsen. ital. p. 82. Glycer.

O. secundiflora Benth. *fl. rar. Ital. Sic. II. p. 42.* et *univ. Ital. p. 10.* *Spag.*
densiflora Desf. *choix de cost. de Tournef. p. 11.* *Pinia cylindracea* Bion.
densiflora Desf. *choix de cost. de Tournef. p. 11.* *Pinia cylindracea* Bion.
densiflora Desf. *choix de cost. de Tournef. p. 11.* *Pinia cylindracea* Bion.

Demiflora *desf.* *inca* *de* *ca.*
l.c. *Todar.* *orchid.* *ficul.* *p. 7.* *Grav.* *syn. fl. sic.* *II. p. 540.*
Son. *Desf.* *l.c.* *st. 6.* *Boudan.* *gaulh. sic.* *t. 32 fl.* *t. 138 fruit.*

Icon. Desf. l. c. tab. 6. Bondu. pangh. sic. t. 32. fol. t. 138. fruct.

p. 173. post Orchidem conopsea L.

Orchis alba Scop. fl. carn. ed. II. p. 201. Hort fl. austr. II. p. 534.

B. labello integerrimo trilobo, lobis auratis intermediis majore obtuso, perianthii laciniis
ovato-oblongis conniventibus, calice crassiusculo pendulo ovario triglo breviori,
bracteis trinerviis ovarium aequantibus spica elongata laxiuscula, foliis lanceo-
latis, radicibus tuberibus fasciculatis.

Syn. *Gynadenia alba* Rich. mim. du Mex. IV. p. 35. *Habenaria alba* R. Br. in Ait. h.

Kew. ed. II. v. 3. p. 195. *Reichb. fl. exc. p.* 130. *Satyrium albidum* L. sp. pl. ed. II. p. 1338.

— Icon. Th. F. Nees gen. pl. fasc. V. n. 6. f. 13-20. Sow. engl. bot. t. 505. Oed. fl. dan.
t. 115. Hall. ic. pl. helv. tab. XXV. f. 1.

Hab. in graminosis montibus ditioribus Napaeinae, unde misit Fr. Neumayer. Fl. parvi albae.
Fl. aequalis. 4.

Haab. in paucis insulae Lesina, et circa Sagulum. Fl. carni lividius rubellus
pichi, suaviter. Fl. virens. 4.

Ob. Radix fibrosa ramentibus ordo-substantia^{integr} geminis. Caulis erectus pro-jectus
tunc foliatus, superius subinde compressus-angulatus, rubrum per se maculatus.
Hinc^{per se maculatus} ramenta hinc majora patenti obliqua, superius erecta obliqua-lanceolata
ante scopin vivos. Flores exigui in spicam cylindricam^{recundam} densam congesti.
Bractea membranacea ordo-lanceolata acuminata ovato brevior. Petala
hinc breviora aucta connatis et infimus inclinata. Labelli breviora
breviter stipitata, medio lineari^{late} apice obliqua^{aut} hinc-^{late} infima. Gynae-
cium ellipticum rectum. — Semina hinc depressa ab Ovulid, et prope ab
Acroate fl. br. cui valde simile ovato recto, et granulatis pollinibus
flo. superius: ab Acroate ramenta differt labelli calcarato. — Nomen
speciem jam ab anno 1800 a cl. Desfontaines plantae a se
firmum invenitae impositionem prioribus jure servandum. Nomen
autem generum quod juxta legendi promissionem convenit
posuit in nomen animalculi nunc nostri, aequius proinde scribi
nuncitur *Tivaca*.

p. 125 Ante *Spiranthes* Rich. inseritur

Goodera fl. Brown

in *Art. h. Kew. ed. III. t. V. p. 197.*

Char. Perianthium ringens rotans laciniis interioribus connatis, rectis extor-
tus patulis, labelli porrecto lobi concavo et in ligulem canaliculatum acuminato
calcarato. Gynostemium brevius erectum. Anthera libera loculis divortis. Pollina
vix hinc caudata. Staminodia nulla. Nothium bilamellatum. Stigma subordi-
natum^{ovarium rectum} Clasi. *Synandria* Ord. *Diandria* Syst. sex.
tum marginatum^{Host fl. austr. II. p. 541}

Goodera regens fl. br. t. c. p. 198. *Haab. fl. ex. p. 131.*

fl. foliis radicalibus^{orbs} pilolatis^{orbs} vixis-
vixis-
recundis pubescentibus, perianthii laciniis labellum lanceolatis.

Syn. *Schizium* regens L. sp. pl. ed. II. p. 1839. *Neottia* regens Sw. act. holm. 1800 p.
226. — Don. *Nov. engl. bot. t. 289.* *Tag. fl. austr. IV t. 369.* *Hall. ic. pl. helv. t.*
XXII. f. media.

Haab. in sylvestribus montani Ditionis *Tagessera*, unde sine loci specialis indicatione misit
fl. Neumayer. fl. albid. fl. deserte. 4.

p. 177. ad *Ophrydium* asipsum Huds. add. iconem *Pedus* en. orb. in *Neil. t. 2. f. 1-2*, et ad
O. brachylophum Lk. ic. *Pedus. ib. t. 2. f. 3-4.*

Pag. 179. Post *Ophrydem flavicantem* Vis. addenda

Ophrys *Dommasinii* Vis.

O. ^{bas. integro,} ~~labello~~ ^{circum} elliptico, medio glabro nervoso, ~~lanceolato~~ citrino-piloso, ~~bas. integro,~~
~~apice~~ apice emarginato, laciniis perianthii internis late lanceolatis planis
glabris gynostemium cernuum apice oblique attenuatum subaequantibus.
Hab. in apris saxosis insulae San Pier d' Nembro prope Lecco fl. parvi
rubro-purpureo labello circum circa piloso-citrino, dorso lineis brevibus lineis brevibus
~~parallelis~~ formam litterarum X simulantibus picto. fl. April. 2.

Descr. Tubum bina subtendens aut elliptica, fibris radicalibus crassis circumdata.
Folia infima ovato-lanceolata ^{acuta}, caulina vaginantis adpressa ~~multa~~
lanceolata-acuminata. ~~Chapm.~~ spithameus, repens 3-5 florig. Bractea
lanceolata-linearis florum subaequant. florig. labello bilineari. Perianthii laciniis
externae viridulae oblique. — Similis quodammodo *Ophrydi* arvensis (Thunb.)
que differt labello foveoliformi non ad medium angustato basi breviter perianthii
laciniis interioribus gynostemio ^{cernuo obliquo} ~~perfecto~~ ^{perfecto} ~~rotundato~~ ^{rotundato} ~~obliquo~~ ^{obliquo} ~~longioribus~~
nec longioribus, et maculae labelli forma. ~~Thunb.~~ ^{habitu} ~~quodammodo~~ refert icon
1154 icon bot. cent. IX cl. Schb., sed differt gynostemio ~~perfecto~~ et forma
maculae ~~f.~~ Similis quoque *O. arvensis* Schb. fl. succ. p. 140. et icon.
bot. f. 1153, ^{ix.} ~~ut~~ tamen differt labello ~~rotundato~~ ^{obliquo} ~~obtusato~~ nec emarginato,
~~et gynostemio~~ ^{et macula} ~~perfecto~~ ^{ut in O. arvensis}. *Hybridam* ~~Thunb.~~ ex ~~describ.~~ ^{esse} ~~hinc~~ ^{genitam} ~~putat~~ ^{esse} cl. Schb. fil. in litt. ad cl. Dommasinium, sed ~~pl.~~
~~hybridam~~ ^{vide} ~~in Orchideis~~ ^{possibilem aut minimam} ~~esse~~ ^{difficultatem} ~~credo~~, ~~pollinis~~ ^{pollinis} ~~naturae~~ ^{naturae} ~~adversae~~
sante. Erant ~~scilicet~~ ^{scilicet} ~~loci~~ ^{loci} ~~citati~~ ^{citati}, unde ~~circiter~~ ^{circiter} ~~hinc~~ ^{hinc} ~~specimina~~ ^{specimina} ~~bona~~
tunc cl. Dommasini ~~per~~ ^{per} ~~ultram~~ ^{ultram} ~~transmissionem~~ ^{transmissionem} ~~inducunt~~ ^{ad O. arvensiformem}
inducunt praebent.

mandata a Lippi ad Thunb. pro a tunc Lippi.

et de Lippi Botterii p. 16.

Lemna L.

Gen. pl. ed. II. p. 396.

Char. Decm. ac. Ordrip.

Lemna gibba L. sp. pl. ed. II. p. 1377. Hort fl. austr. II. p. 33. Pchb. fl. germ. p. 10.

L. caulis articulis foliiformibus obortis super ^{compressis} ~~planis~~ ^{hemisphaericis} ~~subtis~~ ^{concrevis} ~~gongio~~ ^{is}, ^{radice} solitaria.

Syn. *Delmatophace gibba* Schleid. prod. mon. Lemn. in Linnaea vol. XIII p. 391. — Don.

Th 7. Nées gen. pl. fasc. VI. n. 9. ^{p. 13-18} Pchb. ic. fl. germ. VII. tab. XIV. Sav. engl. bot.

t. 1233.

Hab. in aquariis botanorum circa Sclavico. Vene. C.

p. 189. ^{Apote} ~~Act~~ *Loterum nanum* Roth addenda

Lotum marina L. sp. pl. ed. II. p. 1374. Hort fl. austr. I. p. 1. Pchb. fl. exc. p. 6.

L. caule teretifaculo, foliis linearibus, spathe foliaceae pedunculo superne latiore.

Don. Pchb. ic. fl. germ. VII. tab. IV. Th 7. Nées gen. pl. VI. n. 14. p. 15-26. Lam. circ. bot. p. 135. Sav. engl. bot. t. 467.

Hab. in fundo maris tunc orae Dalmatiae, praesentia circa Zara et Sclavico. fl. aestale. 2.

2 p. 190. Pot. *Potamogeton luculentum* add. ejusdem

var. *P. longifolia*: foliis lanceolatis elongatis spinuloso-acuminatis — Syn. *P. longifolius* Gay in Bot. enc. bot. suppl. IV. p. 555. Pchb. ic. fl. germ. VII. p. 24. *P. macrophyllus* Wolff. in Noem. et Sch. 1958. vj. mant. p. 358. — Don Pchb. l. c. tab. XI. p. 70.

Hab. in fluvio Narvata cum quercu.

p. 205. lin. 22 post nec ne add. = *punctum*

ib. lin. 43. post icones add. = *Illyr. Tidovina*. =

p. 240. lin. 22 post vstra lanceolatis integris, nervis spicis elongatis et.

Vol. II

p. 8. Ante *Stichem cancellatam* Bernh. cui proae prioritate praefari debet St. dictyophora Tausch. in syll. pl. a Societ. vob. 1836. ed. ¹⁸³⁸ ~~1836~~ Don. II. p. 254 inferenda est

Statice minuta L. mant. pl. p. 59 (excl. syn.) *P. pubescens* Boiss. in DC. prod. 1958. nat. XII. p. 655.

S. pubescens, foliis lanceolatis-obovatis oblongis basi longius attenuatis univerviis oblongis recurvis, raso, tereti a basi ramoso, nervis articulis parvis erectis, infirmis sterilibus simplicibus, floribus laxo breviterque spicatis, dentibus albis oblongis.

Syn. *St. pubescens* Pchb. ic. bot. II. p. 89 et fl. exc. p. 191. na DC. et excl. syn. et icon. — Don Pchb. l. c.

pl. 182 f. 326

maritimum

Hab. in sacris agris, circa pagum Nervinus in insula Osero, ubi legit doct. Sandner.

um frequenter crescentem

fl. lilacini fl. autumnus t.

cei
m
c

Obs. Hinc similis Statice pedunculata DC. differt praesertim ramis sterilibus
dorsif. bractea superiora ^{nigram ex hirs} ~~valis~~ ^{pedunculata} ~~repente~~ et angustis marginatis,
speculis confertis nec densis.

p. 11. Echinosi Neumayeri videri similis est E. Daggetianus Boiss. et Heldr. Viagn. fl.

et foran idem

juv. orient. VI. p. 97.

p. 79. Lure sit Anthem. salmationis Scheele in Linnaea XVIII. p. 464 ne propter latet.

p. 80. Lure sit Anthem. salmationis Scheele in Linnaea XVIII. p. 464 ne propter latet.

p. 84. Lure sit Anthem. salmationis Scheele in Linnaea XVIII. p. 464 ne propter latet.

confer propter observationes d. Koch in syn. fl. germ. d. II. p. 1026 usque
in Formis italicae anni praecedentis editae informantis.

p. 85. Chamaemela quibus hic loci enumeratis super Chamaemelum inas-
dorem var. b. maritimum et addere sequentis.

p. 348.

fl. cav.

Chamaemelum maritimum Vis. sem. var. h. patav. 1846 p. 4.

Ch. acheniis cuneatis subtriangulis sub dorsif apice impresso-bislandulosis,
pappi brevissimo, disco epigyni terminali latitudinem apicis achenii
aequante, nigro bidentato, costis obtusis concoloribus approximatis.

nto. 7.

ys in

Syn. Chrysanthemum inodorum B. L. fl. pl. ed. II. p. 1283. Chamaemelum
inodorum B. maritimum Vis. fl. dalm. II. p. 85. Drifloropernum
maritimum Koch syn. fl. germ. d. II. p. 1026.

Hab. ad litora maris septentrionalis et baltici.

Obs. In Ch. inodoro hinc simili acheniorum costae sunt acutae, disco
longi et distinctae: lacinae etiam plerumque brevioresque.

Chamaemelum discoides Vis. sem. var. h. patav. 1846 p. 4.

Ch. acheniis cuneatis subtriangulis in dorsif apice impresso-bislandulosis,
pappi brevissimo, disco epigyni terminali latitudinem apicis achenii
aequante obsolete bidentato, costis acutiusculis discoloribus distinctis.

Syn. Matricaria discidea DC. prod. syst. nat. VI. p. 50.

Hab. in California

Chamaemelum confusum Vis. loc. cit. quae est Matricaria confusa
H. Perssonii.

p. 101. Picris laevigata Vis. est eadem ac P. Sprengeriana Alch. fl. jard.
p. 206 ex herb. non Lam. Ontus et in monte Dinam ubi vocatur Mtjizra
ex herb. non DC.

p. 103. Thymus hirs. Alch. fl. jard. p. 206 est idem ac Leontodon hispidum.

p. 104. Leontodon inermis L. v. Leontodon inermis L.

p. 113. Ad Lactucan Scardium L. addendum est etiam L. quercina Alch.
fl. jard. p. 205 ex herb. non L.

Pag. 49. Post *Cirsium palustre* Scop. inserto

Cirsium pannonicum Gaed. fl. helv. VI. p. 363.

C. caule superne nudo ~~apice subramoso~~^{subramoso} ~~mono~~^{tri}trifido, foliis ~~super~~ oblongo-lanceola-
tis indivisis spinuloso-ciliatis, infimis in petiolum attenuatis, caulibus semiaflexi-
caulibus oblique breviterque decurrentibus, capitulis ¹⁻³ longe pedunculatis, ovatis, squamis
orbis-lanceolatis ^{apice} ~~spinuloso~~^{spinuloso}-mucronatis recurvis.

Syn. *Cnicus pannonicus* Host fl. austr. II. p. 444. *Carduus pannonicus* L. fl. suppl. p. 348.
et excl. syn. Frut.

C. serratuloides Jacq. fl. austr. II. p. 16. non L. *Cirsium serratuloides* Scop. fl. carn.
Rehb. fl. ex. p. 284.
II. p. 127. — Jacq. l. c. hb. (127). Clus. hist. pl. v. p. CXLVIII f. 1.

Hab. in partibus montis Dinan ubi legitur ^{legit} ~~exstitit~~ Doct. A. Mazzoleni. fl. perquaci. fl. asplen. 2.

Obs. *Cirsium caucum*^{hib.} huic affine distinguitur foliis radicalibus incrassatis fusiformibus, ut in
nostro filiformes sunt.

Pag. 54. *Adenostyles alpina* Cass. ortus et in monte Javor Drago ditiois Spalatensis.

Pag. 185 Post *Mentham sylvestrem* L. addatur

Mentha pubescens W. en. h. bot. II p. 608. Rehb. fl. ex. p. 308.

*M. foliis petiolatis ovatis pubescentibus subhis caespitibus, spicis oblongo-
cylindricis inferne internodiis, calycibus villosis - caespitibus.*

Syn. *M. nepetoides* Lej. rev. de la fl. de Spa p. 116. *M. aquatica-
sylvestris* Mey. chlor. bot. p. 289. — Icon. Rehb. ii. bot. X. p. 1309.

Hab. in agris ^{et feniis} circa Narvuta, unde missi ex. Matth. Boten.

Fl. aestate. Fl. lilacina. 4.

Obs. Planta probabiliter hybrida ex *M. aquatica* quam refert
Fluc, et *M. sylvestris* cujus habet floris spicatis.

Page 229. lin. 31. loco peduncul. ramoso affinis lege peduncul. junis affinis.

Page 241. Ad Cynoglossum chinensis Ten. ^{reclut. de} ~~physogonum~~ ad C. pictum Ait. partes
C. pictum Alsch. fl. par. p. 48 ex ipsius specimen.

Page 250. lin. 40 loco ubi absunt lege ubi adsunt.

Page 262. lin. 12 loco antecedent corr. antecedent.

Page 11. Asperulae Mahiane Ord. VII. III. Flor. detm. III. 12.

Page 69. Laricidium verticillatum Walst. et Kit. videtur idem ac Physogonum
actaeifolium Presl. det. prae. p. 128. Differentiae loc. cit. adductae inter
Physogonum et Alschingera ex capsularum forma, pygmae crassitie et in tra-
rum latitudine, ^{minoris pichi miki} ~~minima constant~~ vixae sunt post accuratam comparisonem
fructuum Physogoni ^{actaeifolii} ~~verticillati~~ cum illis Alschingerae nec tanti facienda
ut ipsi novum genus repositui possit. Planta hae igitur jure prioribus
namque speciei Kitaibelloni in postum vocandae erit Physogonum
verticillatum - Characteres perperam generis, aliquantisper reformandus ^{part}
juxta illum Alschingerae, cujus fructus ^{quod nos} ~~maximam~~ planissimum aperi-
unt pygmae ^{adco} ~~evoluti~~ ~~fructus~~ praebuerunt, ut ab illis Physogonis
descriptis satis distinctae viderentur -

Page 93. Post Ranunculaceae inseratur Legum. Ord.

Ordo LXXI. bis Droseraceae

D.C. theor. elem. p. 214.

Floris hermaphroditi regulari. calyx lobis pentagonalibus aestivatione imbricatis, sepalis
ima bafi ischaeranthibus. corolla petalis quinque hypogynis. Stamina quinque hyp-
gyna, antheris inrosis bilocularibus, vixae longitudinali duplici delipentibus.
Pistillum ovario libero uniloculari planicollato placenta 3-5 parietalibus, stylo
3-5-partito, stigmatibus integris emarginatis. capsula uniloculari, bivalve
deliscens tri-quinquevalvis, seminibus plurimis ~~testa reticulata~~ ^{laevissima},
cavofo involufo, radiula ad hilum vixae, embryone ~~pari recto~~ ^{abnormi}

Hecinae palustres foliis alternis glandulosis pilosis, junonibus circunatis, floribus saepe
vacuosis subnatis.

Drosera L.

Gen. pl. ed. II. p. 108.

Char. formae ac ordinis

Drosera longifolia L. sp. pl. ed. II. p. 501 ex part. Hort. fl. austr. I. p. 415. Rob. fl. europ. 111

L. foliis cuneatis oblongis in petiolum attenuatis, seminibus oblongis testa reticulata laevissima
involuto

Icon. Cons. et Germ. fl. par. atl. tab. IX. f. 3-5. opt. Ved. fl. dan. t. 1093. Bull. herb. d. la Tr. t. 181.
Hab. in paludibus circa Narvica ubi legit Dr. Neesmaeyer. Fl. alb. aut rosei. Fl. aestate. 2.

Pag. 101. In Clavis analytica generum loco Fuchsia sequitur complanata Vis.
Nr. 1. lye Vis. Nr. 5.

Pag. 135 Brasica Bottemi Vis. adda iconem Hort. Delm. LII p. 1. et post
adda sequentem.

Brasica mollis Vis.

B. superbula velutino-canescent, folijs obovatis, inferioribus obovatis lyratis inac-
liter grossiusque crenato-lobatis petiolatis, superioribus lanceolatis ligulatis inte-
gris semialexicaulibus, calyce laxo, siliquis tetragonis crenatis polypermis,
style conico subulato apertus.

Hab. ad rupes scopulorum innulea Carzola circumstantium, ubi legit agr.

Math. Bottem. fl. luteo. fl. Mayo. 5

Deser. caulis teretifolius stratiss. superioribus inferioribus mammillatis,
media et superioribus lappis acumulatis. Siliqua 2-foveolatus ad quatuor
decim lineas longa, pedicello sua duplo longiore longior, tetragona
lobatis superioribus concoloribus, style lineam longa subulato, inferioribus aut
longioribus concoloribus in superioribus locis 12-16 subobtusis perispermis. — Apertus Brasica
Bottemi Vis., a qua probè differt siliqua tetragona 12-16-perispermis, style
subulato valeris plus duplo angustioris, dem in illa siliqua concoloribus bravibus
subhexapermis, et style ovatis siliquis latitudinem aequat.

Pag. 136 Post Brasica Ragam (incanthes

Brasica adpressa Vis.

B. caulis polys ovoidibus superioribus linearibus lanceolatis, calycibus patulis,
siliquis vix adpressis tumulis teretibus vix oblongis seminiferis, style bre-
vis, stigmate emarginato.

Syn. Sinapis incana L. sp. pl. ed. II. p. 934. Myagrum hispanicum L. p. pl. ed. II. p.
893. Hirschfeldia adpressa Moench meth. fl. p. 264. Cordylacarpus puckers-
im. fl. gr. prodr. II. p. 33. — Leon. Reich. II. fl. germ. II. p. 4423. Lucq.
herb. ind. II. p. 169. Herm. prodr. bot. p. 155. f. 39. Bonum. panph. III. p. 50.

Hab. ad rupes apertas innulea Carzola unde visit Al. Salis. fl. luteo. fl. Mayo. 5.

Pag. 146. Helianthemum montanum Vis. mutetur in H. Rhodae Heud. nom. bot. II. p. 735.

quid nomen anterioris.

Pag. 163. Lianthus viridescens Vis. Addatur icon fl. delm. tab. LIII p. 2

Pag. 166. Silene remotiflora Vis. Addatur icon fl. delm. tab. LIII p. 1