


~~13~~

Bolbitius



111

(Bolting)

Conspectus synopticus specimen

A₁ - Stipes albo-squamosus; pileus vitellinus 1 B. vitellinus
A₂ - Stipes glaber.

B₁ - Pileus lviso argenteo-albus; lamellae
denum ferrugineae 2 B. conocephalus
B₂ - Pileus albo-coloratus; lamellae non
ferrugineae.

C₁ - Stipes albus 3 B. tener

C₂ - Stipes flavus hitecens
D₁ - Lamellae e purpura
luteo fuscæ 4 B. leucanthus

D₂ - Lamellæ e luteo
uninamomeæ 5 B. fragilis

Bolting

Bolbitius

~~Gomphidioides~~

Fr. Epicr. pg. 253

Sacc. Syll. Fung. I pg. 1073

(inter Melano-sporas)

(*Etym.* ~~bolbitos~~, ^{bolbiton} firmus)

Hymenophorum subdiscretum. Volum universale nullum, partiale in plurimis obsoletum. Lamellae membranaceae, molles, liquecentes (nec diffluentes) e sporis secedentibus pulverulentae. Sporae ovatae, leves, subferrugineae.

(— Fungi mucidi, nudi, brevi marcescentes (nec reviviscentes), plurimi fumicolae vel locis stereoratis vi- gentes, lutescentes.

Oss. Piccoli greggi molto naturale, del tipo *Coprinus* per la vegetazione, *Cortinarius* per la fruttificazione, quindi intermedio fra queste due e strettamente appartenente alle Boletacee. Le sue specie non sono né commestibili, né velenose.

1 *Bolbitis wellingii* (Pers.) Fr. Epior. p. 254 (1836) Sav. Sylt. fung
IV p. 1074, *Agaricus wellingii* Pers. Syn. p. 402 (1801)
Zon. Gr. A. Champ. Fr. t. suppl., Cooke M. Br. F. t. 928
Bitt. Bres. MS., 1663. ~~key~~
Hab. ad terram detritum v. stercoretam — Trentino: Magra et supra
Ornago (Bres. MS.), Warne (v. Höhn.). Aestate

Vitellinus 1074

Pileo carnoso-membranaceo, ex ovato esplanato, viscido, vitellino, primo levi, dein
marginie fiso-sulcato, 4-6 cm. lato, refran-
do, unicolore;
lamellis leviter adnexis, ochraceo-argillaceis;

Lipite fistulato, aequali, alto-guanoso, 5-8
cm. longo, 4-7 mm. crasso;
~~Sporif subobtuseis~~ 13-14-8, ochraceo-
ferrugineis vel flavo-ochraceis, subamygdale
liformibus, apice truncatis 12-15=7-9 (Mrs. M.S.)


I.H. Quasi aspettico, maggiore di *B. fragilis*, *bulbillus* ec. Ha tutto
il portamento di *Coprinus micaceus*
(Bref. N.S.)

~~Aestuare~~



2 *Bolbitius conocephalus* (Bull.) Fr. Sp. in p. 205-
Jan. J.M. Zang. V, 1075, *Agaricus conocephalus* Bull. Champ t. 563 f 1
Type Bull s.c. Cooke M. Br. F. t. 1160, Hoffm. Sc. anal. t. ~~105~~ 7 f 2
gr. H. et Champ. Fr. t. 383

Bull. 90, 397, 398, 417, 418, 585-



Hab. ad terram humosam — Liguria, Piemonte, Lombardia, Trentino. Autunno

Exs. Klot. Herb. myc. 1901 (Vercelli: Cesati)

conocephalus 1075

Limpida, gracilis, fragilis e livido
argillaceo-albus;
Pileo membranaceo, conico, hygroscopico,
disco lev, vistidulo, margine striato, 1 cm. latu;

lamellis liberis, ventricosis, e supercente
ferrugineis;

stipite fistuloso, aequali, glabro, nitido,
senacello albo; spona oblongo-transformata
10 p longa, cremeo-ochracea. —

Oss. *Leucostoma* *Polyphaena saginata*

2 Bolbitis Lener Berk. Bull. p. 183, t. 12 f. 2 (1860), Van. Syst. Fung

I p. 1076. Berk. P.c.

Icon. Fr. Ion. Hymen. t. 130 f. 4, Cooke M Br. F. t. 691.

Bull. Presad. Ms.

Hab. in pratis subhamata, pinguis - Trentino: sopra Ormea (Pres. Ms.)
area dist. Europa

aerata

tener 1076

Pileo tenerino, elongato-conico, liso, nudo,
albo, ex incarnato albicans, vel albido-lu-
scolo;
lamellae liberas, postice attenuatas, rubellij;

stipe fusculo, albo, lassi leviter
incrassato, ^{non distincte bulboso,} 5 cm. usque longo

sporif 10 - 12 = 5 - 8; ex presad. MS.
ovato-ellipsoideis, sub microscopicis flavo-
ochraceis, ~~11-14~~ = 8 - 10; basidiis

21 - 24 = 10 - 12

Restato



Dolichus titubans (Bull.) Fr. Spec. p. 256 (1836)

(1182)

Jan. Synt. fung. IV p. 1075, Agaricus titubans Bull. Champ. t. 425 (1)

Non. Bull. p.c., Cooke W. M. F. t. 690, Pitt. Champ. F. t. 380, Patouillard

Tab. anal. fig. 657. Sov. Fig. t. 128.

Bull. 90, 585, Pres. M.S. 6 Treat. no: Megras at prope Massa Bellunum Trento (Pres. M.S.)

Hab. in elevatis ^{herbosis vel} ad finium → Liguria

area sicc. Europe, Australia.

(Aeratae)

titubans 1075

Fragilissimus, tremulo titubans;

Plo membranaceo, ovato-campanulato,

dein fuso explanato, flavo, discoideo, 2,5 cm.
lato; lamellis leviter leviter adnexis, e pale
lide purpurascente fasciis (postea quam obscur-
eas ut habetur in Bull.)

Spirite graciis, stricto, luscenti, nitido,
11-14 cm. longo, 2-4 mm. crasso.

sporij incarnato-salmonis 8-9-4-5;


ex Bro. M.S. sporis ferrugineo-rufescensibus,
ut microscopio. flavis, ovoides 12-13 = 8-9;
et etiam subracies, obovato-elongatis vel
subamygdaliformibus, apice truncatis, 11-15 = 7-8,5;
basiis 22-24 = 10-11 / - <

(Bro. M.S)

021. Presunto in M.S. ^{sospette la sua identità} ~~grave con Galera lateritia?~~

Asperula

5 Bolbitius fragilis (Linn.) Fr. Sp. or. p. 256 (1836)
See Syl. fung. V p. 1074, Agaricus fragilis Linn. Sp. pl. ed I p. 1185 (1753)
Trem. Bolt. Fung. t. 65, Cooke M. Br. F. t. 720 f. A., Hoffm. Icon. myc.
t. 21 f. 2.

Rob. 43° 866, 1062, 1161  Piemonte

Hab. in campus et hookis — (Toscana, Emilia, Lazio, Napoletano. Vere
area sicc. Europa, Apennina aust. Australia.

fragilis 107H

Fragilis, marefrey, e flavo expallens;
Pileo submembranaceo, viscido, peltato
nico, margine strato, disco subumbonato,
1-2 cm. lato;
lamellis attenuato-adnexis, e buccolo
dilute cinnamomeis;
stipe attenuato, nudo, glabro, lutes, 8
cm. longo.
sporif ferrugineis (sunt lente flavif
vel flavidif), subspphaeroideo-ellipsoi
des 14-15 = 8-9 - ↙

DH. Più solite di Bolt. Boltiorii