

Melanomma

Sphaeria Ericae Fr. S.M. Ezech. 2, p. 101. Sparsa

erumpens, peritheciis globosis purpureis fuscis, ostiolo
conico papillato; sporis leviter fuscidulis, 3-septatis,
demum vix contractis, oblongis, 13-16 x 4-5, monostichis
in ascis cylindraceis, 90 x 6. - In ramis Ericae vulgaris
Brioune. (Euv).

Physalospora Sphaeria Solidaginis & S.M. El. 2, p. 106. Subimata

orbiculari Depressa, fuliginosa atra, disco erumpente albo, papilla
centrali atra umbilicata. Ascis cylindraceo-clavulatis,
paraphysatis, 7 x 8, sporis hyalinis ^{oblique} monostichis ovalibus 2 guttulatis
14 x 7. Perithecia per epidermidem tegumenta translucentia.
In caulis emortuis Solidaginis; Brioune (Euv)

Sphaeria rosicola Kehl. Symb. M. 114. In Rosacan.

Rouen ?

Sphaeria persistens Cooke H.B. 888. In Rose canina

Rouen. an Didymella ?

Clypeosporia Notarisii Felt. l.c. 117. In Rubus.

Rouen. Sporis monostichis, elliptico-fusiformis rectis
v. curvulis, brunneis, 3-septatis 20-23 x 5-6. (passim
4-5-septatis) An Sporia maritima ?

Rhaphidophora Sur Sieraria vulgaris

Ceci 170. Sp. 100-120 d. lutescentes.

Prothemiella trifida Mall. Conceptaculis 2 parvis
convexiusculis pallidis-lutescentibus; Conidiis aggregato-

3-stellatis, obclavulatis septatis, torulosis, hyalinis,
30-35 x 3, cellula centrali elliptica, alio ad apicem
attenuatis. In foliis scabiosa succisa; vere. Brioune

emortuis }

(Euv)

Sphaeria pruri Socia

Asc. 58 x 14

Sp. 25-30 x 4-6, 5

CERCLE PRATIQUE

D'HORTICULTURE

et de

BOTANIQUE

du

Département

de la

Seine-Inférieure.

Reuen, le

184

Monsieur