

Peziza araneosa ^{on Hook.} Berk. *H. Tasm.* II p. 275;
cupulis primum subglobosis demum ex hemisphaerico expansis fissilibus.
inter araneosis floccis pubes super matricem repentes orundis, hymenio luteo-
aurantiaco, sporidiis curculis multiseptatis. Hab. On dead wood, Archer.

Minute, at first globose then hemispherical, with border inflexed, then ex-
panded, white and lycopid externally, attached to the matrix by radiating
lycopid flocci. Hymenium pale-orange. Asci clavate. Sporidia linear,
curved, containing six or more nuclei, $\frac{1}{500}$ inch long.

Peziza Archeri Berk. n. Hook. *H. Tasm.* II p. 274 f. 18 f. 72;

cupulis, innatarina, primum concava, demum expansa,
undulata, margine libero.

Hab. On dead leaves of some succulent plant, Archer.
Bright crimson. Cup $\frac{1}{4}$ inch across, sessile, at first depressed, then expanded,
undulatus and umbilicate, paler below, slightly tomentose. Asci cylindrical.
Sporidia globose, $\frac{1}{3500}$ inch across, with a large nucleus. - The fruit is just like
that of *P. endocarpeoides* B. The species is almost intermediate between the sections *Humaria* and
Hellisia. The sporidia ~~are~~ very rarely in pairs in the same ascus.

NB. Sul lato anteriore si scrive soltanto l'indirizzo.

CARTOLINA POSTALE

RISPOSTA



NB. Questa parte può essere spedita isolata.

Al prof. P. A. Suardo

Art. botanico

Padova

(Provincia di _____)