

11/ Lelidium placophyllum (Sng.) Rehm

Synon: Abrochallus placophyllus Sng. / Anal. lich. rar. p. 26

Thallus alienus, apothecia minuta, planiuscula, immarginata, imbricata, tandem maculariformiter confluentia, atra, Epitheliumque fusco, paraphysibus crassis subcohaerentibus. Sporae in acis cylindricis vel clavatis 6, saepe 1 seriatas, oblongae, 2 loculares, pellucidae, $8-9/3-4 \mu$. Gelatina hymenialis 2 distincte coeruleis.

En thallo Lophyridii placophylli. Alpi Gavia (Comoria) Ital. 209.
/ Simulat Ruelliam scabram.

12/ Karschia homoclinella (Nyl.) Rehm

Synon: Lecidea homoclinella Nyl. / Flora 1872 p. 261

Apothecia parasitica parva, c. 9, 3 mult., opaca, demum planiuscula, immarginata, juniora margine subobsoleta. Sporae 8, ellipsoideae vel oblongo-ellipsoideae, 1 septatae, $7-11/3-4 \mu$, fuscae. Paraphyses molles, subarticulatae, apice lutescenti-fuscae, inde Epithelium luteo-fuscescens. Hypothecium non crassum, infuscescens. Gelatina hymenialis vinosae rubens vel fulvescens praecedente coalescentia levi.

Supra Lecanoram atryneam Cetrulicolum. Fens.

13/ Karschia adjuncta (Th. Fr.) Rehm

Synon Ruellia adjuncta Th. Fr. / Prot. nos. 1866 p. 14-18

Ep. Trompeltuber / Flora 1866 p. 266

Thallo nullo, apotheciis minutis, 8, 3-8, $4 \frac{1}{2}$ diam., adnatis, concavae marginatisque, variis demum planiusculis immarginatisque, atris, nudis, hypothecio sordido, paraphysibus laevibus cohaerentibus, nigro capitatis; acis inflato clavatis, gelatina hymenialis 2 + sporis 8, ellipsoideis, utrinque obtusis, $13-16/7-8 \mu$.

Parasitica in thallo Placodi stromminei. Puccia.

- 20) Arthonia circinata Th. Fr. (Bot. not. 1865 p. 100)
 cf. Almqvist Mon. Arthon. in Vet. Sk. Handl. 1880 p. 56
 subparasitica in Gyrophora
- 21) Arthonia caerulea Almq. (Mon. Arthon. p. 59)
 In Apothecia Lecanora variae sp. corticolae. Luccia.

- 23) Scabula epizema (Nyl.) Melius IX
 Synon: Lecidea epizema Nyl. (Prod. Lich. p. 125)
 cf. Lamy (Bull. soc. bot. Fr. 1863 LXX p. 428, Richard (Lich. Deux Leves 1878
 p. 11. Annot. Flor. 1874 p. 97), Exl. Lich. 1194.
 Apothecia 5-12 congregata, raritate sparsa, c. 0,5^m diam., primivus urceolatis,
 distincte marginata, demum plera, demum convexa et imarginata, nigra.
 Asci ovato-pyriformes, 35/10 μ (porae 8, ellipsoideae vel elongatae
 oblongae, 1 septatae, hyalinae, 10-18/4-5 μ . Paraphyses obiter
 apice dilatatae et subfuscae. Hypothecium fuscum.
 In thallo Lecanora calcareae et Verrucariae nigrescentis. Gallia.

19, Leciographa plumbea (Anzi) (Manip. p. 29)

Thallus proprius nullus. Apothecia parasitica, parvula,
sessilia, atra, solitaria vel aggregata, plana, margine ter-
decim convexa et immarginata; lamina lichenina praedisa, su-
perne viridula, paraphysibus conglutinatis, hypothecio fusco
Sporae in ascis clavatis 8, anguste fusiformes, 4 circulares
pellucidae, 20/34. Hymenium 1+.

In thallo Panariae plumbeae ad ramos Rearum. Italia.

(Asci 45-50/10-12 μ , sporae 18-24/4-5 μ . Paraphyses apice subqu-
boae, viridulae - 6 μ st.)

8. Tryblidaria nivalis (Raggl. et Car.) Pellus

Synon: Leciographa nivalis Raggl. et Car. Annot. Ach. Val. p. 215 tab. 4 (58)
Dactylopora nivalis Arnold / Ledb. Fragm. XV in Flora 1874
Ep. Arnold flora 1881 p. 324

Prothecia in thallo alieno primum innata pumeliformia, cito e-
merso sephila, vulgo rotundata, convexa, primum alba conspersa, demum
nuda, atra, margine tenui mox evanescente. Asci ellipsoideo ventriculosi,
6-8 spori, paraphysibus validioribus apicem versus fusciscentibus ob-
vallati. Spores ellipsoideae obovataeve aut subrotundae, fusciscentes, 2-
ut plurimum 4 loculares, non raro septis verticalibus intercom divisa, 6-10
loculares, ad dissepimenta quandoque leviter emarginatae, 15-20/9-10 μ .
In thallo Amphitomatia elegantia. Ospicio di Valdobbia, Dale 1881

26 Orbilia tenuifirma Pat. (Bull. soci. myc. IV 3 p. 99) non Speg.
(Sacc. syl. VIII p. 625)

Apothecia gregaria, sessilia, orbicularia, patellaria vel cupuliformia,
0,5-1^m, pellucida, flavoaurantiaca, sicca rubro-nigra, margine integro
vel undulato. Asci clavati, 30/1-3 μ , 8 spori. Spores subglobosae, hy-
alinae, 0,25-0,5 μ , ibiduae. Paraphyses filiformes, apice globulosae,
hyalinae.

Ad lignum emortuum. Venezuela.

= O. Gardard Sacc. VIII

Apothecia saepe arcte connata, striae longae in ligno formantia. h. 629

25 Thurmeria olivacea (Cuiet in lit.) Cooke / Mycogr. p. 231 pl. 109 f. 389 (1879)
Syn.: Alenia olivacea Cuiet / Endvir. f. p. 278
non Bond. (1897)

~~Aspergillus~~ ~~sepioides~~ ~~applanatus~~, orbiculatus vel flexuosus, lacteus
olivaceus, margine fere niteo elevato, 8-10^{um} diam.; ~~asci~~ cylindrici;
Sporidia elliptica, utrinque subattenuata, nucleata levib., hyalina
2.5-10-12, 8, monokiliis; paraphysis supra incrassata, oli-
vacea limbo.

(Hb. S. terramⁱⁿ Gallia olivacea
(Diversa videtur Peziza Cuiet / Bull. soc. bot. fr. 1878 XXV p. 29
apothecia 1^{um} diam.)

1) Aulographum quercinum Ell. et Mart. (An. nat. Decb 1883
p. 1282)

Ex Ell. et Ev. N. Am. p. 78.

Ex: Ellis N. Am. f. 1284.

Ad folia Quercus virens Florida.

Descript. in Sac. 24. 1

Lomboni pyrenaeica Ell. et Mart.
XIV

3) Gloniella pyrenaeica Petrus (Rev. hb. Stry

in Hedwigia 1886. p. 34)

Sporothecia in maculis ligni denigratis immersa,
sparsa, necata, 2-3 μ long., $\frac{1}{4}$ μ lat. Asci clavati 70/15 μ ,
8 spori. Sporae fusiformes, subacutae et subcurvatae,
hyalinae, 4 cellulares, - 18/5 μ , distichae. Paraphyses
ramosae, fasciculatae. ?

Ad lignum Loniceræ Periclymeni St. Antoine de
Galamus. Pyren. ex.

IX h 1116

27 / Phaeospora arachnoidea (Speg. et Lac.) Peck.

Lyn: Peziza arachnoidea Speg. et Lac. / Rizzoz. Myc. Ven. I p. 223

Apothecia gregaria, sessilia, sicca patellariformiter explanata,
humida 2 fenoides, pallide aurantiaca, encipulo pallide pilis,
setis filiformibus, i cellularibus, 110-120/8 μ . Paraphyses filis
formis, ad apicem clavulatae. Sporae 2 stidacum superiore acie parte,
globosae, gelabrae, 15/4-15 μ . = Pyrenopeziza
aranea (Speg.) Sacc.

In stercore humano et canino. Italia.

f. canina: aeri 100-120/24-26 μ , sporae 11-12/10 μ .

34 Verpa speciosa Vittad. / Mang. p. 120 tab. 15 (16)

Stipes conico campanulatus, subobtusus, plicatus, superne
costatus, alveolatus, fuscus, subtus planus et floccosus, etc.
pes latus, albidus, floccosus, intus fibris tenuissimis replens.

cf. Grevillea p. 57 =

Nothella bohemica R. K.
gigas Fr.

15/ Leciographa conveca Th. Fr. | Reum

Cyni Puellia conveca Th. Fr. | Len. arct. p. 234, Gih. Spitzg. p. 45
Thallus alienus. Apothecia adnata vel adpressa, parvula, disco prae-
sulis subplano et margine crassiusculo cincto, non convexo margi-
nemque excludente, atro, nudo. Asci inflato-clavati vel venter-
covi, paraphysibus caesae cohaerentibus, apice fusciscenti capitulo
cincti. Sporae 8, oblongae, saepe curvulae, 4 blastae, septo uno al-
terove longitudine nunquam divisa, fuliginose fuscae, 13-24
6-9 μ . Hypothecium nigricanti fusciscent. Gelatina hymenium
9+.

Supra thallum. Physciae caesiae saxicolae, et obscurae, Hanshorstae
elegantis et Acetabularia cheilophacae. Luecia.

16/ Leciographa Weissii Körber | Zool. bot. Verh. 1867 p. 207.

Thallus alienus. Apothecia enata, a thallo quandoque coro-
nata, alevina, opaca, inmarginata, difformia, psellariae seu
cephaloidea, saepius lineatim sulcata. Sporae in ascis cylindra-
cis 1 serialiter 6-8, mediocres, tandem majusculae, dacty-
loideae, 4 blastae, diametro 4-6 plo longiores, obscure fuscae.
Parasitica in thallo Pertusariae et Cubrolechiae. Dalmatia.

17 Lecanographa furfuracea (Anzi) Pellum

Synon.: Elidium furfuraceum Anzi / Cat. Lich. Londr.
Eus. Anzi Lich. var. Langob. 249.

Thallus extraneus. Apothecia lichenina destituta, puncti-
minutissima, basi saepe a thallo extraneo marginata, abra-
vaco pulverulenta, planimucosa et marginata, vel convexa ex-
ginata, sporae in ascis cylindricis-clavatis 8, exsertus
lineae, subseriatae, oblongae, 4 localiter, 11, 2 - 13 / 5, 6

Ad thallum Lecidae nivalis, Rhizocarpi subincubici, Le-
scoidae in alpinis Vallis Tellinae

(² syn: Elidium incitivum Korb.)

Pellum

Indicaco. Thallo cylindrico. Anzi.
Anzi. Thallo Tellinae.

Retin

F. calmullescens, Dein arava
Supra lignum veduesam

Fenia
Wohl 29n

Donkotrema Lindl. Powell

Handwritten notes at the top edge of the paper, partially obscured.

Handwritten notes at the top edge of the paper, partially obscured.

6/ Odontotrema anodontum Nyl. (Flora 1869 p. 411)

Simile fere *O. subintegro* Nyl., sed dignoscitur jam sporis
ellipsoideis, 3-5 septatis (denuaque accedente septulo rathum
longitudinali) majoribus, 14-15/5-8 μ . ~~Ab~~ Suci apice Todi
opre coeruleo, cum gelatina hymeniali.

Ad lignum Juniperi. Lappland.

7/ Odontotrema phacidiellum Nyl. (Flora 1874 p. 316)

Simile *O. phacidoi* Nyl., sed minus, sporis 1 septatis, paullo
senioribus, 11-15/2,5-3,5 μ . Epithecium incolore. Gelatina hymenial
I. caeruleo, dein vinose fulvescens. mit inosität)

Supra lignum vesucum Fentia (wohl 29n) Odontotrema Dini Monell