

Filices Farvisinae.

- cum Selaginibus, Hydropteridibus atque Calamariis.
- | | | | |
|-----|---------------------------------------|----|--|
| 1 | <i>Pteris aquilina</i> Linn. | 23 | <i>Equisetum arvense</i> Linn. |
| 2 | <i>Blechnum Spicant</i> Smith. | 24 | " <i>fluviatile</i> Linn. |
| 3 | <i>Scelopendrium officinale</i> Sm. | 25 | " <i>hyemale</i> Linn. |
| 4 | <i>Asplenium Trichomanes</i> L. | 26 | " <i>palustre</i> Linn. |
| 5 | " <i>Puta muraria</i> L. | 34 | " <i>silvaticum</i> L. |
| 6 | " <i>Adiantum nigrum</i> L. | 46 | " <i>ramosum</i> Schlecht. |
| 7 | " <i>acutum</i> Bory. | 41 | " <i>variegatum</i> Schlecht. |
| 8 | <i>Attherium Filix femina</i> Roth. | | |
| 9 | <i>Aspidium fragile</i> Sw. | 27 | <i>Chara vulgaris</i> Linn. |
| | (<i>Cystopteris fragilis</i> Bernh.) | 35 | " <i>flexilis</i> L. (<i>mucomata</i> A.B.) |
| 10 | <i>Polystichum aculeatum</i> Roth. | 36 | " <i>opaca</i> Ag. (<i>midacea</i> Mill.) |
| 11 | " <i>Filix mas</i> DC. | | |
| 12 | " <i>Thelypteris</i> Roth. | | |
| 13 | <i>Polypodium calcareum</i> Smith. | | |
| | (<i>Clus. hist. II p. 212</i>) | | |
| 14 | " <i>vulgare</i> Linn. | 31 | <i>Aspidium montanum</i> Sw. |
| 15 | " <i>Thaectopteris</i> Linn. | 37 | " <i>Oreopteris</i> Sw. |
| 16 | <i>Ceterach officinarum</i> Baub. | 38 | <i>Asplenium vire</i> Sw. |
| 17 | <i>Adiantum Caput Veneris</i> L. | 39 | — <i>frustum</i> Walld. K.T. |
| 28 | <i>Asplenium Breynii</i> Swartz. | | |
| 6/6 | " <i>Puta muraria</i> L. | | |
| | — var. <i>trijunata</i> | | |
| 29 | <i>Botrychium ternaria</i> Sw. | | |
| 30 | <i>Polystichum dilatatum</i> DC. | | |
| | — var. <i>cyuapifolium</i> | | |
| | (<i>Polypod. Hoffm.</i>) | | |
| 10 | " <i>Londensis</i> | | |
| 18 | <i>Lycopodium clavatum</i> L. | | |
| 19 | " <i>helveticum</i> Linn. | | |
| 20 | " <i>complanatum</i> L. | | |
| 32 | " <i>Selaginoides</i> L. | | |
| 33 | " <i>Selago</i> L. | | |
| 42 | " <i>mundana</i> | | |
| 21 | <i>Martilea quadifolia</i> L. | | |
| 22 | <i>Salvinia natans</i> All. | | |

In Filicibus, huiusque affinitibus plantis Observationes.

- 1 Vulgo: Speltea. In s. Martello vulgarissima, nec deest in plantis repitibus, locisque dimetatis. Sa utuntur vltis ad stramen totus, equis aliisq. animalibus parand₂ dum.
- 2 In sylva Martello rarum admodum. Habet ex valle ejusdem Dell'Agua dicta.
- 3 In sylvae Martelli umbrosis, tenosis, humidisq. Nascentur etiam in humilioris provinciae puteis et muris vetustis. Vulgo dicitur: Lengue et serps. In sylva Martello legi varietatem crispam, q_{ue} rara.
- 4 In murorum veterum vitiis frequentissimum.
- 5 In muris vetustis, rupibus commune.
- 6 In quercuum caudicibus frequens praesertim in sylva Martello.
- 7 Legi in truncis quercus prope et forane de Giavara in sylva Martello. haec frequens.
- 8 In uliginosis Martelli, collium et montium vulg₂ lum: vulgo Speldin.
- 9 Legi in ruderat_{is} humentibus alle vat delle Fontana a Cornuda.
- 10 In umbrosis Martellianis frequens: in collibus omnibus.
- 11 In Martello et dimetis collium et montanis comm₂ ne; frequens etiam ad sepes humidas plantis.
- 12 Legi ad fanas Bavaria albae dictas: sed est infrequens: in Sagnis sylva Martello, et Aquino.
- 13 In lapidosis ^{calcareis} a Corvato ^{Corvato} Dryopteris Linn, cui ster₂ pen meam antea referebam, huius simillimum est et auctores plerique cum hoc confundunt.
- 14 Vulgo: Pimpinimpolis, pimpinimpolis. Rhizoma com₂ dicitur a pueris, quia subdole.
- 15 In Martelli collibus umbrosis frequens.
- 16 Legi ad rupium fessuras prope Serravalle, in inferior₂ provincia haud frequens.
- 17 In specubus humentibus Martelli, ^{Corvato} Co. Goro etc: inter₂ dum in puteis.
- 18 In collium Martelli graminoso et macro caumine frequens reperitur; nec non in montibus fere omnibus. Dicitur vulgo: Erba pedocchina, quia monticulae herbae decocto et pulvere utuntur ad pediculas enecandas.
- 19 In rupium aqua humentium prope ad Plain₂ frequens.
- 20 Cum Lycopodi clavato frequens.

48	.00
6	.00
150	.00
56	.00
26	.00
108	.00
36	.00
200	.00
286	.90
628	.43
102	.50
86	.00
1727	.80

1730
500
1230
60
1170
205
965

60	88
10	295

Muri. Kensterni. exsiccati. decur.

+ <i>Bombacum</i> <i>indifiatum</i>		<i>Hypnum</i> <i>fluitans</i>	
<i>Bryum</i> <i>bimum</i> 1		" <i>Kneiffi</i>	
" <i>gallienum</i> 2		" <i>commutatum</i> 17	
+ " <i>torquens</i>		" <i>pilicinum</i>	
" <i>capillare</i> 47		+ " <i>intermedium</i>	
<i>Brachythecium</i> <i>salicifolium</i> 3		" <i>univittatum</i> 18	
<i>Barbula</i> <i>dederingii</i> 4		" <i>Halleri</i> 19	
" <i>Halleriana</i> 5		" <i>Schreberi</i> 20	
<i>Barbula</i> <i>tortuosa</i> 6		" <i>vicinatum</i> 21	
<i>Antitrichia</i> <i>contropendula</i> 7		+ " <i>cardifolium</i>	
+ <i>Monodon</i> <i>distans</i> 48		" <i>trisetum</i>	
<i>Amblystegium</i> <i>irriguum</i> 49		" <i>brevistre</i> 22	
<i>Fornelii</i> <i>aut</i> <i>piperitica</i> 49		+ <i>Hornelia</i> <i>schumanniana</i> 23	
<i>Xichodontium</i> <i>johannicum</i> 8		<i>Hornelothecium</i> <i>Phylloporum</i> 24	
<i>Eurhynchium</i> <i>cinnatum</i> 9		<i>Mnium</i> <i>punctatum</i> 25	
" <i>striatum</i> 46		" <i>orthorrhynchum</i> 25	
<i>Xiracella</i> <i>heteromalla</i> 10		<i>Nacken</i> <i>pumila</i> 26	
<i>Xiracium</i> <i>repens</i> 11		<i>Orthotrichum</i> <i>stramineum</i> 27	
" <i>undulatum</i> 12		" <i>speciosum</i> 28	
" <i>Sauteri</i> 13		<i>Leptotrichum</i> <i>glaucescens</i> 29	
<i>Xiracodontium</i> <i>longistria</i>		" <i>flexicaule</i> 30	
<i>X-dymondii</i> <i>rubellum</i> 14		<i>Philotrichum</i> <i>maritimum</i> 31	
<i>Distichum</i> <i>capillare</i> 15		<i>Plagithecium</i> <i>pleurocarum</i> 32	
+ <i>Cylindrothecium</i> <i>cinnamum</i> 56		<i>Pogonotrichum</i> <i>nanum</i> 33	
<i>Ceratodon</i> <i>purpureus</i> 16		<i>Polytrichum</i> <i>commune</i> 34	
		" <i>formosum</i> 35	

- Pteris gymnantha filiformis* 36
Phlox Ciste-castensis 37
Rhynchospora dep. sp.
Sphagnum cymbifolium 38
 " *acutifolium* 39
Cystopteris crispus 40
Tetraphis pellucida 41
Trichostema anacardium 42
 " *crispulum* 43
Licor julace 44
Barbula fragilis
Anomodon attenuatus stenellus 51
Amblystegium chrysophyllum ster. 52
 " *uncinatum fructif.* — 53